

Question			Expected Answers	Marks	Additional Guidance
1	a	i	^{118}Sn 50p 68n 50e Complete row ✓	1	
		ii	$^{120}_{50}\text{Sn}$ has (two) more neutrons / 70 neutrons ✓ ora	1	ALLOW There is a different number of neutrons IGNORE correct reference to protons / electrons DO NOT ALLOW incorrect references to protons / electrons ALLOW ECF for stated number of neutrons from 1a(i)
	b	i	The (weighted) mean mass of an atom (of an element) OR The (weighted) average mass of an atom (of an element) ✓ compared with 1/12th (the mass) ✓ of (one atom of) carbon-12 ✓	3	ALLOW average atomic mass DO NOT ALLOW mean mass of an element ALLOW mean mass of isotopes OR average mass of isotopes DO NOT ALLOW the singular; 'isotope' For second and third marking points ALLOW compared with (the mass of) carbon-12 which is 12 ALLOW mass of one mole of atoms ✓ compared to 1/12th ✓ (mass of) one mole OR 12g of carbon-12 ✓ ALLOW $\frac{\text{mass of one mole of atoms}}{1/12\text{th mass of one mole OR 12g of carbon-12}}$
	c		moles of Sn = $\frac{2080}{118.7} = 17.52$ ✓ $17.52 \times 6.02 \times 10^{23} = 1.05 \times 10^{25}$ atoms ✓	2	ALLOW 17.5 up to (correctly rounded) calculator value of 17.52316765 DO NOT ALLOW use of 118, which makes moles of Sn = 17.63 ALLOW 105×10^{23} atoms DO NOT ALLOW answers which are not to three sig figs for second marking point ALLOW two marks for answer only of 1.05×10^{25} ALLOW one mark for answer only if not 3 sig figs up to calculator value of $1.054894693 \times 10^{25}$ Eg 100×1 ALLOW ECF for any calculated moles of Sn (based on use of any A_r value) $\times 6.02 \times 10^{23}$ if shown to 3 sig figs DO NOT ALLOW mass of Sn $\times 6.02 \times 10^{23}$



Question		Expected Answers	Marks	Additional Guidance
1	d	$\frac{78.8}{118.7}$ and $\frac{21.2}{16.0}$ OR $= 0.66(4)$ and $= 1.3(25)$ ✓ $\frac{0.66(4)}{0.66(4)} = 1$ $\frac{1.325}{0.66(4)} = 2$ ans = SnO ₂ ✓	2	ALLOW SnO ₂ for one mark if no working shown ALLOW use of 118 for this part IGNORE incorrect rounding provided given to two sig figs IGNORE incorrect symbols e.g. T or Ti for Tin, as long as correct A _r of tin (118.7 or 118) used ALLOW Sn ₂ O for 1 mark ECF if both inverted mole calculations are shown ALLOW Sn ₃ O ₅ with evidence of use of both atomic numbers for one mark ALLOW 2 marks if candidate has adopted the following approach 78.8% of mass = 118.7 100% of mass = 118.7/0.788 = 150.6 (151) 150.6 – 118.7 = 31.9 (32) Both masses would get one mark 31.9/16 = 2
Total			9	


Question			Expected Answers	Marks	Additional Guidance
2	a	i	Any two from ✓✓ H ⁺ SO ₄ ²⁻ HSO ₄ ⁻	2 max	DO NOT ALLOW OH ⁻ IGNORE state symbols Charge is essential ALLOW H ₃ O ⁺ for H ⁺ and SO ₄ ⁻² for SO ₄ ²⁻ One answer incorrect = 1 mark max Two answers incorrect = 0 marks
		ii	Effervescence OR fizzing OR bubbling OR gas produced ✓ K ₂ CO ₃ dissolves OR disappears OR colourless solution is formed ✓ H ₂ SO ₄ + K ₂ CO ₃ → K ₂ SO ₄ + CO ₂ + H ₂ O ✓	3	DO NOT ALLOW 'carbon dioxide produced' without 'gas' DO NOT ALLOW incorrectly named gas produced DO NOT ALLOW 'precipitate forms' = CON ALLOW 'it' for K ₂ CO ₃ DO NOT ALLOW mark for 'dissolves' from state symbols in equation DO NOT ALLOW 'potassium' IGNORE state symbols ALLOW ionic equation
	b	i	$\frac{24.6 \times 0.100}{1000} = 0.00246 \text{ mol}$ ✓ ($2.46 \times 10^{-3} \text{ mol}$)	1	DO NOT ALLOW 0.0025 as this would lead to 100% in part (iii) DO NOT ALLOW 0.0024 due to incorrect rounding
		ii	$0.00246 \times 2 = 0.00492 \text{ mol}$ ✓ ($4.92 \times 10^{-3} \text{ mol}$)	1	ALLOW ECF for ans (i) × 2
		iii	Moles of NaOH in 250 cm ³ = $0.00492 \times \frac{250}{25} = 0.0492 \text{ mol}$ ✓ Mass of NaOH in original sample = $0.0492 \times 40.0 = 1.968 \text{ g}$ ✓ % purity $\frac{1.968}{2.00} \times 100 = 98.4\%$ ✓	3	ALLOW ECF for ans (ii) × 10 ALLOW 1.97g ALLOW ECF for moles of NaOH × 40 ALLOW 98.5% (from use of 1.97) ALLOW ECF for $\frac{\text{mass of NaOH}}{2.00} \times 100$ DO NOT ALLOW ECF for 3rd marking point if answer >100% ALLOW ECF for 3rd marking point if answer = 100% ALLOW molar approach for second and third marks i.e. mol of (expected) NaOH in 2.00 g = $2/40 = 0.05(00) \text{ mol}$ ($0.0492/0.0500$) × 100 = 98.4% 1.6% (the percentage of the impurity present) is likely to be 2 marks, but please check 9.84% has not multiplied up by 10 for first marking point is likely to be 2 marks, but please check
Total				10	

Question		Expected Answers	Marks	Additional Guidance
3	a	3d 4p ✓	1	Correct order is essential ALLOW '3D'
	b	i	1	ALLOW 'can be found' for 'can hold' ALLOW 'area' OR 'volume' OR 'space' for region DO NOT ALLOW 'place' for region DO NOT ALLOW path of an electron IGNORE references to 'orbitals being parts of sub-shells'
		ii	1	
	c	18 ✓	1	
	d	i	2	Mark as pairs IGNORE references to 12th and 13th Three answers with one correct pair = 1 mark Four answers with one correct pair = 1 mark Five answers with both pairs correct = 1 mark Five answers with only one pair correct = 0 marks Six (or more) answers = 0 marks
		ii	2	ALLOW $\text{Al}^{2+}(\text{g}) - \text{e}^- \rightarrow \text{Al}^{3+}(\text{g})$ for 2 marks ALLOW 1 mark for $\text{Al}(\text{g}) \rightarrow \text{Al}^{3+}(\text{g}) + 3\text{e}^-$ as states are correct ALLOW 1 mark for $\text{Al}^{2+}(\text{g}) + 2\text{e}^- \rightarrow \text{Al}^{3+}(\text{g}) + 3\text{e}^-$ as states are correct ALLOW 1 mark if symbol of Al is incorrect, but equation is otherwise fully correct. ALLOW e for electron (i.e. no charge) IGNORE states on electron
Total			8	

Question			Expected Answers	Marks	Additional Guidance
4	a	i	1 = purple / lilac / violet / pink / mauve ✓ 3 = orange ✓	2	ALLOW any combination of these but no others for 1 ALLOW yellow as an alternative for 3 DO NOT ALLOW 'precipitate' in either
		ii	$\text{Cl}_2 + 2\text{Br}^- \longrightarrow 2\text{Cl}^- + \text{Br}_2$ ✓	1	IGNORE state symbols ALLOW correct multiples, including fractions
		iii	Addition of $\text{Br}_2(\text{aq})$ to $\text{I}^-(\text{aq})$ ions ✓	1	ALLOW Addition of bromine to iodide (i.e. aqueous not needed) DO NOT ALLOW Addition of bromine to iodine ALLOW Addition of I_2 to Br^- , but NOT if accompanied by description of displacement of bromine ALLOW $\text{Br}_2 + \text{I}^-$ even if seen in an unbalanced equation
	b	i	Cl_2 is 0 AND HCl is -1 AND HClO is $(+1)$ ✓ Chlorine has been both oxidised and reduced OR Chlorine's oxidation state has increased and decreased ✓ Chlorine has been oxidised (from 0) to $+1$ AND chlorine has been reduced (from 0) to -1 ✓ (These two points together subsume the second marking point)	3	ALLOW $1-$ ALLOW $1+$ Oxidation states may be seen above the equation DO NOT ALLOW Cl^- in HCl DO NOT ALLOW Cl^+ in HClO in text of answer DO NOT ALLOW chlorIDE in place of 'chlorine' IF CORRECT OXIDATION STATES ARE SEEN, ALLOW second and third marking points for: Chlorine is oxidised to form HClO Chlorine is reduced to form HCl ALLOW Cl or Cl_2 for 'chlorine' IGNORE reference to electron loss / gain if correct DO NOT ALLOW 3rd mark for reference to electron loss / gain if incorrect ALLOW one mark for 'disproportionation is when a species is both oxidised and reduced' if chlorine / chloride is not mentioned
		ii	Kills bacteria OR 'kills germs' kills micro-organisms OR makes water safe to drink OR sterilises water ✓ OR 'disinfects'	1	ALLOW to make water potable ALLOW 'removes' for 'kills' IGNORE 'virus' IGNORE 'purifies water'
	c	i	Thermal decomposition ✓	1	DO NOT ALLOW just 'decomposition' or 'thermodecomposition'
		ii	$1.47 = 0.0174$ mol of MgCO_3 ✓ 84.3 $0.0174 \times 24.0 = 0.418$ dm ³ OR (Calculator value $\times 24.0$) = 0.419 dm ³ ✓	2	ALLOW mol of MgCO_3 as calculator value of 0.017437722 or correct rounding to 2 sig figs or more DO NOT ALLOW 0.0175 (this has taken M_r of MgCO_3 as 84) ALLOW , for 2nd mark calculated moles of $\text{MgCO}_3 \times 24(.0)$ as calculator value or correct rounding to 2 sig figs or more [e.g. $0.017 \times 24(.0) = 0.408$] DO NOT ALLOW 84.3 or $1.47 \times 24(.0)$ as no mole calculation has been done ALLOW two marks for correct answer with no working shown

Question			Expected Answers	Marks	Additional Guidance
4	c	iii	The ease of (thermal) decomposition decreases (down the group) ora ✓	1	ALLOW (thermal) stability increases IGNORE more heat would be needed IGNORE 'takes longer' or 'is slower' IGNORE reference to trend in reactivity IGNORE answers which include 'more / less mol of CO ₂ '
			Total	15	

Question	Expected Answers	Marks	Additional Guidance
5 a	 <p>Diagram showing a regular arrangement of labelled 'Li⁺' or '+ ions' with some attempt to show electrons ✓</p> <p>Scattering of labelled electrons between other species OR a statement anywhere of delocalised electrons (can be in text or in diagram) ✓</p> <p>The attraction between + ions and e⁻ is strong OR metallic bonding is strong ✓</p>	3	<p>Lattice diagram must have at least two rows of correctly charged ions and a minimum of 2 ions per row</p> <p>ALLOW as label: + ions, positive ions, cations If '+' is unlabelled in diagram, award label from a correct statement within the text below</p> <p>DO NOT ALLOW 2+, 3+ etc ions DO NOT ALLOW for label or in text: nuclei OR positive atom OR protons</p> <p>ALLOW e⁻ OR e as label for electron</p> <p>ALLOW a lot of energy is needed to break the (metallic) bond</p> <p>DO NOT ALLOW incorrect particles or incorrect attraction e.g. 'intermolecular attraction' or 'nuclear attraction'</p>
b i	 <p>Dot and cross bond + 6 matching electrons on each F atom ✓</p>	1	<p>ALLOW diagram consisting of all dots OR all crosses Circles not essential ALLOW 'F' for fluorine</p>
	<p>ii F₂ has induced dipoles OR temporary dipoles OR van der Waals' forces (between the molecules) ✓ which are weak ✓</p>	2	<p>ALLOW little energy needed to overcome intermolecular bonding for second mark ALLOW 'weak' intermolecular bonding for second mark ALLOW max 1 mark if structure is referred to as giant with first and second marking points correct Award no marks if 'weak' is applied to incorrect bonding. E.g. ionic, covalent, metallic or unspecified bonding</p>

Question			Expected Answers	Marks	Additional Guidance
5	c	i	 <p>Li shown with either 2 or 0 electrons and F shown with 8 electrons with 7 crosses and one dot (or <i>vice versa</i>) ✓ correct charges on both ions ✓</p>	2	<p>For first mark, if 2 electrons are shown in the cation then the 'extra' electron in the anion must match symbol chosen for electrons in the cation IGNORE inner shell electrons ALLOW 'F' for fluorine Circles not essential DO NOT ALLOW Li⁺ with 8 electrons</p> <p>Second mark is independent</p>
		ii	<p>Ions cannot move in a solid ✓</p> <p>Ions can move OR are mobile when molten ✓</p>	2	<p>ALLOW ions are fixed in place IGNORE electrons IGNORE 'charge carriers' or 'charged particles'</p> <p>DO NOT ALLOW ions can move when in solution IGNORE charge carriers IGNORE 'delocalised ions' or 'free ions' ALLOW 'ions can only move when molten' for one mark Any mention of electrons moving when molten is a CON</p>
	d	i	$2\text{B} + 3\text{F}_2 \longrightarrow 2\text{BF}_3$ ✓	1	<p>ALLOW B₂ ALLOW multiples including fractions</p>
		ii	<p><i>Shape</i>: trigonal planar ✓ <i>Bond angle</i>: 120° ✓</p> <p><i>Explanation</i>: Pairs of electrons repel (one another equally) ✓</p> <p>Boron has 3 bonded pairs (and 0 lone pairs) ✓</p>	4	<p>'Trigonal planar' must be seen and spelt correctly at least ONCE</p> <p>DO NOT ALLOW 'atoms repel' or 'electrons repel' ALLOW 'bonds repel'</p> <p>ALLOW diagram showing B atom with three dot-and-cross pairs of electrons, but no lone pairs for 4th mark Must refer to boron / central atom ALLOW 'bonds' for 'bonded pairs'</p>

Question		Expected Answers	Marks	Additional Guidance
5	e	F is more electronegative than N OR $\delta^- \text{F}-\text{N}^{\delta+}$ ✓ Dipoles do not cancel OR NF ₃ is pyramidal (in words) / asymmetrical ✓	2	ALLOW F attracts electrons more than N ALLOW N has a partial positive charge and F has a partial negative charge (partial must be seen) DO NOT ALLOW diagrams that contradict statements about polarity ALLOW unsymmetrical, non-symmetrical etc
	f	(As you go across the period) The atomic radii decreases ✓ The nuclear charge increases OR protons increase ✓ electrons are added to the same shell OR shielding remains the same ✓ greater (nuclear) attraction on (outer) electrons / (outer) shell(s) ✓	4	Use annotations with ticks, crosses ECF etc. for this part Assume 'across the period from Li to F' ALLOW (outer shell) electrons get closer (to nucleus) IGNORE 'atomic number increases', but ALLOW 'proton number' increases IGNORE 'nucleus gets bigger' 'Charge increases' is insufficient ALLOW 'effective nuclear charge increases' OR 'shielded nuclear charge increases' Nuclear OR proton(s) OR nucleus spelt correctly ONCE and used in context of 2nd marking point ALLOW shielding is similar ALLOW screening for shielding DO NOT ALLOW 'subshells' DO NOT ALLOW 'distance is similar' This will CON first marking point ALLOW 'greater (nuclear) pull for greater nuclear attraction' DO NOT ALLOW 'pulled in more' as this is a restatement of the first marking point
Total			21	

OCR (Oxford Cambridge and RSA Examinations)
1 Hills Road
Cambridge
CB1 2EU

OCR Customer Contact Centre

14 – 19 Qualifications (General)

Telephone: 01223 553998

Facsimile: 01223 552627

Email: general.qualifications@ocr.org.uk

www.ocr.org.uk

For staff training purposes and as part of our quality assurance programme your call may be recorded or monitored

Oxford Cambridge and RSA Examinations
is a Company Limited by Guarantee
Registered in England
Registered Office; 1 Hills Road, Cambridge, CB1 2EU
Registered Company Number: 3484466
OCR is an exempt Charity



OCR (Oxford Cambridge and RSA Examinations)
Head office
Telephone: 01223 552552
Facsimile: 01223 552553