

Mark Scheme (Results)

January 2013

GCE Core Mathematics C2 (6664/01)

#### **Edexcel and BTEC Qualifications**

Edexcel and BTEC qualifications come from Pearson, the world's leading learning company. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at <a href="https://www.edexcel.com">www.edexcel.com</a> or <a href="https://www.edexcel.com">www.btec.co.uk</a> for our BTEC qualifications.

Alternatively, you can get in touch with us using the details on our contact us page at www.edexcel.com/contactus.

If you have any subject specific questions about this specification that require the help of a subject specialist, you can speak directly to the subject team at Pearson.

Their contact details can be found on this link: www.edexcel.com/teachingservices.

You can also use our online Ask the Expert service at <a href="www.edexcel.com/ask">www.edexcel.com/ask</a>. You will need an Edexcel username and password to access this service.

### Pearson: helping people progress, everywhere

Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: <a href="https://www.pearson.com/uk">www.pearson.com/uk</a>

January 2013
Publications Code US034362
All the material in this publication is copyright
© Pearson Education Ltd 2013

## **General Marking Guidance**

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.
- Unless indicated in the mark scheme a correct answer with no working should gain full marks for that part of the question.

ALWAYS LEARNING PEARSON

#### **EDEXCEL GCE MATHEMATICS**

## General Instructions for Marking

- 1. The total number of marks for the paper is 75.
- 2. The Edexcel Mathematics mark schemes use the following types of marks:
- **M** marks: method marks are awarded for 'knowing a method and attempting to apply it', unless otherwise indicated.
- A marks: Accuracy marks can only be awarded if the relevant method (M) marks have been earned.
- **B** marks are unconditional accuracy marks (independent of M marks)
- Marks should not be subdivided.

In some instances, the mark distributions (e.g. M1, B1 and A1) printed on the candidate's response may differ from the final mark scheme.

### 3. Abbreviations

These are some of the traditional marking abbreviations that will appear in the mark schemes and can be used.

- bod benefit of doubt
- ft follow through
- the symbol will be √used for correct ft
- cao correct answer only
- cso correct solution only. There must be no errors in this part of the question to obtain this mark
- isw ignore subsequent working
- awrt answers which round to
- SC: special case
- oe or equivalent (and appropriate)
- dep dependent
- indep independent
- dp decimal places
- sf significant figures
- \* The answer is printed on the paper
- The second mark is dependent on gaining the first mark
- 4. All A marks are 'correct answer only' (cao.), unless shown, for example, as A1 ft to indicate that previous wrong working is to be followed through. After a misread however, the subsequent A marks affected are treated as A ft, but incorrect answers should never be awarded A marks.
- 5. For misreading which does not alter the character of a question or materially simplify it, deduct two from any A or B marks gained, in that part of the question affected.
- 6. If a candidate makes more than one attempt at any question:

- If all but one attempt is crossed out, mark the attempt which is NOT crossed out.
- If either all attempts are crossed out or none are crossed out, mark all the attempts and score the highest single attempt.
- 7. Ignore wrong working or incorrect statements following a correct answer.
- 8. The maximum mark allocation for each question/part question(item) is set out in the marking grid and you should allocate a score of '0' or '1' for each mark, or "trait", as shown:

|     | 0 | 1 |
|-----|---|---|
| аМ  |   | • |
| аA  | • |   |
| bM1 |   | • |
| bA1 | • |   |
| bB  | • |   |
| bM2 |   | • |
| bA2 |   | • |
|     |   |   |

# January 2013 6664 Core Mathematics C2 Mark Scheme

| Question<br>Number | Scheme  |  |             |
|--------------------|---|--|-------------|
| 1.                 | $(2-5x)^6$  |  |             |
|                    | $(2^6 =) 64$  | Award this when first seen (not $64x^0$ )  | B1          |
|                    | $+6 \times (2)^{5} (-5x) + \frac{6 \times 5}{2} (2)^{4} (-5x)^{2}$  | Attempt binomial expansion with correct structure for at least one of these terms. E.g. a term of the form: $\binom{6}{p} \times \left(2\right)^{6-p} \left(-5x\right)^p$ with $p=1$ or $p=2$ consistently. Condone sign errors. Condone missing brackets if later work implies correct structure and allow alternative forms for binomial coefficients e.g. $\binom{6}{1}$ or even $\left(\frac{6}{1}\right)$ | M1          |
|                    | -960 <i>x</i>   | Not $+-960x$   | A1 (first)  |
|                    | $(+)6000x^2$  |  | A1 (Second) |
|                    |   |  |             |
|                    |   |  |             |
|                    |   |  |             |
|                    |   |  | (4)         |
|                    |   |  |             |
| Way 2              | 64(1±)  | 64 and $(1 \pm Award when first seen.$   | B1          |
|                    | $\left(1 - \frac{5x}{2}\right)^6 = 1 - 6 \times \frac{5x}{2} + \frac{6 \times 5}{2} \left(-\frac{5x}{2}\right)^2$ | brackets if later work implies correct<br>structure but it must be an expansion of $(1-kx)^6$ where $k \neq \pm 5$   | M1          |
|                    | -960x   | Not $+-960x$   | A1          |
|                    | $(+)6000x^2$  |  | A1          |
|                    |   |  | (4)         |
|                    |   |  |             |
|                    |   |  |             |
|                    |   |  |             |
|                    |   |  |             |

| Question<br>Number | Scheme  |   | Marks |
|--------------------|---|---|-------|
| 2.                 |   |   |       |
| (a)                | f(1) = a+b-4-3 = 0 or $a+b-7=0$                             | Attempt $f(\pm 1)$                                | M1    |
|                    | a+b=7 *   | Must be $f(1)$ and = $0$ needs to be seen         | A1    |
|                    |   |   |       |
|                    |   |   | (2)   |
| <b>(b)</b>         | $f(-2) = a(-2)^3 + b(-2)^2 - 4(-2) - 3 = 9$                 | Attempt $f(\pm 2)$ <b>and uses</b> $f(\pm 2) = 9$ | M1    |
|                    | -8a + 4b + 8 - 3 = 9  | Correct equation with exponents of (-2) removed   | A1    |
|                    | (-8a + 4b = 4)  |   |       |
|                    | Solves the <b>given equation from part</b> (a)              |   |       |
|                    | and their equation in $a$ and $b$ from part                 |   | M1    |
|                    | (b) as far as $a = \dots$ or $b = \dots$                    | D. d.   | A 1   |
|                    | a = 2 and $b = 5$   | Both correct                                      | A1    |
|                    |   |   | (4)   |
|                    |   |   | [6]   |
|                    | Long Divis  | sion  |       |
|                    | $(ax^3 + bx^2 - 4x - 3) \div (x - 1) = ax^2 + px + q$       |   |       |
|                    |   | ,   |       |
| (a)                | where $p$ and $q$ are in terms of $a$ or $b$ or both        |   | M1    |
|                    | and sets their remainder = 0                                |   |       |
|                    | NB Quotient = $ax^2 + (a+b)x + (a+b-4)$                     |   |       |
|                    | a+b=7   | *   | A1    |
|                    |   |   | (2)   |
|                    | $\left(ax^3 + bx^2 - 4x - 3\right) \div \left(x + 1\right)$ | $2) = ax^2 + px + q$                              |       |
| <b>4</b>           | where $p$ and $q$ are in terms o                            | f a or b or both                                  | 3.61  |
| <b>(b)</b>         | and sets their remainder = 9                                |   | M1    |
|                    | NB Quotient = $ax^2 + (b-2a)x + (4a-4-2b)$                  |   |       |
|                    | 4b - 8a + 5   | 5 = 9   | A1    |
|                    | Follow scheme for final 2 marks                             |   |       |
|                    |   |   |       |
|                    |   |   |       |
|                    |   |   |       |
|                    |   |   |       |
|                    |   |   |       |
|                    |   |   |       |
|                    | 1   |   | 1     |

| 3.         |   |   |       |     |
|------------|---|---|-------|-----|
| (a)        | $120000 \times (1.05)^3 = 138915 *$   | Or 120000 × 1.05 × 1.05 × 1.05 = 138915<br>Or 120000, 126000, 132300, 138915  | B1    |     |
|            |   | Or $a = 120000$ and $a \times (1.05)^3 = 138915$  |       |     |
|            |   |   |       | (1) |
| <b>(b)</b> | $120000 \times (1.05)^{n-1} > 200000$   | Allow $n$ or $n - 1$ and ">", "<", or "=" etc.  | M1    |     |
|            | $\log 1.05^{n-1} > \log \left(\frac{5}{3}\right)$   | Takes logs correctly Allow $n$ or $n-1$ and ">", "<", or "=" etc.   | M1    |     |
|            | $(n-1>)\frac{\log\left(\frac{5}{3}\right)}{\log 1.05} \text{ or equivalent}$ $\text{e.g } (n>)\frac{\log\left(\frac{7}{4}\right)}{\log 1.05}$ | Allow $n$ or $n - 1$ and ">", "<", or "=" etc. Allow $1.\dot{6}$ or awrt $1.67$ for $5/3$ .                             | A1    |     |
|            |   | M1: Identifies a calendar year using their value  |       |     |
|            | 2024  | of <i>n</i> or <i>n</i> - 1   | M1A1  |     |
|            |   | A1: 2024  |       |     |
|            |   |   |       |     |
|            |   |   |       |     |
|            |   |   |       | (5) |
|            | $\frac{a(1-r^n)}{1-r} = \frac{120000(1-1.05^{11})}{1-1.05}$   | M1: Correct sum formula with $n = 10$ , 11 or 12  |       |     |
| (c)        | $\frac{1-r}{1-r} = \frac{1-1.05}{1-1.05}$   | A1: Correct numerical expression with $n = 11$  | M1 A1 |     |
|            | 1704814   | Cao (Allow 1704814.00)  | A1    |     |
|            |   |   |       | (3) |
|            |   |   |       | [9] |
|            |   | or trial/improvement in (b)   |       |     |
|            |   | $U_{11} = 195 \ 467.36$ , $U_{12} = 205 \ 240.72$<br>$1^{th}$ or $12^{th}$ terms correctly using a common ratio of 1.05 | 3.51  |     |
|            | (all the  | terms need <b>not</b> be listed)  | M1    |     |
|            |   | gression correctly to reach a term > 200 000  | M1    |     |
|            |   | wrt 195 500 <b>and</b> a "12 <sup>th</sup> " term of awrt 205 200   | A1    |     |
|            | Uses their numbe  | r of terms to identify a calendar year  | M1    |     |
|            |   | 2024  | A1    |     |
|            |   |   |       | (5) |
|            |   |   |       |     |
|            |   |   |       |     |
|            |   |   |       |     |

| 4. |  |   |     |
|----|--|---|-----|
|    | $\cos^{-1}(-0.4) = 113.58 \ (\alpha)$                    | Awrt 114  | B1  |
|    | $3x - 10 = \alpha \Rightarrow x = \frac{\alpha + 10}{3}$ | Uses their $\alpha$ to find $x$ .<br>Allow $x = \frac{\alpha \pm 10}{3} \mathbf{not} \frac{\alpha}{3} \pm 10$ | M1  |
|    | 41.2   |   | A 1 |
|    | x = 41.2   | Awrt  | A1  |
|    | $(3x-10=)360-\alpha$ (246.4)                             | $360 - \alpha$ (can be implied by 246.4)  | M1  |
|    | x = 85.5   | Awrt  | A1  |
|    | $(3x-10=)360+\alpha (=473.57)$                           | $360 + \alpha$ (Can be implied by 473.57)   | M1  |
|    | x = 161.2  | Awrt  | A1  |
|    |  |   |     |
|    |  |   | 1   |
|    |  |   |     |
|    |  |   |     |
|    |  |   |     |
|    |  |   |     |
|    |  |   |     |
|    |  |   |     |
|    |  |   |     |
|    |  |   |     |
|    |  |   |     |
|    |  |   |     |
|    |  |   |     |
|    |  |   |     |
|    |  |   |     |
|    |  |   |     |
|    |  |   |     |
|    |  |   |     |
|    |  |   |     |
|    |  |   |     |
|    |  |   |     |
|    |  |   |     |
|    |  |   |     |
|    |  |   |     |
|    |  |   |     |
|    |  |   |     |
|    |  |   |     |

| 5.           |   |   |          |            |
|--------------|---|---|----------|------------|
| (a)          |   |   |          |            |
| (i)          | The centre is at (10, 12)   | B1: $x = 10$<br>B1: $y = 12$  | B1 B1    |            |
| (ii)         | Uses $(x-10)^2 + (y-12)^2 =$  | $-195+100+144 \Rightarrow r = \dots$  | M1       |            |
|              | Completes the square for both $x$<br>$(x \pm "10")^2 \pm a$ and $(y \pm "12")^2$<br>Allow errors in obtaining their | $\pm b$ and $+195 = 0$ , $(a, b \neq 0)$<br>$r^2$ but must find square root   |          |            |
|              | $r = \sqrt{10^2 + 12^2 - 195}$  | A correct numerical expression for <i>r</i> including the square root and can implied by a correct value for <i>r</i> | A1       |            |
|              | r = 7   | Not $r = \pm 7$ unless – 7 is rejected  | A1       |            |
|              |   |   |          | (5)        |
|              | Compares the given equation with $x^2 + y^2 + 2gx + 2fy + c = 0$ to write   | B1: $x = 10$  | B1B1     |            |
| (a)<br>Way 2 | down centre $(-g, -f)$ i.e. $(10, 12)$  | B1: <i>y</i> = 12   | DIDI     |            |
| ·            | Uses $r = \sqrt{(\pm "10")^2 + (\pm "12")^2 - c}$   |   | M1       |            |
|              | $r = \sqrt{10^2 + 12^2 - 195}$  | A correct numerical expression for <i>r</i>   | A1       |            |
|              | r = 7   | _   | A1       |            |
|              |   |   |          | (5)        |
|              |   |   |          |            |
| (b)          | $MN = \sqrt{(25 - "10")^2 + (32 - "12")^2}$   | Correct use of Pythagoras   | M1       |            |
|              | $MN\left(=\sqrt{625}\right)=25$   |   | A1       |            |
|              |   |   |          | (2)        |
| (c)          | $NP = \sqrt{("25"^2 - "7"^2)}$  | $NP = \sqrt{(MN^2 - r^2)}$  | M1       |            |
|              | $NP\left(=\sqrt{576}\right)=24$   |   | A1       |            |
|              |   |   |          | (2)        |
| (c)<br>Way 2 | $\cos(NMP) = \frac{7}{"25"} \Rightarrow NP = "25" \sin(NR)$   | MP) Correct strategy for finding NP   | M1       |            |
| •            | NP = 24   |   | A1       |            |
|              |   |   |          | <b>(2)</b> |
|              |   |   |          | [9]        |
|              |   |   |          |            |
|              |   |   |          |            |
|              |   |   |          |            |
|              | l   | l   | <u> </u> |            |

| 6.  |   |   |       |
|-----|---|---|-------|
| (a) | $2\log(x+15) = \log(x+15)^2$  |   | B1    |
|     | $\log(x+15)^2 - \log x = \log \frac{(x+15)^2}{x}$                   | Correct use of $\log a - \log b = \log \frac{a}{b}$                                       | M1    |
|     |   |   |       |
|     | $2^6 = 64 \text{ or } \log_2 64 = 6$                                | 64 used in the correct context  | B1    |
|     | $\log_2 \frac{(x+15)^2}{x} = 6 \Rightarrow \frac{(x+15)^2}{x} = 64$ | Removes logs correctly  | M1    |
|     |   |   |       |
|     | $\Rightarrow x^2 + 30x + 225 = 64x$ or $x + 30 + 225x^{-1} = 64$    | Must see expansion of $(x+15)^2$ to score the final mark.                                 |       |
|     | $\therefore x^2 - 34x + 225 = 0 *$                                  | score the final mark.   | A1    |
|     | 3+1 223 = 0   |   | (5)   |
| (b) | $(x-25)(x-9) = 0 \Rightarrow x = 25 \text{ or } x = 9$              | M1: Correct attempt to solve the <b>given</b> quadratic as far as $x =$ A1: Both 25 and 9 | M1 A1 |
|     |   |   | (2)   |
|     | 1   |   | [7]   |
|     |   |   |       |
|     |   |   |       |
|     |   |   |       |
|     |   |   |       |
|     |   |   |       |
|     |   |   |       |
|     |   |   |       |
|     |   |   |       |
| _   |   |   |       |
|     |   |   |       |
| _   |   |   |       |
|     |   |   |       |
|     |   |   |       |
|     |   |   |       |
|     |   |   |       |
|     |   |   | †     |
|     |   |   |       |
|     |   |   |       |
|     |   |   |       |

|            |  |  | I        | [12] |
|------------|--|--|----------|------|
|            |  |  |          | (4)  |
|            | Perimeter = 27.2 or 27.3   | Awrt 27.2 or awrt 27.3   | A1       |      |
|            | Perimeter = $ZY + WY + Arc Length$   | 9 + 2 + Any Arc  | M1       |      |
| (d)        | Or $8\pi - 4 \times 2.22$  | Or circumference – minor arc A1: Correct ft expression                   | M1A1ft   |      |
| (4)        | Arc length = $4 \times 4.06 (= 16.24)$   | M1: $4 \times their(2\pi - 2.22)$  | M1 A 1ft |      |
|            |  |  |          | (3)  |
|            | $= 42.1 \text{ cm}^2 \text{ or } 42.0 \text{ cm}^2$  | Awrt 42.1 or 42.0 (Or <u>just</u> 42)                                    | A1       |      |
|            | So area required = "9.56" + "32.5"   | Their Triangle XYZ + (part (b) answe or correct attempt at major sector) | r M1     |      |
| (c)        | Area of triangle = $\frac{1}{2} \times 4 \times 6 \times \sin 2.22 (= 9.56)$   | Correct expression for the area of triangle XYZ                          | B1       | (3)  |
|            | = 32.5   | Awrt 32.5  | A1       | (2)  |
|            | $\pi \times 4^{2}$ $\pi \times 4^{2} - \frac{1}{2} \times 4^{2} \times 2.22 = 32.5$  | Correct method for circle - minor sector area                            | M1       |      |
|            | $\pi \times 4^2$   | Correct expression for circle area                                       | B1       |      |
| (b) Way2   | Circle – Mino  | or sector  |          | (3)  |
|            | 32.5   | Awrt 32.5  | A1       | (2)  |
|            | $\frac{1}{2} \times 4^2 \times "4.06"$   | Correct method for major sector area.                                    | M1       |      |
| <b>(b)</b> | $2\pi - 2.22 (= 4.06366)$  | $2\pi - 2.22$ or awrt 4.06   | B1       |      |
|            | <i>XY</i> = 9.00   |  | A1       | (2)  |
|            | $XY^2 = 81.01$   |  | A 1      |      |
| (a) Way 2  | $XY^2 = 4^2 + 6^2 - 2 \times 4 \times 6 \cos 2.22 \Rightarrow XY^2 =$ Correct use of cosine rule leading to a value for $XY^2$ |  | M1       | ` /  |
|            | $\alpha = 2.22$ * (NB $\alpha = 2.219516005$ )   | Cso (2.22 must be seen here)   | Λ1       | (2)  |
|            | $\cos \alpha = \frac{4^2 + 6^2 - 9^2}{2 \times 4 \times 6} \left( = -\frac{29}{48} = -0.604 \right)$ $\alpha = 2.22  *$        |  | A1       |      |
| (a)        | $9^2 = 4^2 + 6^2 - 2 \times 4 \times 6 \cos \alpha \Rightarrow \cos \alpha = \dots$  | leading to a value for cos $\alpha$                                      | M1       |      |
| 7.         |  |  |          |      |

| 8.          | y = 6  | $-3x - \frac{4}{x^3}$ $M1: x^n \to x^{n-1}$   |           |
|-------------|--|---|-----------|
| (a)         | 1  dy $12$   |   | M1 A1     |
|             | $\frac{dy}{dx} = 0 \Rightarrow -3 + \frac{12}{x^4} = 0 \Rightarrow x = \dots \text{ or }$ $\frac{dy}{dx} = -3 + \frac{12}{\sqrt{2}^4}$   | y' = 0 and attempt to solve for $xMay be implied by \frac{dy}{dx} = -3 + \frac{12}{x^4} = 0 \Rightarrow \frac{12}{x^4} = 3 \Rightarrow x = \dots \text{ or} Substitutes x = \sqrt{2} into their y'$ | M1        |
|             | So $x^4 = 4$ and $x = \sqrt{2}$ or $\frac{dy}{dx} = -3 + \frac{12}{\left(\sqrt{2}\right)^4} \text{ or } -3 + 12\left(\sqrt{2}\right)^{-4} = 0$   | Correct completion to answer with no errors by solving their $y' = 0$ or substituting $x = \sqrt{2}$ into their $y'$  | A1        |
|             |  |   |           |
| <i>a</i> .) | \[ \sum_{\text{\sigma}} \]   |   | (4)       |
| <b>(b)</b>  | $x = -\sqrt{2}$  | Awrt -1.41  | B1 (1)    |
| (c)         | $\frac{d^2 y}{dx^2} = \frac{-48}{x^5} \text{ or } -48x^{-5}$   | Follow through their first derivative from part (a)   | B1ft      |
| (d)         | An appreciation that either $y'' > 0 \Rightarrow$ a minimum or $y'' < 0 \Rightarrow$ a maximum   |   | (1)<br>B1 |
|             | Maximum at P as $y'' < 0$  | Cso   | B1        |
|             | Need a fully correct solution for this mark. $y''$ need not be evaluated but must be correct and there must be reference to P or to $\sqrt{2}$ and negative or < 0 and maximum. There must be no incorrect or contradictory statements (NB allow $y''$ = awrt-8 or -9) |   |           |
|             | Minimum at Q as $y'' > 0$  | Cso   | B1        |
|             | · ·  | k. $y''$ need not be evaluated but must be d there must be reference to P or to $-\sqrt{2}$ must be no incorrect or contradictory   |           |
|             |  |   | (3)       |
|             | Other methods for identifying the nature of the for finding values of y or dy/dx either side of B1's for fully correct solutions to identify the   |   | [9]       |

| 9.  | y = 27 - 2x   | $x - 9\sqrt{x} - \frac{16}{x^2}$            |                                   |          |  |
|-----|---|---|-----------------------------------|----------|--|
| (a) | 6.272 , 3.634   |   |                                   | B1, B1   |  |
|     |   |   | 1                                 |          |  |
|     | 1 1 1   |   |                                   | (2)      |  |
| (b) | $\frac{1}{2} \times \frac{1}{2}$ or $\frac{1}{4}$   |   |                                   | B1       |  |
|     | $\{(0+0)+2(5.866+"6.272"+5.210$   | +"3.634"+1.856)}                            | Need {} or implied later for A1ft | M1A1ft   |  |
|     |   |   |                                   |          |  |
|     | $\frac{1}{2} \times 0.5 \{ (0+0) + 2(5.866 + "6.272" + 5.210 + "3.634" + 1.856) \}$   |   |                                   |          |  |
|     | $=\frac{1}{4} \times 45.676$  |   |                                   |          |  |
|     | = 11.42   | A1  |                                   |          |  |
|     |   |   |                                   |          |  |
|     |   |   |                                   |          |  |
|     |   |   |                                   | (4)      |  |
|     | $\int y  dx = 27x - x^2 - 6x^{\frac{3}{2}} + 16x^{-1} \left( +c \right)$  | M1: $x^n \to x^{n+1}$ on<br>A1: $27x - x^2$ | any term                          |          |  |
|     | $\int y  dx = 27x - x - 6x^2 + 16x  (+c)$   | A1: $-6x^{\frac{3}{2}}$                     |                                   | M1A1A1A1 |  |
| (0) |   | A1: $+16x^{-1}$                             |                                   |          |  |
| (c) |   |   |                                   |          |  |
|     | $\left(27(4)-(4)^2-6(4)^{\frac{2}{2}}+16(4)^{-1}\right)$  | Attempt to subtra                           |                                   |          |  |
|     | $ \left(27(4) - (4)^{2} - 6(4)^{\frac{3}{2}} + 16(4)^{-1}\right) - \left(27(1) - (1)^{2} - 6(1)^{\frac{3}{2}} + 16(1)^{-1}\right) $ | round using the line Dependent on the       |                                   | dM1      |  |
|     |   | - 36)                                       |                                   |          |  |
|     | 12  | Cao   |                                   | A1       |  |
|     |   |   |                                   | (6)      |  |
|     |   |   |                                   | [12]     |  |

Further copies of this publication are available from Edexcel Publications, Adamsway, Mansfield, Notts, NG18 4FN

Telephone 01623 467467 Fax 01623 450481 Email <u>publication.orders@edexcel.com</u> Order Code US034362 January 2013

For more information on Edexcel qualifications, please visit our website  $\underline{www.edexcel.com}$ 

Pearson Education Limited. Registered company number 872828 with its registered office at Edinburgh Gate, Harlow, Essex CM20 2JE





