

June 2006
6663 Core Mathematics C1
Mark Scheme

Question number	Scheme	Marks
	$2x + \frac{x^{\frac{1}{2}}}{\frac{1}{2}} \quad (+c)$ $= 2x^3 + 2x + 2x^{\frac{1}{2}}$ <p>for some attempt to integrate $x^n \rightarrow x^{n+1}$</p> <p>1st A1 for either $\frac{6}{3}x^3$ or $\frac{x^{\frac{1}{2}}}{\frac{1}{2}}$ or better</p> <p>for all terms in x correct. Allow $2\sqrt{x}$ and $2x^1$.</p> <p>for $+c$, when first seen with a changed expression.</p>	Total 4 marks

Critical Values

$$(x \pm a)(x \pm b) \text{ with } ab=18 \quad \text{or} \quad x = \frac{7 \pm \sqrt{49 - 72}}{2} \quad \text{or}$$



$$\left(x - \frac{7}{2}\right)^2 \pm \left(\frac{7}{2}\right)^2 - 18$$

$$(x - 9)(x + 2) \quad \text{or} \quad x = \frac{7 \pm 11}{2} \quad \text{or} \quad x = \frac{7}{2} \pm \frac{11}{2}$$

Solving Inequality $x > 9$ or $x < -2$ Choosing “outside”

Total 4 marks

For attempting to find critical values.
Factors alone are OK for M1, $x =$ appearing somewhere for the formula and as written for completing the square
1st A1 Factors alone are OK. Formula or completing the square need $x =$ as written.
2nd M1 For choosing outside region. Can f.t. their critical values. They must have two different critical values.
– $2 > x > 9$ is M1A0 but ignore if it follows a correct version
– $-2 < x < 9$ is M0A0 whatever the diagram looks like.
2nd A1 Use of \geq in final answer gets A0

Question number	Scheme	Marks
	U shape touching x -axis	B1
	$(-3, 0)$	B1
		B1 (3)
	Translated parallel to y -axis up $(0, 9 + k)$	(2)
Total 5 marks		
<p>2nd B1 They can score this even if other intersections with the x-axis are given.</p> <p>2nd B1 & 3rd B1 The -3 and 9 can appear on the sketch as shown</p> <p>M1 Follow their curve in (a) up only. If it is not obvious do not give it. e.g. if it cuts y-axis in (a) but doesn't in (b) then it is M0.</p> <p>B1f.t. Follow through their 9</p>		

Question number	Scheme	Marks
(a)	$a_2 = 4$ $a_3 = 3 \times a_2 - 5 = 7$	B1f.t. (2)
(b)	$a_4 = 3a_3 - 5 (= 16)$ and $a_5 = 3a_4 - 5 (= 43)$ $3 + 4 + 7 + 16 + 43$ $= 73$	A1c.a.o. (3)
Total 5 marks		

- 2nd B1f.t. Follow through their a_2 but it must be a value.
 $3 \times 4 - 5$ is B0. Give wherever it is first seen.
- 1st M1 For two further attempts to use of $a_{n+1} = 3a_n - 5$, wherever seen. Condone arithmetic slips
- 2nd M1 For attempting to add 5 relevant terms (i.e. terms derived from an attempt to use the recurrence formula) or an expression. Follow through their values for $a_2 - a_5$
- Use of formulae for arithmetic series is M0A0 but could get 1st M1 if a_4 and a_5 are correctly attempted.

Question number	Scheme	Marks
5.	<p>(a) $(y = x^4 + 6x^{\frac{1}{2}} \Rightarrow y' =) 4x^3 + 3x^{-\frac{1}{2}}$ or $4x^3 + \frac{3}{\sqrt{x}}$ A1 (3)</p> <p>(b) $(x+4)^2 = x^2 + 8x + 16$ $\frac{(x+4)^2}{x} = x + 8 + 16x^{-1}$ (allow 4+4 for 8) $(y = \frac{(x+4)^2}{x} \Rightarrow y' =) 1 - 16x^{-2}$ o.e. M1A1 (4)</p>	
Total 7 marks		
	<p>(a) M1 For some attempt to differentiate $x^n \rightarrow x^{n-1}$ 1st A1 For one correct term as printed. 2nd A1 For both terms correct as printed. $4x^3 + 3x^{-\frac{1}{2}} + c$ scores M1A1A0</p> <p>(b) 1st M1 For attempt to expand $(x+4)^2$, must have x^2, x, x^0 terms and at least 2 correct e.g. $x^2 + 8x + 8$ or $x^2 + 2x + 16$ 1st A1 Correct expression for $\frac{(x+4)^2}{x}$. As printed but allow $\frac{16}{x}$ and $8x^0$. 2nd M1 For some correct differentiation, any term. Can follow through their simplification. N.B. $\frac{x^2 + 8x + 16}{x}$ giving rise to $(2x + 8)/1$ is M0A0</p> <p>ALT <u>Product or Quotient rule</u> (If in doubt send to review) M2 For correct use of product or quotient rule. Apply usual rules on formulae. 1st A1 For $\frac{2(x+4)}{x}$ or $\frac{2x(x+4)}{x^2}$ 2nd A1 for $-\frac{(x+4)^2}{x^2}$</p>	

Question number	Scheme	Marks
6.	<p>(a) $16 + 4\sqrt{3} - 4\sqrt{3} - (\sqrt{3})^2$ or $16 - 3$ $= 13$</p> <p>(b) $\frac{26}{4 + \sqrt{3}} \times \frac{4 - \sqrt{3}}{4 - \sqrt{3}}$ $= \frac{26(4 - \sqrt{3})}{13} = \underline{8 - 2\sqrt{3}}$ or $8 + (-2)\sqrt{3}$ or $a = 8$ and $b = -2$</p>	<p>M1 A1c.a.o (2)</p> <p>M1 A1 (2)</p> <p>Total 4 marks</p>
(a)	<p>M1 For 4 terms, at least 3 correct e.g. $8 + 4\sqrt{3} - 4\sqrt{3} - (\sqrt{3})^2$ or $16 \pm 8\sqrt{3} - (\sqrt{3})^2$ or $16 + 3$ 4^2 instead of 16 is OK $(4 + \sqrt{3})(4 + \sqrt{3})$ scores M0A0</p>	
(b)	<p>M1 For a correct attempt to rationalise the denominator can be implied NB $\frac{-4 + \sqrt{3}}{-4 + \sqrt{3}}$ is OK</p>	

Question number	Scheme	Marks
7.	$a + (n - 1)d = k$ $(u_{11} =) a + 10d = 9$ $\frac{n}{2}[2a + (n - 1)d] = 77$ or $\frac{(a + l)}{2} \times n = 77$ $(S_{11} =) \frac{11}{2}(2a + 10d) = 77$ or $\frac{(a + 9)}{2} \times 11 = 77$ e.g. $a + 10d = 9$ $a + 5d = 7$ or $a + 9 = 14$ $a = 5$ and $d = 0.4$ or exact equivalent	M1 A1c.a.o. M1 A1 M1 A1 A1 Total 7 marks
<p>1st M1 Use of u_n to form a linear equation in a and d. $a + nd = 9$ is M0A0</p> <p>1st A1 For $a + 10d = 9$.</p> <p>2nd M1 Use of S_n to form an equation for a and d (LHS) or in a (RHS)</p> <p>2nd A1 A correct equation based on S_n.</p> <p>For 1st 2 Ms they must write n or use $n = 11$.</p> <p>3rd M1 Solving (LHS simultaneously) or (RHS a linear equation in a)</p> <p>Must lead to $a = \dots$ or $d = \dots$ and depends on one previous M</p> <p>3rd A1 for $a = 5$</p> <p>4th A1 for $d = 0.4$ (o.e.)</p> <p><u>ALT</u> Uses $\frac{(a + l)}{2} \times n = 77$ to get $a = 5$, gets second and third M1A1 i.e. 4/7</p> <p>Then uses $\frac{n}{2}[2a + (n - 1)d] = 77$ to get d, gets 1st M1A1 and 4th A1</p> <p><u>MR</u> Consistent MR of 11 for 9 leading to $a = 3$, $d = 0.8$ scores M1A0M1A0M1A1ftA1ft</p>		


Question number	Marks	Scheme	
8. (a)	$b^2 - 4ac = 4p^2 - 4(3p + 4) = 4p^2 - 12p - 16 (=0)$ M1, A1 or $(x + p)^2 - p^2 + (3p + 4) = 0 \Rightarrow p^2 - 3p - 4 (=0)$ $(p - 4)(p + 1) = 0$ M1 $p = (-1 \text{ or } 4)$ A1c.s.o. (4)		
(b)	$x = \frac{-b}{2a}$ or $(x + p)(x + p) = 0 \Rightarrow x = \dots$ M1 $x (= -p) = \underline{-4}$ A1f.t. (2)		
	6		
(a)	1 st M1 For use of $b^2 - 4ac$ or a full attempt to complete the square leading to a 3TQ in p . May use $b^2 = 4ac$. One of b or c must be correct. 1 st A1 For a correct 3TQ in p . Condone missing “=0” but all 3 terms must be on one side . 2 nd M1 For attempt to solve their 3TQ leading to $p = \dots$ 2 nd A1 For $p = 4$ (ignore $p = -1$). $b^2 = 4ac$ leading to $p^2 = 4(3p + 4)$ and then "spotting" $p = 4$ scores 4/4.		
(b)	M1 For a full method leading to a repeated root $x = \dots$ A1f.t. For $x = -4$ (- their p)		

Trial and Improvement

M2 For substituting values of p into the equation and attempting to factorize.

(Really need to get to $p = 4$ or -1)

A2c.s.o. Achieve $p = 4$. Don't give without valid method being seen.

Question number	Marks	Scheme
<p>9. (a)</p> <p>= 1</p> <p>(b)</p> <p>(c)</p>	<p> $f(x) = x[(x-6)(x-2)+3]$ or $x^3 - 6x^2 - 2x^2 + 12x + 3x = x($ M1 $f(x) = x(x^2 - 8x + 15)$ A1 A1 (3) $(x^2 - 8x + 15) = (x-5)(x-3)$ M1 $f(x) = x(x-5)(x-3)$ A1 (2) B1  </p>	<p> $x^3 - 6x^2 - 2x^2 + 12x + 3x = x($ $b = -8$ or $c = 15$ both and a Shape their 3 <u>or</u> their 5 B1f.t. </p>

implication

(3)
and (0,0) by

both their 3 and their 5 B1f.t.

0 3 5 x

8

(a) M1 for a correct method to get the factor of x . $x($ as printed is the minimum.

1st A1 for $b = -8$ or $c = 15$.

-8 comes from -6-2 and must be coefficient of x , and 15 from $6x^2+3$ and must have no x s.

2nd A1 for $a = 1$, $b = -8$ and $c = 15$. Must have $x(x^2 - 8x + 15)$.

(b) M1 for attempt to factorise their 3TQ from part (a).

A1 for all 3 terms correct. They must include the x .

For part (c) they must have at most 2 non-zero roots of their $f(x) = 0$ to

~~fit their 3 and their 5.~~

(c) 1st B1 for correct shape (i.e. from bottom left to top right and two turning points.)

2nd B1f.t. for crossing at their 3 or their 5 indicated on graph or in text.

3rd B1f.t. if graph passes through (0, 0) [needn't be marked] and both their 3 and their 5.

Question number	Marks	Scheme
<p>10.(a)</p> <p>$\frac{1}{3}$</p>	<p> $f(x) = \frac{2x^2}{2} + \frac{3x^{-1}}{-1} (+c)$ M1A1 $(3, 7\frac{1}{2})$ gives $\frac{15}{2} = 9 - \frac{3}{3} + c$ M1A1f.t. $c = -\frac{1}{2}$ A1 (5) </p> <p>(b) $f(-2) = 4 + \frac{3}{2} - \frac{1}{2} \quad (*)$ B1c.s.o. (1)</p> <p>(c) $m = -4 + \frac{3}{4}, = -3.25$ M1,A1 Equation of tangent is: $y - 5 = -3.25(x + 2)$ M1 $4y + 13x + 6 = 0$ A1 (4)</p>	<p>$-\frac{3}{x}$ is OK</p> <p>3^2 or 3^{-1} are OK instead of 9 or</p> <p>o.e.</p>
<p>(a)</p> <p>(b)</p> <p>(c)</p>	<p>10</p> <p> 1st M1 for some attempt to integrate $x^n \rightarrow x^{n+1}$ 1st A1 for both x terms as printed or better. Ignore $(+c)$ here. 2nd M1 for use of $(3, 7\frac{1}{2})$ or $(-2, 5)$ to form an equation for c. There must be some correct substitution. No $+c$ is M0. Some changes in x terms of function needed. 2nd A1f.t. for a correct equation for c. Follow through their integration. They must tidy up fraction/fraction and signs (e.g. - - to +). </p> <p> (b) B1cso If $(-2, 5)$ is used to find c in (a) B0 here unless they verify $f(3)=7.5$. </p> <p> (c) 1st M1 for attempting $m = f'(\pm 2)$ 1st A1 for $-\frac{13}{4}$ or -3.25 </p>	

2nd M1 for attempting equation of tangent at $(-2, 5)$, f.t. their m , based
on $\frac{dy}{dx}$.

2nd A1 o.e. must have a , b and c integers and $= 0$.

Treat (a) and (b) together as a batch of 6 marks.

Question number	Marks	Scheme
11.(a)	$m = \frac{8-2}{11+1} \quad (= \frac{1}{2})$ <p>M1 A1</p> $y - 2 = \frac{1}{2}(x - -1) \quad \text{or} \quad y - 8 = \frac{1}{2}(x - 11) \quad \text{o.e.}$ <p>M1</p> $y = \frac{1}{2}x + \frac{5}{2}$ <p>accept exact equivalents</p> <p>e.g. $\frac{6}{12}$ A1 c.a.o. (4)</p>	
(b)	<p>Gradient of $l_2 = -2$</p> <p>M1</p> <p>Equation of l_2: $y - 0 = -2(x - 10) \quad [y = -2x + 20]$</p> <p>M1 $\frac{1}{2}x + \frac{5}{2} = -2x + 20$</p> <p>M1</p> <p><u>$x = 7$ and $y = 6$</u></p> <p>Ms A1, A1 (5)</p> <p>depend on all 3</p>	
(c)	$RS^2 = (10 - 7)^2 + (0 - 6)^2 (= 3^2 + 6^2)$ <p>M1</p> $RS = \sqrt{45} = 3\sqrt{5} \quad (*)$ <p>A1 c.s.o. (2)</p>	
(d)	$PQ = \sqrt{12^2 + 6^2} = 6\sqrt{5} \quad \text{or} \quad \sqrt{180} \quad \text{or} \quad PS = 4\sqrt{5} \quad \text{and} \quad SQ = 2\sqrt{5}$ <p>M1, A1</p> $\text{Area} = \frac{1}{2}PQ \times RS = \frac{1}{2}6\sqrt{5} \times 3\sqrt{5}$ <p>dM1</p>	

$$= 45$$

A1 c.a.o. (4)

15

(a)	1 st M1	for attempting $\frac{y_1 - y_2}{x_1 - x_2}$, must be y over x. No formula condone
	one sign slip, but if correct substitution.	formula is quoted then there must be some
	1 st A1	for a fully correct expression, needn't be simplified.
	2 nd M1	for attempting to find equation of l_1 .
(b)	1 st M1	for using the perpendicular gradient rule
	2 nd M1	for attempting to find equation of l_2 . Follow their gradient
	provided different.	
	3 rd M1	for forming a suitable equation to find S.
(c)	M1	for expression for RS or RS^2 . Ft their S coordinates
(d)	1 st M1	for expression for PQ or PQ^2 .
	$PQ^2 = 12^2 + 6^2$ is M1 but $PQ = 12^2 + 6^2$ is M0	
		Allow one numerical slip.
	2 nd dM1	for a full, correct attempt at area of triangle. Dependent on
	previous M1.	
