

Mark Scheme Summer 2009

GCE

GCE Mathematics (8371/8374; 9371/9374)

Edexcel is one of the leading examining and awarding bodies in the UK and throughout the world. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers.

Through a network of UK and overseas offices, Edexcel's centres receive the support they need to help them deliver their education and training programmes to learners.

For further information, please call our GCE line on 0844 576 0025, our GCSE team on 0844 576 0027, or visit our website at www.edexcel.com.

If you have any subject specific questions about the content of this Mark Scheme that require the help of a subject specialist, you may find our **Ask The Expert** email service helpful.

Ask The Expert can be accessed online at the following link:

<http://www.edexcel.com/Aboutus/contact-us/>

Summer 2009

Publications Code UA021531

All the material in this publication is copyright

© Edexcel Ltd 2009

Contents

1.	6663 Core Mathematics 1	5
2.	6664 Core Mathematics 2	17
3.	6665 Core Mathematics 3	27
4.	6666 Core Mathematics 4	39
5.	6667 Further Pure Mathematics 1 (new)	47
6.	6668 Further Pure Mathematics 2 (new)	55
7.	6669 Further Pure Mathematics 3 (new)	65
8.	6674 Further Pure Mathematics 1 (legacy)	73
9.	6675 Further Pure Mathematics 2 (legacy)	81
10.	6676 Further Pure Mathematics 3 (legacy)	87
11.	6677 Mechanics 1	93
12.	6678 Mechanics 2	97
13.	6679 Mechanics 3	103
14.	6680 Mechanics 4	109
15.	6681 Mechanics 5	115
16.	6683 Statistics 1	121
17.	6684 Statistics 2	129
18.	6691 Statistics 3	137
19.	6686 Statistics 4	145
20.	6689 Decision Mathematics 1	151
21.	6690 Decision Mathematics 2	161

June 2009
6663 Core Mathematics C1
Mark Scheme

Question Number	Scheme	Marks
Q1 (a)	$(3\sqrt{7})^2 = 63$	B1 (1)
(b)	$(8 + \sqrt{5})(2 - \sqrt{5}) = 16 - 5 + 2\sqrt{5} - 8\sqrt{5}$ $= 11, -6\sqrt{5}$	M1 A1, A1 (3) [4]
(a)	B1 for 63 only	
(b)	M1 for an attempt to expand <u>their</u> brackets with ≥ 3 terms correct. They may collect the $\sqrt{5}$ terms to get $16 - 5 - 6\sqrt{5}$ Allow $-\sqrt{5} \times \sqrt{5}$ or $-(\sqrt{5})^2$ or $-\sqrt{25}$ instead of the -5 These 4 values may appear in a list or table but they should have minus signs included The next two marks should be awarded for the final answer but check that correct values follow from correct working. Do not use ISW rule 1 st A1 for 11 from $16 - 5$ <u>or</u> $-6\sqrt{5}$ from $-8\sqrt{5} + 2\sqrt{5}$ 2 nd A1 for <u>both</u> 11 and $-6\sqrt{5}$. <u>S.C - Double sign error in expansion</u> For $16 - 5 - 2\sqrt{5} + 8\sqrt{5}$ leading to $11 + \dots$ allow <u>one</u> mark	

Question Number	Scheme	Marks
Q2	$32 = 2^5$ or $2048 = 2^{11}$, $\sqrt{2} = 2^{\frac{1}{2}}$ or $\sqrt{2048} = (2048)^{\frac{1}{2}}$ $a = \frac{11}{2}$ (or $5\frac{1}{2}$ or 5.5)	B1, B1 B1 [3]
	<p>1st B1 for $32 = 2^5$ or $2048 = 2^{11}$ This should be explicitly seen: $32\sqrt{2} = 2^a$ followed by $2^5\sqrt{2} = 2^a$ is OK Even writing $32 \times 2 = 2^5 \times 2 (= 2^6)$ is OK but simply writing $32 \times 2 = 2^6$ is NOT</p> <p>2nd B1 for $2^{\frac{1}{2}}$ or $(2048)^{\frac{1}{2}}$ seen. This mark may be implied</p> <p>3rd B1 for answer as written. Need $a = \dots$ so $2^{\frac{11}{2}}$ is B0</p> <p>$a = \frac{11}{2}$ (or $5\frac{1}{2}$ or 5.5) with no working scores full marks.</p> <p>If $a = 5.5$ seen then award 3/3 unless it is clear that the value follows from totally incorrect work.</p> <p>Part solutions: e.g. $2^5\sqrt{2}$ scores the first B1.</p> <p><u>Special case:</u></p> <p>If $\sqrt{2} = 2^{\frac{1}{2}}$ is not explicitly seen, but the final answer includes $\frac{1}{2}$, e.g. $a = 2\frac{1}{2}$, $a = 4\frac{1}{2}$, the second B1 is given by implication.</p>	

Question Number	Scheme	Marks
Q3 (a)	$\frac{dy}{dx} = 6x^2 - 6x^{-3}$	M1 A1 A1 (3)
(b)	$\frac{2x^4}{4} + \frac{3x^{-1}}{-1} (+C)$	M1 A1
	$\frac{x^4}{2} - 3x^{-1} + C$	A1 (3) [6]
(a)	<p>M1 for an attempt to differentiate $x^n \rightarrow x^{n-1}$</p> <p>1st A1 for $6x^2$</p> <p>2nd A1 for $-6x^{-3}$ or $-\frac{6}{x^3}$ Condone $+ -6x^{-3}$ here. Inclusion of $+c$ scores A0 here.</p>	
(b)	<p>M1 for some attempt to integrate an x term of the given y. $x^n \rightarrow x^{n+1}$</p> <p>1st A1 for both x terms correct but unsimplified- as printed or better. Ignore $+c$ here</p> <p>2nd A1 for both x terms correct and simplified and $+c$. Accept $-\frac{3}{x}$ but <u>NOT</u> $+ -3x^{-1}$</p> <p>Condone the $+c$ appearing on the first (unsimplified) line but missing on the final (simplified) line</p> <p>Apply ISW if a correct answer is seen</p> <p>If part (b) is attempted first and this is clearly labelled then apply the scheme and allow the marks. Otherwise assume the first solution is for part (a).</p>	

Question Number	Scheme	Marks
Q4 (a)	$5x > 10, x > 2$ [Condone $x > \frac{10}{2} = 2$ for M1A1]	M1, A1 (2)
(b)	$(2x + 3)(x - 4) = 0$, 'Critical values' are $-\frac{3}{2}$ and 4 $-\frac{3}{2} < x < 4$	M1, A1 M1 A1ft (4)
(c)	$2 < x < 4$	B1ft (1) [7]
(a)	M1 for attempt to collect like terms on each side leading to $ax > b$, or $ax < b$, or $ax = b$ Must have a or b correct so eg $3x > 4$ scores M0	
(b)	1 st M1 for an attempt to factorize or solve to find critical values. Method must potentially give 2 critical values 1 st A1 for $-\frac{3}{2}$ and 4 seen. They may write $x < -\frac{3}{2}$, $x < 4$ and still get this A1 2 nd M1 for choosing the "inside region" for their critical values 2 nd A1ft follow through their 2 distinct critical values Allow $x > -\frac{3}{2}$ with "or" " , " \cup " " $x < 4$ to score M1A0 but "and" or " \cap " score M1A1 $x \in (-\frac{3}{2}, 4)$ is M1A1 but $x \in [-\frac{3}{2}, 4]$ is M1A0. Score M0A0 for a number line or graph only	
(c)	B1ft Allow if a correct answer is seen or follow through their answer to (a) and their answer to (b) but their answers to (a) and (b) <u>must be regions</u> . Do not follow through single values. If their follow through answer is the empty set accept \emptyset or $\{\}$ or equivalent in words If (a) or (b) are not given then score this mark for cao NB You may see $x < 4$ (with anything or nothing in-between) $x < -1.5$ in (b) and empty set in (c) for B1ft Do not award marks for part (b) if only seen in part (c) Use of \leq instead of $<$ (or \geq instead of $>$) loses one accuracy mark only, at first occurrence.	




Question Number	Scheme	Marks
Q5 (a)	$a + 9d = 2400 \quad a + 39d = 600$ $d = \frac{-1800}{30} \quad d = -60 \quad (\text{accept } \pm 60 \text{ for A1})$	M1 M1 A1 (3)
(b)	$a - 540 = 2400 \quad a = 2940$	M1 A1 (2)
(c)	$\text{Total} = \frac{1}{2}n\{2a + (n-1)d\} = \frac{1}{2} \times 40 \times (5880 + 39 \times -60) \quad (\text{ft values of } a \text{ and } d)$ $= \underline{70\,800}$	M1 A1ft A1cao (3) [8]
	<p><u>Note:</u> If the sequence is considered 'backwards', an equivalent solution may be given using $d = 60$ with $a = 600$ and $l = 2940$ for part (b). This can still score full marks. Ignore labelling of (a) and (b)</p> <p>(a) 1st M1 for an attempt to use 2400 and 600 in $a + (n-1)d$ formula. Must use both values i.e. need $a + pd = 2400$ and $a + qd = 600$ where $p = 8$ or 9 and $q = 38$ or 39 (any combination) 2nd M1 for an attempt to solve <u>their</u> 2 linear equations in a and d as far as $d = \dots$ A1 for $d = \pm 60$. Condone correct equations leading to $d = 60$ or $a + 8d = 2400$ and $a + 38d = 600$ leading to $d = -60$. They should get penalised in (b) and (c). NB This is a "one off" ruling for A1. Usually an A mark must follow from their work. ALT 1st M1 for $(30d) = \pm (2400 - 600)$ 2nd M1 for $(d =) \pm \frac{(2400 - 600)}{30}$ A1 for $d = \pm 60$ $a + 9d = 600, a + 39d = 2400$ only scores M0 BUT if they solve to find $d = \pm 60$ then use ALT scheme above.</p> <p>(b) M1 for use of <u>their</u> d in a correct linear equation to find a leading to $a = \dots$ A1 their a must be compatible with their d so $d = 60$ must have $a = 600$ and $d = -60, a = 2940$ So for example they can have $2400 = a + 9(60)$ leading to $a = \dots$ for M1 but it scores A0 Any approach using a list scores M1A1 for a correct a but M0A0 otherwise</p> <p>(c) M1 for use of a correct S_n formula with $n = 40$ and at least one of a, d or l correct or correct ft. 1st A1ft for use of a correct S_{40} formula and both a, d or a, l correct or correct follow through ALT $\text{Total} = \frac{1}{2}n\{a + l\} = \frac{1}{2} \times 40 \times (2940 + 600) \quad (\text{ft value of } a) \quad \text{M1 A1ft}$ 2nd A1 for 70800 only</p>	

Question Number	Scheme	Marks
Q6	$b^2 - 4ac$ attempted, in terms of p . $(3p)^2 - 4p = 0$ o.e. Attempt to solve for p e.g. $p(9p - 4) = 0$ Must potentially lead to $p = k, k \neq 0$ $p = \frac{4}{9}$ (Ignore $p = 0$, if seen)	M1 A1 M1 A1cso [4]
	<p>1st M1 for an attempt to substitute into $b^2 - 4ac$ or $b^2 = 4ac$ with b or c correct Condone x's in one term only. This can be inside a square root as part of the quadratic formula for example. Use of inequalities can score the M marks only</p> <p>1st A1 for any correct equation: $(3p)^2 - 4 \times 1 \times p = 0$ or better</p> <p>2nd M1 for an attempt to factorize or solve their quadratic expression in p. Method must be sufficient to lead to their $p = \frac{4}{9}$.</p> <p>Accept factors or use of quadratic formula or $(p \pm \frac{2}{9})^2 = k^2$ (o.e. eg) $(3p \pm \frac{2}{3})^2 = k^2$ or equivalent work on <u>their</u> eqn.</p> $9p^2 = 4p \Rightarrow \frac{9p^2}{p} = 4 \text{ which would lead to } 9p = 4 \text{ is OK for this 2nd M1}$ <p>ALT <u>Comparing coefficients</u></p> <p>M1 for $(x + \alpha)^2 = x^2 + \alpha^2 + 2\alpha x$ and A1 for a correct equation eg $3p = 2\sqrt{p}$</p> <p>M1 for forming solving leading to $\sqrt{p} = \frac{2}{3}$ or better</p> <p><u>Use of quadratic/discriminant formula (or any formula) Rule for awarding M mark</u> If the formula is quoted accept some correct substitution leading to a partially correct expression. If the formula is not quoted only award for a fully correct expression using their values.</p>	

Question Number	Scheme	Marks
Q7	<p>(a) $(a_2 =) 2k - 7$</p> <p>(b) $(a_3 =) 2(2k - 7) - 7$ or $4k - 14 - 7, = 4k - 21$ (*)</p> <p>(c) $(a_4 =) 2(4k - 21) - 7 (= 8k - 49)$</p> $\sum_{r=1}^4 a_r = k + "(2k - 7)" + (4k - 21) + "(8k - 49)"$ $k + (2k - 7) + (4k - 21) + (8k - 49) = 15k - 77 = 43 \quad k = 8$	<p>B1 (1)</p> <p>M1, A1cso (2)</p> <p>M1</p> <p>M1</p> <p>M1 A1 (4)</p> <p>[7]</p>
	<p>(b) M1 must see $2(\text{their } a_2) - 7$ or $2(2k - 7) - 7$ or $4k - 14 - 7$. Their a_2 must be a function of k. A1cso must see the $2(2k - 7) - 7$ or $4k - 14 - 7$ expression and the $4k - 21$ with no incorrect working</p> <p>(c) 1st M1 for an attempt to find a_4 using the given rule. Can be awarded for $8k - 49$ seen. Use of formulae for the sum of an arithmetic series scores M0M0A0 for the next 3 marks. 2nd M1 for attempting the sum of the 1st 4 terms. Must have "+" not just , or clear attempt to sum. Follow through their a_2 and a_4 provided they are linear functions of k. Must lead to linear expression in k. Condone use of their linear $a_3 \neq 4k - 21$ here too. 3rd M1 for forming a linear equation in k using their sum and the 43 and attempt to solve for k as far as $pk = q$ A1 for $k = 8$ only so $k = \frac{120}{15}$ is A0</p> <p><u>Answer Only</u> (e.g. trial improvement) Accept $k = 8$ <u>only if</u> $8 + 9 + 11 + 15 = 43$ is seen as well</p> <p><u>Sum</u> $a_2 + a_3 + a_4 + a_5$ or $a_2 + a_3 + a_4$ Allow: M1 if $8k - 49$ is seen, M0 for the sum (since they are not adding the 1st 4 terms) then M1 if they use their sum along with the 43 to form a linear equation and attempt to solve but A0</p>	

Question Number	Scheme	Marks
Q8 (a)	$AB: m = \frac{2-7}{8-6}, \left(= -\frac{5}{2} \right)$ Using $m_1 m_2 = -1: m_2 = \frac{2}{5}$ $y - 7 = \frac{2}{5}(x - 6), \quad 2x - 5y + 23 = 0 \quad (\text{o.e. with integer coefficients})$	B1 M1 M1, A1 (4)
(b)	Using $x = 0$ in the answer to (a), $y = \frac{23}{5}$ or 4.6	M1, A1ft (2)
(c)	Area of triangle = $\frac{1}{2} \times 8 \times \frac{23}{5} = \frac{92}{5}$ (o.e.) e.g. $\left(18\frac{2}{5}, 18.4, \frac{184}{10} \right)$	M1 A1 (2)
		[8]
(a)	B1 for an expression for the gradient of AB . Does not need the $= -2.5$ 1 st M1 for use of the perpendicular gradient rule. Follow through their m 2 nd M1 for the use of (6, 7) and their changed gradient to form an equation for l . Can be awarded for $\frac{y-7}{x-6} = \frac{2}{5}$ o.e. Alternative is to use (6, 7) in $y = mx + c$ to <u>find a value</u> for c . Score when $c = \dots$ is reached.	
	A1 for a correct equation in the required form and must have “= 0” and integer coefficients	
(b)	M1 for using $x = 0$ in their answer to part (a) e.g. $-5y + 23 = 0$ A1ft for $y = \frac{23}{5}$ provided that $x = 0$ clearly seen <u>or</u> $C(0, 4.6)$. Follow through their equation in (a) If $x=0, y = 4.6$ are clearly seen but C is given as (4.6,0) apply ISW and award the mark. This A mark requires a simplified fraction or an exact decimal Accept their 4.6 marked on diagram next to C for M1A1ft	
(c)	M1 for $\frac{1}{2} \times 8 \times y_C$ so can follow through their y coordinate of C . A1 for 18.4 (o.e.) but their y coordinate of C must be positive	
	<u>Use of 2 triangles or trapezium and triangle</u> Award M1 when an expression for area of OCB only is seen <u>Determinant approach</u> Award M1 when an expression containing $\frac{1}{2} \times 8 \times y_C$ is seen	

Question Number	Scheme	Marks
Q9 (a)	$\left[(3-4\sqrt{x})^2 = \right] 9-12\sqrt{x}-12\sqrt{x}+(-4)^2 x$ $9x^{-\frac{1}{2}}+16x^{\frac{1}{2}}-24$	M1 A1, A1 (3)
(b)	$f'(x) = -\frac{9}{2}x^{-\frac{3}{2}}, +\frac{16}{2}x^{-\frac{1}{2}}$	M1 A1, A1ft (3)
(c)	$f'(9) = -\frac{9}{2} \times \frac{1}{27} + \frac{16}{2} \times \frac{1}{3} = -\frac{1}{6} + \frac{16}{6} = \frac{5}{2}$	M1 A1 (2)
		[8]
(a)	<p>M1 for an attempt to expand $(3-4\sqrt{x})^2$ with at least 3 terms correct- as printed or better</p> <p>Or $9-k\sqrt{x}+16x$ ($k \neq 0$) . See also the MR rule below</p> <p>1st A1 for their coefficient of $\sqrt{x} = 16$. Condone writing $(\pm)9x^{(\pm)\frac{1}{2}}$ instead of $9x^{-\frac{1}{2}}$</p> <p>2nd A1 for $B = -24$ or their constant term = -24</p>	
(b)	<p>M1 for an attempt to differentiate an x term $x^n \rightarrow x^{n-1}$</p> <p>1st A1 for $-\frac{9}{2}x^{-\frac{3}{2}}$ <u>and</u> their constant B differentiated to zero. NB $-\frac{1}{2} \times 9x^{-\frac{3}{2}}$ is A0</p> <p>2nd A1ft follow through their $Ax^{\frac{1}{2}}$ but can be scored without a value for A, i.e. for $\frac{A}{2}x^{-\frac{1}{2}}$</p>	
(c)	<p>M1 for some correct substitution of $x = 9$ in <u>their</u> expression for $f'(x)$ including an attempt at $(9)^{\pm\frac{k}{2}}$ (k odd) somewhere that leads to some appropriate multiples of $\frac{1}{3}$ or 3</p> <p>A1 accept $\frac{15}{6}$ or any exact equivalent of 2.5 e.g. $\frac{45}{18}, \frac{135}{54}$ or even $\frac{67.5}{27}$</p> <p><u>Misread (MR)</u> Only allow MR of the form $\frac{(3-k\sqrt{x})^2}{\sqrt{x}}$ N.B. Leads to answer in (c) of $\frac{k^2-1}{6}$</p> <p>Score as M1A0A0, M1A1A1ft, M1A1ft</p>	

Question Number	Scheme	Marks
Q10 (a)	$x(x^2 - 6x + 9)$ $= x(x - 3)(x - 3)$	B1 M1 A1 (3) B1
(b)	 <p>Shape </p> <p><u>Through</u> origin (<u>not</u> touching) Touching x-axis only once Touching at (3, 0), or 3 on x-axis [Must be on graph not in a table]</p>	B1 B1 B1ft (4)
(c)	 <p>Moved horizontally (either way) (2, 0) and (5, 0), or 2 and 5 on x-axis</p>	M1 A1 (2)
		[9]
(a)	<p>B1 for correctly taking out a factor of x</p> <p>M1 for an attempt to factorize their 3TQ e.g. $(x + p)(x + q)$ where $pq = 9$. So $(x - 3)(x + 3)$ will score M1 but A0</p> <p>A1 for a fully correct factorized expression - accept $x(x - 3)^2$ If they “solve” use ISW If the only correct linear factor is $(x - 3)$, perhaps from factor theorem, award B0M1A0 Do not award marks for factorising in part (b)</p> <p>For the graphs “Sharp points” will lose the 1st B1 in (b) but otherwise be generous on shape Condone (0, 3) in (b) and (0, 2), (0,5) in (c) if the points are marked in the correct places.</p>	
S.C.		
(b)	<p>2nd B1 for a curve that starts or terminates at (0, 0) score B0</p> <p>4th B1ft for a curve that touches (not crossing or terminating) at $(a, 0)$ where their $y = x(x - a)^2$</p>	
(c)	<p>M1 for their graph moved horizontally (only) <u>or</u> a fully correct graph Condone a partial stretch if ignoring their values looks like a simple translation</p> <p>A1 for their graph translated 2 to the right <u>and</u> crossing or touching the axis at 2 and 5 only</p> <p>Allow a fully correct graph (as shown above) to score M1A1 whatever they have in (b)</p>	

Question Number	Scheme	Marks
Q11 (a) (b) (c)	$x = 2: y = 8 - 8 - 2 + 9 = 7 \quad (*)$ $\frac{dy}{dx} = 3x^2 - 4x - 1$ $x = 2: \frac{dy}{dx} = 12 - 8 - 1 (= 3)$ $y - 7 = 3(x - 2), \quad \underline{y = 3x + 1}$ $m = -\frac{1}{3} \quad \left(\text{for } -\frac{1}{m} \text{ with their } m\right)$ $3x^2 - 4x - 1 = -\frac{1}{3}, \quad 9x^2 - 12x - 2 = 0 \quad \text{or} \quad x^2 - \frac{4}{3}x - \frac{2}{9} = 0 \quad (\text{o.e.})$ $\left(x = \frac{12 + \sqrt{144 + 72}}{18}\right) (\sqrt{216} = \sqrt{36} \sqrt{6} = 6\sqrt{6}) \text{ or } (3x - 2)^2 = 6 \rightarrow 3x = 2 \pm \sqrt{6}$ $x = \frac{1}{3}(2 + \sqrt{6}) \quad (*)$	B1 (1) M1 A1 A1ft M1, <u>A1</u> (5) B1ft M1, A1 M1 A1cso (5) [11]
(a) (b) (c) ALT	B1 there must be a clear attempt to substitute $x = 2$ leading to 7 e.g. $2^3 - 2 \times 2^2 - 2 + 9 = 7$ 1 st M1 for an attempt to differentiate with at least one of the given terms fully correct. 1 st A1 for a fully correct expression 2 nd A1ft for sub. $x = 2$ in <u>their</u> $\frac{dy}{dx} (\neq y)$ accept for a correct expression e.g. $3 \times (2)^2 - 4 \times 2 - 1$ 2 nd M1 for use of their “3” (provided it comes from their $\frac{dy}{dx} (\neq y)$ and $x=2$) to find equation of tangent. Alternative is to use (2, 7) in $y = mx + c$ to <u>find a value</u> for c . Award when $c = \dots$ is seen. No attempted use of $\frac{dy}{dx}$ in (b) scores 0/5 1 st M1 for forming an equation from their $\frac{dy}{dx} (\neq y)$ and their $-\frac{1}{m}$ (must be changed from m) 1 st A1 for a correct 3TQ all terms on LHS (condone missing =0) 2 nd M1 for proceeding to $x = \dots$ or $3x = \dots$ by formula or completing the square for a 3TQ. Not factorising. Condone \pm 2 nd A1 for proceeding to given answer with no incorrect working seen. Can still have \pm . <u>Verify (for M1A1M1A1)</u> 1 st M1 for attempting to square need ≥ 3 correct values in $\frac{4+6+4\sqrt{6}}{9}$, 1 st A1 for $\frac{10+4\sqrt{6}}{9}$ 2 nd M1 Dependent on 1 st M1 in this case for substituting in all terms of their $\frac{dy}{dx}$ 2 nd A1cso for cso <u>with a full comment</u> e.g. “the x co-ord of Q is ...”	

