

GCE Examinations  
Advanced Subsidiary

# Core Mathematics C1

Paper K

## MARKING GUIDE

This guide is intended to be as helpful as possible to teachers by providing concise solutions and indicating how marks could be awarded. There are obviously alternative methods that would also gain full marks.

Method marks (M) are awarded for knowing and using a method.

Accuracy marks (A) can only be awarded when a correct method has been used.

(B) marks are independent of method marks.



*Written by Shaun Armstrong*

© *Solomon Press*

*These sheets may be copied for use solely by the purchaser's institute.*

## C1 Paper K – Marking Guide

<p>1. <math>(2^2)^{y+3} = 2^3</math>  <math>2y + 6 = 3</math>  <math>y = -\frac{3}{2}</math></p>	<p>M1 M1 A1</p>	<p>(3)</p>
--	-------------------------	------------


---

<p>2. <math>= \int (3x^2 + \frac{1}{2}x^{-2}) dx</math>  <math>= x^3 - \frac{1}{2}x^{-1} + c</math></p>	<p>B1 M1 A2</p>	<p>(4)</p>
---	---------------------	------------

---

<p>3. <math>\frac{EH}{AD} = \frac{EF}{AB} \therefore \frac{EH}{\sqrt{5}} = \frac{1+\sqrt{5}}{3-\sqrt{5}}</math>  <math>\frac{1+\sqrt{5}}{3-\sqrt{5}} = \frac{1+\sqrt{5}}{3-\sqrt{5}} \times \frac{3+\sqrt{5}}{3+\sqrt{5}} = \frac{3+\sqrt{5}+3\sqrt{5}+5}{9-5} = 2 + \sqrt{5}</math>  <math>\therefore EH = \sqrt{5}(2 + \sqrt{5}) = 5 + 2\sqrt{5}</math></p>	<p>M1 M2 A1 M1 A1</p>	<p>(6)</p>
---	-------------------------------	------------

---

<p>4. (a) </p>	<p>B2 B2</p>	
<p>(b) 3 solutions  <math>x^2 - 4x + \frac{1}{x} = 0 \Rightarrow x^2 - 4x = -\frac{1}{x}</math>          and the graphs of <math>y = x^2 - 4x</math> and <math>y = -\frac{1}{x}</math> intersect at 3 points</p>	<p>B1 B1</p>	<p>(6)</p>

---

<p>5. (a) <math>(x+k)^2 - k^2 + 4 = 0</math>  <math>(x+k)^2 = k^2 - 4</math>  <math>x+k = \pm\sqrt{k^2-4}</math>  <math>x = -k \pm \sqrt{k^2-4}</math></p>	<p>M1 A1 M1 A1</p>	
<p>(b) <math>k = 3 \therefore x = -3 \pm \sqrt{3^2-4}</math>  <math>= -3 \pm \sqrt{5}</math></p>	<p>M1 A1</p>	<p>(6)</p>

---

<p>6. (a) AP: <math>a = 77, l = -70</math>  <math>S_{50} = \frac{50}{2}[77 + (-70)] = 25 \times 7 = 175</math></p>	<p>B1 M1 A1</p>	
<p>(b) AP: <math>a = 2, d = \frac{1}{2}</math>  <math>S_n = \frac{n}{2}[4 + \frac{1}{2}(n-1)]</math>  <math>= \frac{1}{4}n[8 + (n-1)] = \frac{1}{4}n(n+7) \quad [k = \frac{1}{4}]</math></p>	<p>B2 M1 A1</p>	<p>(7)</p>

---

<p>7. <math>x - 3y + 7 = 0 \Rightarrow x = 3y - 7</math>          sub. into <math>x^2 + 2xy - y^2 = 7</math>  <math>(3y-7)^2 + 2y(3y-7) - y^2 = 7</math>  <math>y^2 - 4y + 3 = 0</math>  <math>(y-1)(y-3) = 0</math>  <math>y = 1, 3</math>  <math>\therefore x = -4, y = 1</math> or <math>x = 2, y = 3</math></p>	<p>M1 M1 A1 M1 A1 M1 A1</p>	<p>(7)</p>
---	---	------------

---

8.	(a)	$\frac{dy}{dx} = 1 - 4x^{-3}$	B1
		$\frac{d^2y}{dx^2} = 12x^{-4}$	M1 A1
	(b)	$y = \int (1 - 4x^{-3}) dx$	
		$y = x + 2x^{-2} + c$	M1 A2
		$x = -1, y = 0 \therefore 0 = -1 + 2 + c$	
		$c = -1$	M1
		$y = x + 2x^{-2} - 1$	
		when $x = 2, y = 2 + \frac{1}{2} - 1 = \frac{3}{2}$	M1 A1 (9)

9.	(a)	$y = x - 6\sqrt{x} + 9$	M1 A1
		$\frac{dy}{dx} = 1 - 3x^{-\frac{1}{2}} = 1 - \frac{3}{\sqrt{x}}$	M1 A1
	(b)	$x = 4 \therefore y = 1$	
		grad of tangent $= 1 - \frac{3}{2} = -\frac{1}{2}$	M1
		grad of normal $= \frac{-1}{-\frac{1}{2}} = 2$	M1 A1
		$\therefore y - 1 = 2(x - 4)$	M1
		$y = 2x - 7$	A1
	(c)	at intersect: $x - 6\sqrt{x} + 9 = 2x - 7$	
		$x + 6\sqrt{x} - 16 = 0$	M1
		$(\sqrt{x} + 8)(\sqrt{x} - 2) = 0$	M1
		$\sqrt{x} = -8, 2$	A1
		$\sqrt{x} = 2 \Rightarrow x = 4$ (at P)	
		$\sqrt{x} = -8 \Rightarrow$ no real solutions $\therefore$ normal does not intersect again	A1 (13)

10.	(a)	$y - 4 = 3(x + 6)$	M1
		$y = 3x + 22$	A1
	(b)	at B, $x = 0 \therefore y = 2 \Rightarrow B(0, 2)$	B1
		at C, $x - 7(3x + 22) + 14 = 0$	M1
		$x = -7$	A1
		$\therefore C(-7, 1)$	A1
	(c)	grad AB $= \frac{2-4}{0-(-6)} = -\frac{1}{3}$	M1 A1
		grad AC $= \frac{1-4}{-7-(-6)} = 3$	
		grad AB $\times$ grad AC $= -\frac{1}{3} \times 3 = -1$	M1
		$\therefore AB$ perp to $AC \therefore \angle BAC = 90^\circ$	A1
	(d)	$AB = \sqrt{(0+6)^2 + (2-4)^2} = \sqrt{36+4} = \sqrt{40} = 2\sqrt{10}$	M1 A1
		$AC = \sqrt{(-7+6)^2 + (1-4)^2} = \sqrt{1+9} = \sqrt{10}$	
		area $= \frac{1}{2} \times 2\sqrt{10} \times \sqrt{10} = 10$	M1 A1 (14)

Total (75)

## Performance Record – C1 Paper K

Question no.	1	2	3	4	5	6	7	8	9	10	Total
Topic(s)	indices	integr.	surds	curve sketch	compl. square	AP	simul. eqn	diff., integr.	diff., normal	straight lines	
Marks	3	4	6	6	6	7	7	9	13	14	75
Student											