1 Find an equation for the tangent to the curve with equation

$$y = (3-x)^{\frac{3}{2}}$$

at the point on the curve with x-coordinate -1.

(4)

**(2)** 

**2** a Sketch the curve with equation  $y = 3 - \ln 2x$ .

- (2)
- **b** Find the exact coordinates of the point where the curve crosses the *x*-axis.
- c Find an equation for the tangent to the curve at the point on the curve where x = 5. (4)

This tangent cuts the x-axis at A and the y-axis at B.

- **d** Show that the area of triangle OAB, where O is the origin, is approximately 7.20 (3)
- 3 Differentiate with respect to x

**a** 
$$(3x-1)^4$$
, (2)

$$\mathbf{b} \quad \frac{x^2}{\sin 2x} \,. \tag{3}$$

4 The area of the surface of a boulder covered by lichen,  $A \text{ cm}^2$ , at time t years after initial observation, is modelled by the formula

$$A = 2e^{0.5t}$$
.

Using this model,


**a** find the area of lichen on the boulder after three years,

- (2)
- **b** find the rate at which the area of lichen is increasing per day after three years,
- (2)
- c find, to the nearest year, how long it takes until the area of lichen is 65 cm<sup>2</sup>.
- (1)

**(4)** 

**d** Explain why the model cannot be valid for large values of t.

5



The diagram shows the curve with equation  $y = a \ln x - 4x$ , where a is a positive constant. Find, in terms of a,

- a the coordinates of the stationary point on the curve, (4)
- **b** an equation for the tangent to the curve at the point where x = 1. (3)

Given that this tangent meets the x-axis at the point (3, 0),

c show that 
$$a = 6$$
.

6 Given that  $y = e^{2x} \sin x$ ,

$$\mathbf{a} \quad \text{find } \frac{\mathrm{d}y}{\mathrm{d}x}, \tag{2}$$


**b** show that 
$$\frac{d^2y}{dx^2} - 4\frac{dy}{dx} + 5y = 0.$$
 (3)

7 A curve has the equation  $x = \tan^2 y$ .

**a** Show that 
$$\frac{\mathrm{d}y}{\mathrm{d}x} = \frac{1}{2\sqrt{x}(x+1)}$$
. (5)

**b** Find an equation for the normal to the curve at the point where  $y = \frac{\pi}{4}$ . (3)

8



The diagram shows the curve  $y = \frac{x+2}{\sqrt{x-2}}$ , x > 2, which has a minimum point at P.

a Find and simplify an expression for 
$$\frac{dy}{dx}$$
. (3)

**b** Find the coordinates of 
$$P$$
. (2)

The point Q on the curve has x-coordinate 3.

**c** Show that the normal to the curve at Q has equation

$$2x - 3y + 9 = 0. (3)$$

9 A curve has the equation  $y = e^x(x-1)^2$ .

a Find 
$$\frac{dy}{dx}$$
.

**b** Show that 
$$\frac{d^2y}{dx^2} = e^x(x^2 + 2x - 1)$$
. (2)

c Find the exact coordinates of the turning points of the curve and determine their nature. (4)

**d** Show that the tangent to the curve at the point where x = 2 has the equation

$$y = e^2(3x - 5). {3}$$

10 The curve with equation  $y = \frac{1}{2}x^2 - 3 \ln x$ , x > 0, has a stationary point at A.

a Find the exact x-coordinate of 
$$A$$
. (3)

c Show that the y-coordinate of A is 
$$\frac{3}{2}(1 - \ln 3)$$
. (2)

**d** Find an equation for the tangent to the curve at the point where x = 1, giving your answer in the form ax + by + c = 0, where a, b and c are integers. (3)

11  $f(x) = \frac{6x}{(x-1)(x+2)} - \frac{2}{x-1}.$ 

a Show that 
$$f(x) = \frac{4}{x+2}$$
. (5)

**b** Find an equation for the tangent to the curve y = f(x) at the point with x-coordinate 2, giving your answer in the form ax + by = c, where a, b and c are integers. (4)