

- 1** Use the identity $\sin^2 x + \cos^2 x \equiv 1$ to obtain the identities
a $1 + \tan^2 x \equiv \sec^2 x$ **b** $1 + \cot^2 x \equiv \operatorname{cosec}^2 x$
- 2** **a** Given that $\tan A = \frac{1}{3}$, find the exact value of $\sec^2 A$.
b Given that $\operatorname{cosec} B = 1 + \sqrt{3}$, find the exact value of $\cot^2 B$.
c Given that $\sec C = \frac{3}{2}$, find the possible values of $\tan C$, giving your answers in the form $k\sqrt{5}$.
- 3** Solve each equation for θ in the interval $0 \leq \theta \leq 2\pi$ giving your answers in terms of π .
a $3 \sec^2 \theta = 4 \tan^2 \theta$ **b** $\tan^2 \theta - 2 \sec \theta + 1 = 0$
c $\cot^2 \theta - 3 \operatorname{cosec} \theta + 3 = 0$ **d** $\operatorname{cosec}^2 \theta + \cot^2 \theta = 3$
e $\sec^2 \theta + 2 \tan \theta = 0$ **f** $\operatorname{cosec}^2 \theta - \sqrt{3} \cot \theta - 1 = 0$
- 4** Solve each equation for x in the interval $-180^\circ \leq x \leq 180^\circ$.
Give your answers to 1 decimal place where appropriate.
a $\tan^2 x - 2 \sec x - 2 = 0$ **b** $2 \operatorname{cosec}^2 x + 2 = 9 \cot x$
c $\operatorname{cosec}^2 x + 5 \operatorname{cosec} x + 2 \cot^2 x = 0$ **d** $3 \tan^2 x - 3 \tan x + \sec^2 x = 2$
e $\tan^2 x + 4 \sec x - 2 = 0$ **f** $2 \cot^2 x + 3 \operatorname{cosec}^2 x = 4 \cot x + 3$
- 5** Solve each equation for x in the interval $0 \leq x \leq 360^\circ$.
a $\cot^2 2x + \operatorname{cosec} 2x - 1 = 0$ **b** $8 \sin^2 x + \sec x = 8$
c $3 \operatorname{cosec}^2 x - 4 \sin^2 x = 1$ **d** $9 \sec^2 x - 8 = \operatorname{cosec}^2 x$
- 6** Prove each of the following identities.
a $\operatorname{cosec}^2 x - \sec^2 x \equiv \cot^2 x - \tan^2 x$ **b** $(\cot x - 1)^2 \equiv \operatorname{cosec}^2 x - 2 \cot x$
c $(\cos x - 2 \sec x)^2 \equiv \cos^2 x + 4 \tan^2 x$ **d** $\sec^2 x - \sin^2 x \equiv \tan^2 x + \cos^2 x$
e $(\tan x + \cot x)^2 \equiv \sec^2 x + \operatorname{cosec}^2 x$ **f** $(\sin x - \sec x)^2 \equiv \sin^2 x + (\tan x - 1)^2$
g $\sec^2 x + \operatorname{cosec}^2 x \equiv \sec^2 x \operatorname{cosec}^2 x$ **h** $\sec^4 x + \tan^4 x \equiv 2 \sec^2 x \tan^2 x + 1$
- 7** Prove that there are no real values of x for which
$$4 \sec^2 x - \sec x + 2 \tan^2 x = 0.$$
- 8** **a** Prove the identity
$$\operatorname{cosec} x \sec x - \cot x \equiv \tan x.$$

b Hence, or otherwise, find the values of x in the interval $0 \leq x \leq 360^\circ$ for which
$$\operatorname{cosec} x \sec x = 3 + \cot x,$$

giving your answers to 1 decimal place.