Centre No.			Paper Reference			Surname	Initial(s)				
Candidate No.			6	6	6	5	/	0	1	Signature	

6665/01

Edexcel GCE

Core Mathematics C3 Advanced Level

Monday 12 June 2006 – Afternoon

Time: 1 hour 30 minutes

Materials	required	for	examination

Mathematical Formulae (Green)

Items included with question papers

Candidates may use any calculator EXCEPT those with the facility for symbolic algebra, differentiation and/or integration. Thus candidates may NOT use calculators such as the Texas Instruments TI 89, TI 92, Casio CFX 9970G, Hewlett Packard HP 48G.

Instructions to Candidates

In the boxes above, write your centre number, candidate number, your surname, initial(s) and signature.

Check that you have the correct question paper.

When a calculator is used, the answer should be given to an appropriate degree of accuracy.

You must write your answer for each question in the space following the question.

If you need more space to complete your answer to any question, use additional answer sheets.

Information for Candidates

A booklet 'Mathematical Formulae and Statistical Tables' is provided.

The marks for individual questions and the parts of questions are shown in round brackets: e.g. (2). Full marks may be obtained for answers to ALL questions.

There are 8 questions in this question paper. The total mark for this paper is 75.

There are 24 pages in this question paper. Any blank pages are indicated.

Advice to Candidates

You must ensure that your answers to parts of questions are clearly labelled.

You must show sufficient working to make your methods clear to the Examiner. Answers without working may gain no credit.

This publication may be reproduced only in accordance with cel Limited copyright policy.

W850/R6665/57570 3/3/3/3/3/36,300





Turn over

Total

Examiner's use only

Team Leader's use only

Ouestion

1

2

3

4

5

6

7

8

Leave



1	(a)	Simplify	$\frac{3x^2 - x - 2}{x^2 + 1}$
1.	(a)	Simping	$x^2 - 1$

(3)

(h)	Hence or otherwise evaress	$3x^2 - x - 2$	_ 1	as a single fraction in its simplest
(0)	Trenee, or otherwise, express	$x^{2}-1$	x(x+1)	as a single fraction in its simplest
	form			

(3)

2.	Differentiate,	with	respect	to	x.

(a)
$$e^{3x} + \ln 2x$$
,

(3)

(b)
$$(5+x^2)^{\frac{3}{2}}$$
.

(3)

(Total 6 marks)

Q2

Leave blank

3.




Figure 1

Figure 1 shows part of the curve with equation y = f(x), $x \in \mathbb{R}$, where f is an increasing function of x. The curve passes through the points P(0, -2) and Q(3, 0) as shown.

In separate diagrams, sketch the curve with equation

(a)
$$y = |f(x)|$$
, (3)

(b)
$$y = f^{-1}(x)$$
, (3)

(c)
$$y = \frac{1}{2} f(3x)$$
. (3)

Indicate clearly on each sketch the coordinates of the points at which the curve crosses or meets the axes.

	Leave blank
Question 3 continued	

4.	A heated metal ball is dropped into a liquid.	As the ball cools, its temperature, $T^{\circ}C$,
	t minutes after it enters the liquid, is given by	

$$T = 400 e^{-0.05t} + 25, \quad t \geqslant 0.$$

(a) Find the temperature of the ball as it enters the liquid.

(1)

(b) Find the value of t for which T = 300, giving your answer to 3 significant figures.

(4)

(c) Find the rate at which the temperature of the ball is decreasing at the instant when t = 50. Give your answer in °C per minute to 3 significant figures.

(3)

(d) From the equation for temperature T in terms of t, given above, explain why the temperature of the ball can never fall to 20 °C.

(1)

5.






Figure 2 shows part of the curve with equation

$$y = (2x-1)\tan 2x, \ \ 0 \le x < \frac{\pi}{4}.$$

The curve has a minimum at the point P. The x-coordinate of P is k.

(a) Show that k satisfies the equation

$$4k + \sin 4k - 2 = 0.$$

(6)

The iterative formula

$$x_{n+1} = \frac{1}{4}(2 - \sin 4x_n), \ x_0 = 0.3,$$

is used to find an approximate value for k.

(b) Calculate the values of x_1 , x_2 , x_3 and x_4 , giving your answers to 4 decimal places.

(3)

(c) Show that k = 0.277, correct to 3 significant figures.

(2)



6	(0)	Using $\sin^2\theta + \cos^2\theta = 1$	show that $\cos^2\theta = \cot^2\theta - 1$
0.	(a)	Using $\sin^2\theta + \cos^2\theta \equiv 1$,	show that $\csc^2 \theta - \cot^2 \theta \equiv 1$.

(2)

(b) Hence, or otherwise, prove that

$$\csc^4\theta - \cot^4\theta = \csc^2\theta + \cot^2\theta.$$

(2)

(c) Solve, for $90^{\circ} < \theta < 180^{\circ}$,

$$\csc^4\theta - \cot^4\theta = 2 - \cot \theta$$
.

(6)

7. For the constant k, where k > 1, the functions f and g are defined by

f:
$$x \mapsto \ln(x+k)$$
, $x > -k$,
g: $x \mapsto |2x-k|$, $x \in \mathbb{R}$.

(a) On separate axes, sketch the graph of f and the graph of g.

On each sketch state, in terms of k, the coordinates of points where the graph meets the coordinate axes.

(5)

(b) Write down the range of f.

(1)

(c) Find $fg\left(\frac{k}{4}\right)$ in terms of k, giving your answer in its simplest form.

(2)

The curve C has equation y = f(x). The tangent to C at the point with x-coordinate 3 is parallel to the line with equation 9y = 2x + 1.

(d) Find the value of k.

(4)



Leave blank

- **8.** (a) Given that $\cos A = \frac{3}{4}$, where $270^{\circ} < A < 360^{\circ}$, find the exact value of $\sin 2A$. (5)
 - (b) (i) Show that $\cos\left(2x + \frac{\pi}{3}\right) + \cos\left(2x \frac{\pi}{3}\right) \equiv \cos 2x$.

Given that

$$y = 3\sin^2 x + \cos\left(2x + \frac{\pi}{3}\right) + \cos\left(2x - \frac{\pi}{3}\right),$$

(ii) show that $\frac{dy}{dx} = \sin 2x$.

(4)

TOTAL FOR PAPER: 75	MARKS
	2 marks)
	Q