edexcel
advancing learning, changing lives

Mark Scheme (Results)
Summer 2007

GCE

GCE Mathematics

Core Mathematics C3 (6665)

Edexcel Limited. Registered in England and Wales No. 4496750
Registered Office: One90 High Holborn, London WC1V 7BH

A PEARSON COMPANY

June 2007
6665 Core Mathematics C3
Mark Scheme

Question Number		Scheme	Marks
1.	(*a*)	$\ln 3x = \ln 6$ or $\ln x = \ln\left(\frac{6}{3}\right)$ [implied by 0.69…] or $\ln\left(\frac{3x}{6}\right) = 0$	M1
		$x = 2$ (only this answer)	A1 (cso) (2)
	(*b*)	$(e^x)^2 - 4e^x + 3 = 0$ (any 3 term form)	M1
		$(e^x - 3)(e^x - 1) = 0$	
		$e^x = 3$ or $e^x = 1$ Solving quadratic	M1 dep
		$x = \ln 3$, $x = 0$ (or ln 1)	M1 A1 (4)
			(6 marks)

Notes: (a) Answer $x = 2$ with no working or no incorrect working seen: M1A1

Beware $x = 2$ from $\ln x = \frac{\ln 6}{\ln 3} = \ln 2$ M0A0

$\ln x = \ln 6 - \ln 3 \Rightarrow x = e^{(\ln 6 - \ln 3)}$ allow M1, $x = 2$ (no wrong working) A1

(b) 1st M1 for attempting to multiply through by e^x : Allow y, X , even x, for e^x

Be generous for M1 e.g $e^{2x} + 3 = 4$, $e^{x^2} + 3 = 4e^x$,

$3y^2 + 1 = 12y$ (from $3e^{-x} = \frac{1}{3e^x}$), $e^x + 3 = 4e^x$

2nd M1 is for solving quadratic (may be by formula or completing the square) as far as getting two values for e^x or y or X etc

3rd M1 is for converting their answer(s) of the form $e^x = k$ to $x = \ln k$ (must be exact)

A1 is for ln3 **and** ln1 or 0 (Both required and no further solutions)

2. (*a*)	$2x^2 + 3x - 2 = (2x - 1)(x + 2)$ at any stage	B1
	$f(x) = \dfrac{(2x+3)(2x-1)-(9+2x)}{(2x-1)(x+2)}$ f.t. on error in denominator factors (need not be single fraction)	M1, A1√
	Simplifying numerator to quadratic form [$= \dfrac{4x^2+4x-3-9-2x}{(2x-1)(x+2)}$]	M1
	Correct **numerator** $= \dfrac{4x^2+2x-12}{[(2x-1)(x+2)]}$	A1
	Factorising numerator, **with** a denominator $= \dfrac{2(2x-3)(x+2)}{(2x-1)(x+2)}$ o.e. [$= \dfrac{2(2x-3)}{2x-1}$] $= \dfrac{4x-6}{2x-1}$ (❋)	M1 A1 cso (7)
Alt.(*a*)	$2x^2 + 3x - 2 = (2x - 1)(x + 2)$ at any stage B1 $f(x) = \dfrac{(2x+3)(2x^2+3x-2)-(9+2x)(x+2)}{(x+2)(2x^2+3x-2)}$ M1A1 f.t. $= \dfrac{4x^3+10x^2-8x-24}{(x+2)(2x^2+3x-2)}$ $= \dfrac{2(x+2)(2x^2+x-6)}{(x+2)(2x^2+3x-2)}$ or $\dfrac{2(2x-3)(x^2+4x+4)}{(x+2)(2x^2+3x+2)}$ o.e. Any one linear factor × quadratic factor in **numerator** M1, A1 $= \dfrac{2(x+2)(x+2)(2x-3)}{(x+2)(2x^2+3x-2)}$ o.e. M1 $= \dfrac{2(2x-3)}{2x-1}$ $\dfrac{4x-6}{2x-1}$ (❋) A1	
(*b*)	Complete method for $f'(x)$; e.g $f'(x) = \dfrac{(2x-1)\times 4-(4x-6)\times 2}{(2x-1)^2}$ o.e	M1 A1
	$= \dfrac{8}{(2x-1)^2}$ or $8(2x-1)^{-2}$	A1 (3)
	Not treating f^{-1} (for f') as misread	**(10 marks)**

Notes: (a) 1st M1 in either version is for correct method

1st A1 Allow $\dfrac{2x+3(2x-1)-(9+2x)}{(2x-1)(x+2)}$ or $\dfrac{(2x+3)(2x-1)-9+2x}{(2x-1)(x+2)}$ or $\dfrac{2x+3(2x-1)-9+2x}{(2x-1)(x+2)}$ (fractions)

2nd M1 in (main a) is for forming 3 term quadratic in **numerator**

3rd M1 is for factorising their quadratic (usual rules) ; factor of 2 need not be extracted

(❋) A1 is given answer so is cso

Alt (a) 3rd M1 is for factorising resulting quadratic

Notice that B1 likely to be scored very late but on ePen scored first

(b) SC: For M allow ± given expression or one error in product rule

Alt: Attempt at $f(x) = 2 - 4(2x-1)^{-1}$ and diff. M1; $k(2x-1)^{-2}$ A1; A1 as above

Accept $8(4x^2 - 4x + 1)^{-2}$. Differentiating original function – mark as scheme.

Question Number	Scheme	Marks
3. (*a*)	$\frac{dy}{dx} = x^2e^x + 2xe^x$	M1,A1,A1 (3)
(*b*)	If $\frac{dy}{dx} = 0$, $e^x(x^2 + 2x) = 0$ setting (*a*) = 0 $[e^x \neq 0]$ $x(x + 2) = 0$ ($x = 0$) or $x = -2$ $x = 0, y = 0$ **and** $x = -2, y = 4e^{-2}$ (= 0.54…)	M1 A1 A1 √ (3)
(*c*)	$\frac{d^2y}{dx^2} = x^2e^x + 2xe^x + 2xe^x + 2e^x$ $\left[= (x^2 + 4x + 2)e^x \right]$	M1, A1 (2)
(*d*)	$x = 0$, $\frac{d^2y}{dx^2} > 0$ (=2) $x = -2$, $\frac{d^2y}{dx^2} < 0$ [$= -2e^{-2}$ (= –0.270…)] M1: Evaluate, or state sign of, candidate's (c) for at least one of candidate's x value(s) from (b) $\therefore$ minimum $\therefore$ maximum	M1 A1 (cso) (2)
Alt.(*d*)	For M1: Evaluate, or state sign of, $\frac{dy}{dx}$ at two appropriate values – on either side of at least one of their answers from (b) or Evaluate y at two appropriate values – on either side of at least one of their answers from (b) or Sketch curve 	**(10 marks)**

Notes: (a) Generous M for attempt at $f(x)g'(x) + f'(x)g(x)$
1st A1 for one correct, 2nd A1 for the other correct.
Note that x^2e^x on its own scores no marks

(b) 1st A1 ($x = 0$) may be omitted, but for
2nd A1 both sets of coordinates needed ; f.t only on candidate's $x = -2$

(c) M1 requires complete method for candidate's (a), result may be unsimplified for A1

(d) A1 is cso; $x = 0$, min, and $x = -2$, max and no incorrect working seen.,
or (in alternative) sign of $\frac{dy}{dx}$ either side correct, or values of y appropriate to t.p.
Need only consider the quadratic, as may assume $e^x > 0$.
If all marks gained in (a) and (c), and correct x values, give M1A1 for correct statements with no working

Question Number	Scheme	Marks
4. (*a*)	$x^2(3-x)-1=0$ o.e. (e.g. $x^2(-x+3)=1$) $x=\sqrt{\dfrac{1}{3-x}}$ (✱) Note(✱), answer is given: need to see appropriate working and A1 is cso [Reverse process: Squaring and non-fractional equation M1, form f(x) A1]	M1 A1 (cso) (2)
(*b*)	$x_2 = 0.6455$, $x_3 = 0.6517$, $x_4 = 0.6526$ 1st B1 is for one correct, 2nd B1 for other two correct If all three are to greater accuracy, award B0 B1	B1; B1 (2)
(*c*)	Choose values in interval (0.6525, 0.6535) or tighter and evaluate both f(0.6525) = –0.0005 (372… f(0.6535) = 0.002 (101… At least one correct "up to bracket", i.e. -0.0005 or 0.002 **Change of sign,** $\therefore x = 0.653$ is a root (correct) to 3 d.p. Requires both correct "up to bracket" and conclusion as above	M1 A1 A1 (3) **(7 marks)**
Alt (i)	Continued iterations at least as far as x_6 M1 $x_5 = 0.6527$, $x_6 = 0.6527$, $x_7 =$ … two correct to at least 4 s.f. A1 Conclusion : Two values correct to 4 d.p., so 0.653 is root to 3 d.p. A1	
Alt (ii)	If use g(0.6525) = 0.6527..>0.6525 and g(0.6535) = 0.6528..<0.6535 M1A1 Conclusion : Both results correct, so 0.653 is root to 3 d.p. A1	
5. (*a*)	Finding g(4) = k and f(k) = …. or fg(x) = $\ln\left(\dfrac{4}{x-3}-1\right)$ [f(2) = ln(2x2 – 1) fg(4) = ln(4 – 1)] = $\ln 3$	M1 A1 (2)
(*b*)	$y=\ln(2x-1) \Rightarrow e^y = 2x-1$ or $e^x = 2y-1$ $f^{-1}(x) = \frac{1}{2}(e^x+1)$ Allow $y = \frac{1}{2}(e^x+1)$ Domain $x \in \Re$ [Allow $\Re$, all reals, $(-\infty, \infty)$] independent	M1, A1 A1 B1 (4)
(*c*)	 Shape, and x-axis should appear to be asymptote **Equation $x = 3$ needed,** may see in diagram (ignore others) Intercept (0, $\frac{2}{3}$) no other; accept y = ⅔ (0.67) or on graph	B1 B1 ind. B1 ind (3)
(*d*)	$\dfrac{2}{x-3}=3 \Rightarrow x = 3\frac{2}{3}$ or exact equiv. $\dfrac{2}{x-3}=-3$, $\Rightarrow x = 2\frac{1}{3}$ or exact equiv. Note: $2 = 3(x+3)$ or $2 = 3(-x-3)$ o.e. is M0A0	B1 M1, A1 (3)
Alt:	Squaring to quadratic ($9x^2 - 54x + 77 = 0$) **and** solving M1; B1A1	**(12 marks)**

6.	(*a*)	Complete method for *R*: e.g. $R\cos\alpha = 3$, $R\sin\alpha = 2$, $R = \sqrt{(3^2 + 2^2)}$	M1
		$R = \sqrt{13}$ or 3.61 (or more accurate)	A1
		Complete method for $\tan\alpha = \frac{2}{3}$ [Allow $\tan\alpha = \frac{3}{2}$]	M1
		$\alpha = 0.588$ (Allow 33.7°)	A1 (4)
	(*b*)	Greatest value = $\left(\sqrt{13}\right)^4 = 169$	M1, A1 (2)
	(*c*)	$\sin(x + 0.588) = \frac{1}{\sqrt{13}}$ (= 0.27735…) sin(*x* + their α) = $\frac{1}{\text{their } R}$	M1
		(*x* + 0.588) = 0.281(03… or 16.1°	A1
		(*x* + 0.588) = π – 0.28103… Must be π – their 0.281 or 180° – their 16.1°	M1
		or (*x* + 0.588) = 2π + 0.28103… Must be 2π + their 0.281 or 360° + their 16.1°	M1
		x = 2.273 or *x* = 5.976 (awrt) Both (radians only)	A1 (5)
		If 0.281 or 16.1° not seen, correct answers imply this A mark	**(11 marks)**

Notes: (a) 1st M1 on Epen for correct method for R, even if found second
2nd M1 for correct method for tan α
No working at all: M1A1 for √13, M1A1 for 0.588 or 33.7°.
N.B. Rcos α = 2, Rsin α = 3 used, can still score M1A1 for R, but loses the A mark for α.
cosα = 3, sin α = 2: apply the same marking.

(b) M1 for realising sin(x + α) = ±1, so finding R^4.

(c) Working in mixed degrees/rads : first two marks available
Working consistently in degrees: Possible to score first 4 marks
[Degree answers, just for reference, Only are 130.2° and 342.4°]
Third M1 can be gained for candidate's 0.281 – candidate's 0.588 + 2π or equiv. in degrees
One of the answers correct in radians or degrees implies the corresponding M mark.

Alt: (c) (i) Squaring to form quadratic in sin *x* or cos *x* M1
[$13\cos^2 x - 4\cos x - 8 = 0$, $13\sin^2 x - 6\sin x - 3 = 0$]
Correct values for cos *x* = 0.953… , –0.646; or sin *x* = 0.767, 2.27 awrt A1
For any one value of cos *x* or sin*x*, correct method for two values of *x* M1
x = 2.273 or *x* = 5.976 (awrt) Both seen anywhere A1
Checking other values (0.307, 4.011 or 0.869, 3.449) and discarding M1

(ii) Squaring and forming equation of form *a* cos2*x* + *b*sin2*x* = *c*
$9\sin^2 x + 4\cos^2 x + 12\sin 2x = 1 \Rightarrow 12\sin 2x + 5\cos 2x = 11$
Setting up to solve using R formula e.g. √13 cos(2*x* – 1.176) = 11 M1
$(2x - 1.176) = \cos^{-1}\left(\frac{11}{\sqrt{13}}\right) = 0.562(0...$ (α) A1
(2*x* – 1.176) = $2\pi - \alpha$, $2\pi + \alpha$,..... M1
x = 2.273 or *x* = 5.976 (awrt) Both seen anywhere A1
Checking other values and discarding M1

Question Number	Scheme	Marks
7. (*a*)	$\frac{\sin\theta}{\cos\theta}+\frac{\cos\theta}{\sin\theta}=\frac{\sin^2\theta+\cos^2\theta}{\cos\theta\sin\theta}$	M1
	M1 Use of common denominator to obtain single fraction	
	$=\frac{1}{\cos\theta\sin\theta}$	M1
	M1 Use of appropriate trig identity (in this case $\sin^2\theta+\cos^2\theta=1$)	
	$=\frac{1}{\frac{1}{2}\sin 2\theta}$ Use of $\sin 2\theta=2\sin\theta\cos\theta$	M1
	$=2\text{cosec}2\theta$ (✱)	A1 cso (4)
Alt.(*a*)	$\frac{\sin\theta}{\cos\theta}+\frac{\cos\theta}{\sin\theta}=\tan\theta+\frac{1}{\tan\theta}=\frac{\tan^2\theta+1}{\tan\theta}$ M1	
	$=\frac{\sec^2\theta}{\tan\theta}$ M1	
	$=\frac{1}{\cos\theta\sin\theta}=\frac{1}{\frac{1}{2}\sin 2\theta}$ M1	
	$=2\text{cosec}2\theta$ (✱) (cso) A1	
	If show two expressions are equal, need conclusion such as QED, tick, true.	
(*b*)	 Shape (May be translated but need to see 4"sections")	B1
	T.P.s at $y=\pm 2$, asymptotic at correct x-values (dotted lines not required)	B1 dep. (2)
(*c*)	$2\text{cosec}2\theta=3$	
	$\sin 2\theta=\frac{2}{3}$ Allow $\frac{2}{\sin 2\theta}=3$ [M1 for equation in $\sin 2\theta$]	M1, A1
	(2θ) = [41.810…°, 138.189…° ; 401.810…°, 498.189…°]	M1; M1
	1st M1 for α, $180-\alpha$; 2nd M1 adding 360° to at least one of values	
	θ = 20.9°, 69.1°, 200.9°, 249.1° (1 d.p.) awrt	
Note	1st A1 for any two correct, 2nd A1 for other two	A1,A1 (6)
	Extra solutions in range lose final A1 only	
	SC: Final 4 marks: **θ = 20.9°, after M0M0 is B1; record as M0M0A1A0**	
Alt.(*c*)	$\tan\theta+\frac{1}{\tan\theta}=3$ and form quadratic , $\tan^2\theta-3\tan\theta+1=0$ M1, A1	
	(M1 for attempt to multiply through by tanθ, A1 for correct equation above)	
	Solving quadratic [$\tan\theta=\frac{3\pm\sqrt{5}}{2}$ = 2.618… or = 0.3819…] M1	
	θ = 69.1°, 249.1° θ = 20.9°, 200.9° (1 d.p.) M1, A1, A1	**(12 marks)**
	(M1 is for one use of 180° + α°, A1A1 as for main scheme)	

Question Number	Scheme	Marks
8. (*a*)	$D = 10$, $t = 5$, $x = 10e^{-\frac{1}{8}\times 5}$ $= 5.353$ awrt	M1 A1 (2)
(*b*)	$D = 10+10e^{-\frac{5}{8}}$, $t = 1$, $x = 15.3526\ldots \times e^{-\frac{1}{8}}$ $x = 13.549$ (❋)	M1 A1 cso (2)
Alt.(*b*)	$x = 10e^{-\frac{1}{8}\times 6}+10e^{-\frac{1}{8}\times 1}$ M1 $x = 13.549$ (❋) A1 cso	
(*c*)	$15.3526...e^{-\frac{1}{8}T} = 3$ $e^{-\frac{1}{8}T} = \frac{3}{15.3526...} = 0.1954...$ $-\frac{1}{8}T = \ln 0.1954...$ $T = 13.06\ldots$ or 13.1 or 13	M1 M1 A1 (3) **(7 marks)**

Notes: (b) (main scheme) M1 is for $(10+10e^{-\frac{5}{8}})e^{-\frac{1}{8}}$, or {10 + their(a)}$e^{-(1/8)}$

N.B. The answer is given. There are many correct answers seen which deserve M0A0 or M1A0

(If adding two values, these should be 4.724 and 8.825)

(c) 1st M is for $(10+10e^{-\frac{5}{8}})\, e^{-\frac{T}{8}} = 3$

2nd M is for converting $e^{-\frac{T}{8}} = k$ (k > 0) to $-\frac{T}{8} = \ln k$. This is independent of 1st M.

Trial and improvement: M1 as scheme,
M1 correct process for their equation (two equal to 3 s.f.)
A1 as scheme