

1. (a) Find the binomial expansion of

$$\frac{1}{\sqrt[3]{(9-10x)}}, \quad |x| < \frac{9}{10}$$

in ascending powers of x up to and including the term in x^2 .
Give each coefficient as a simplified fraction.

(5)

- (b) Hence, or otherwise, find the expansion of

$$\frac{3+x}{\sqrt{(9-10x)}}, \quad |x| < \frac{9}{10}$$

in ascending powers of x , up to and including the term in x^2 .
Give each coefficient as a simplified fraction.

(3)

This image shows a full page of blank, lined paper. It features approximately 20 evenly spaced horizontal grey lines across its entire width, providing a template for writing or drawing. The background is a solid off-white color.

2.


**Figure 1**

Figure 1 shows a sketch of part of the curve with equation

$$y = (2 - x)e^{2x}, \quad x \in \mathbb{R}$$

The finite region R , shown shaded in Figure 1, is bounded by the curve, the x -axis and the y -axis.

The table below shows corresponding values of x and y for $y = (2 - x)e^{2x}$

x	0	0.5	1	1.5	2
y	2	4.077	7.389	10.043	0

- (a) Use the trapezium rule with all the values of y in the table, to obtain an approximation for the area of R , giving your answer to 2 decimal places. (3)
- (b) Explain how the trapezium rule can be used to give a more accurate approximation for the area of R . (1)
- (c) Use calculus, showing each step in your working, to obtain an exact value for the area of R . Give your answer in its simplest form. (5)





3.

$$x^2 + y^2 + 10x + 2y - 4xy = 10$$

- (a) Find $\frac{dy}{dx}$ in terms of x and y , fully simplifying your answer.

(5)

- (b) Find the values of y for which $\frac{dy}{dx} = 0$

(5)

This image shows a single sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

4. (a) Express $\frac{25}{x^2(2x+1)}$ in partial fractions.

(4)




Figure 2

Figure 2 shows a sketch of part of the curve C with equation $y = \frac{5}{x\sqrt{2x+1}}$, $x > 0$

The finite region R is bounded by the curve C , the x -axis, the line with equation $x = 1$ and the line with equation $x = 4$

This region is shown shaded in Figure 2

The region R is rotated through 360° about the x -axis.

- (b) Use calculus to find the exact volume of the solid of revolution generated, giving your answer in the form $a + b \ln c$, where a , b and c are constants.

(6)

[illegible]



- [You are given that $V = \frac{4}{3}\pi r^3$ and that $S = 4\pi r^2$]

(a) Find $\frac{dr}{dt}$ when the radius of the sphere is 4 cm, giving your answer to 3 significant figures.

(4)

- (2)

[illegible]

6. With respect to a fixed origin, the point A with position vector $\mathbf{i} + 2\mathbf{j} + 3\mathbf{k}$ lies on the line l_1 with equation

$$\mathbf{r} = \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix} + \lambda \begin{pmatrix} 0 \\ 2 \\ -1 \end{pmatrix}, \quad \text{where } \lambda \text{ is a scalar parameter,}$$

and the point B with position vector $4\mathbf{i} + p\mathbf{j} + 3\mathbf{k}$, where p is a constant, lies on the line l_2 with equation

$$\mathbf{r} = \begin{pmatrix} 7 \\ 0 \\ 7 \end{pmatrix} + \mu \begin{pmatrix} 3 \\ -5 \\ 4 \end{pmatrix}, \quad \text{where } \mu \text{ is a scalar parameter.}$$

- (a) Find the value of the constant p . (1)
- (b) Show that l_1 and l_2 intersect and find the position vector of their point of intersection, C . (4)
- (c) Find the size of the angle ACB , giving your answer in degrees to 3 significant figures. (3)
- (d) Find the area of the triangle ABC , giving your answer to 3 significant figures. (2)

This image shows a single sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.





8.




Figure 3

The curve shown in Figure 3 has parametric equations

$$x = t - 4 \sin t, \quad y = 1 - 2 \cos t, \quad -\frac{2\pi}{3} \leq t \leq \frac{2\pi}{3}$$

The point A, with coordinates $(k, 1)$, lies on the curve.

Given that $k > 0$

(a) find the exact value of k , (2)

(b) find the gradient of the curve at the point A. (4)

There is one point on the curve where the gradient is equal to $-\frac{1}{2}$

(c) Find the value of t at this point, showing each step in your working and giving your answer to 4 decimal places.

[Solutions based entirely on graphical or numerical methods are not acceptable.] (6)



Leave
blank

Question 8 continued

Q8

(Total 12 marks)

TOTAL FOR PAPER: 75 MARKS

END

