



1. Use integration by parts to find the exact value of  $\int_1^3 x^2 \ln x \, dx$ . (6)
- 

2. Fluid flows out of a cylindrical tank with constant cross section. At time  $t$  minutes,  $t \geq 0$ , the volume of fluid remaining in the tank is  $V \text{ m}^3$ . The rate at which the fluid flows, in  $\text{m}^3 \text{ min}^{-1}$ , is proportional to the square root of  $V$ .

(a) Show that the depth  $h$  metres of fluid in the tank satisfies the differential equation

$$\frac{dh}{dt} = -k\sqrt{h}, \quad \text{where } k \text{ is a positive constant.} \quad (3)$$

(b) Show that the general solution of the differential equation may be written as

$$h = (A - Bt)^2, \quad \text{where } A \text{ and } B \text{ are constants.} \quad (4)$$

Given that at time  $t = 0$  the depth of fluid in the tank is 1 m, and that 5 minutes later the depth of fluid has reduced to 0.5 m,

(c) find the time,  $T$  minutes, which it takes for the tank to empty. (3)

(d) Find the depth of water in the tank at time  $0.5T$  minutes. (2)

---

3. (a) Use the identity for  $\cos(A + B)$  to prove that  $\cos 2A = 2 \cos^2 A - 1$ . (2)

(b) Use the substitution  $x = 2\sqrt{2} \sin \theta$  to prove that

$$\int_2^{\sqrt{6}} \sqrt{8 - x^2} \, dx = \frac{1}{3}(\pi + 3\sqrt{3} - 6). \quad (7)$$

A curve is given by the parametric equations

$$x = \sec \theta, \quad y = \ln(1 + \cos 2\theta), \quad 0 \leq \theta < \frac{\pi}{2}.$$

(c) Find an equation of the tangent to the curve at the point where  $\theta = \frac{\pi}{3}$ . (5)

---

4.

**Figure 2**




Figure 2 shows a sketch of the curve  $C$  with equation  $y = \frac{4}{x-3}$ ,  $x \neq 3$ .

The points  $A$  and  $B$  on the curve have  $x$ -coordinates 3.25 and 5 respectively.

(a) Write down the  $y$ -coordinates of  $A$  and  $B$ . (1)

(b) Show that an equation of  $C$  is  $\frac{3y+4}{y}$ ,  $y \neq 0$ . (1)

The shaded region  $R$  is bounded by  $C$ , the  $y$ -axis and the lines through  $A$  and  $B$  parallel to the  $x$ -axis. The region  $R$  is rotated through  $360^\circ$  about the  $y$ -axis to form a solid shape  $S$ .

(c) Find the volume of  $S$ , giving your answer in the form  $\pi(a + b \ln c)$ , where  $a$ ,  $b$  and  $c$  are integers. (7)

The solid shape  $S$  is used to model a cooling tower. Given that 1 unit on each axis represents 3 metres,

(d) show that the volume of the tower is approximately  $15\,500 \text{ m}^3$ . (2)

5. Relative to a fixed origin  $O$ , the point  $A$  has position vector  $3\mathbf{i} + 2\mathbf{j} - \mathbf{k}$ , the point  $B$  has position vector  $5\mathbf{i} + \mathbf{j} + \mathbf{k}$ , and the point  $C$  has position vector  $7\mathbf{i} - \mathbf{j}$ .

(a) Find the cosine of angle  $ABC$ . (4)

(b) Find the exact value of the area of triangle  $ABC$ . (3)

The point  $D$  has position vector  $7\mathbf{i} + 3\mathbf{k}$ .

(c) Show that  $AC$  is perpendicular to  $CD$ . (2)

(d) Find the ratio  $AD:DB$ . (2)

---

6.

Figure 2




Figure 2 shows the cross-section of a road tunnel and its concrete surround. The curved section of the tunnel is modelled by the curve with equation  $y = 8 \sqrt{\left(\sin \frac{\pi x}{10}\right)}$ , in the interval  $0 \leq x \leq 10$ . The concrete surround is represented by the shaded area bounded by the curve, the  $x$ -axis and the lines  $x = -2$ ,  $x = 12$  and  $y = 10$ . The units on both axes are metres.

- (a) Using this model, copy and complete the table below, giving the values of  $y$  to 2 decimal places.

$x$	0	2	4	6	8	10
$y$	0	6.13				0

(2)

The area of the cross-section of the tunnel is given by  $\int_0^{10} y \, dx$ .

- (b) Estimate this area, using the trapezium rule with all the values from your table.

(4)

- (c) Deduce an estimate of the cross-sectional area of the concrete surround.

(1)

- (d) State, with a reason, whether your answer in part (c) over-estimates or under-estimates the true value.

(2)

7.

$$f(x) = \frac{25}{(3+2x)^2(1-x)}, \quad |x| < 1.$$

(a) Express  $f(x)$  as a sum of partial fractions.

(4)

(b) Hence find  $\int f(x) \, dx$ .

(5)

(c) Find the series expansion of  $f(x)$  in ascending powers of  $x$  up to and including the term in  $x^2$ . Give each coefficient as a simplified fraction.

(7)

---

END