

Paper Reference (complete below)							
<b>6</b>	<b>6</b>	<b>6</b>	<b>3</b>	<b>/</b>	<b>0</b>	<b>1</b>	

Centre No.					
Candidate No.					

Surname	Initial(s)
Signature	

Paper Reference(s)

**6663**

# Edexcel GCE

## Core Mathematics C3

### Advanced Subsidiary

### Set A: Practice Paper 3

Time: 1 hour 30 minutes

Examiner's use only

--	--	--

Team Leader's use only

--	--	--

**Materials required for examination**

Mathematical Formulae

**Items included with question papers**

Nil

Question Number	Leave Blank
1	
2	
3	
4	
5	
6	
7	
8	
<b>Total</b>	

**Instructions to Candidates**

In the boxes above, write your centre number, candidate number, your surname, initials and signature. You must write your answer for each question in the space following the question. If you need more space to complete your answer to any question, use additional answer sheets.

**Information for Candidates**

A booklet 'Mathematical Formulae and Statistical Tables' is provided.  
Full marks may be obtained for answers to ALL questions.  
This paper has nine questions.

**Advice to Candidates**

You must ensure that your answers to parts of questions are clearly labelled.  
You must show sufficient working to make your methods clear to the examiner.  
Answers without working may gain no credit.

*Turn over*

1.

---

2.

**Figure 2**




Figure 2 shows part of the curve  $C$  with equation  $y = f(x)$ , where

$$f(x) = 0.5e^x - x^2.$$

The curve  $C$  cuts the  $y$ -axis at  $A$  and there is a minimum at the point  $B$ .

(a) Find an equation of the tangent to  $C$  at  $A$ . (4)

The  $x$ -coordinate of  $B$  is approximately 2.15. A more exact estimate is to be made of this coordinate using iterations  $x_{n+1} = \ln g(x_n)$ .

(b) Show that a possible form for  $g(x)$  is  $g(x) = 4x$ . (3)

(c) Using  $x_{n+1} = \ln 4x_n$ , with  $x_0 = 2.15$ , calculate  $x_1$ ,  $x_2$  and  $x_3$ . Give the value of  $x_3$  to 4 decimal places. (2)

---

3. (a) Sketch the graph of  $y = |2x + a|$ ,  $a > 0$ , showing the coordinates of the points where the graph meets the coordinate axes. **(2)**
- (b) On the same axes, sketch the graph of  $y = \frac{1}{x}$ . **(1)**
- (c) Explain how your graphs show that there is only one solution of the equation

$$x|2x + a| - 1 = 0. \quad \mathbf{(1)}$$

- (d) Find, using algebra, the value of  $x$  for which  $x|2x + 1| - 1 = 0$ . **(3)**
- 

4.

**Figure 1**




Figure 1 shows a sketch of the curve with equation  $y = f(x)$ ,  $-1 \leq x \leq 3$ . The curve touches the  $x$ -axis at the origin  $O$ , crosses the  $x$ -axis at the point  $A(2, 0)$  and has a maximum at the point  $B(\frac{4}{3}, 1)$ .

In separate diagrams, show a sketch of the curve with equation

- (a)  $y = f(x + 1)$ , **(3)**
- (b)  $y = |f(x)|$ , **(3)**
- (c)  $y = f(|x|)$ , **(4)**

marking on each sketch the coordinates of points at which the curve

- (i) has a turning point,
- (ii) meets the  $x$ -axis.
-

5. (i) Given that  $\sin x = \frac{3}{5}$ , use an appropriate double angle formula to find the exact value of  $\sec 2x$ . (4)

(ii) Prove that

$$\cot 2x + \operatorname{cosec} 2x \equiv \cot x, \quad \left( x \neq \frac{n\pi}{2}, n \in \mathbb{Z} \right).$$

(4)

---

6. The function  $f$  is defined by  $f: x \rightarrow \frac{3x-1}{x-3}, x \in I, x \neq 3$ .

(a) Prove that  $f^{-1}(x) = f(x)$  for all  $x \in I, x \neq 3$ .

(3)

(b) Hence find, in terms of  $k$ ,  $ff(k)$ , where  $x \neq 3$ .

$I$   
(2)

**Figure 3**




Figure 3 shows a sketch of the one-one function  $g$ , defined over the domain  $-2 \leq x \leq 2$ .

(c) Find the value of  $fg(-2)$ .

(3)

(d) Sketch the graph of the inverse function  $g^{-1}$  and state its domain.

(3)

The function  $h$  is defined by  $h: x \mapsto 2g(x-1)$ .

(e) Sketch the graph of the function  $h$  and state its range.

(3)

7. (i) (a) Express  $(12 \cos \theta - 5 \sin \theta)$  in the form  $R \cos (\theta + \alpha)$ , where  $R > 0$  and  $0 < \alpha < 90^\circ$ . (4)

(b) Hence solve the equation

$$12 \cos \theta - 5 \sin \theta = 4,$$

for  $0 < \theta < 90^\circ$ , giving your answer to 1 decimal place. (3)

(ii) Solve

$$8 \cot \theta - 3 \tan \theta = 2,$$

for  $0 < \theta < 90^\circ$ , giving your answer to 1 decimal place. (5)

- 
8. The curve  $C$  has equation  $y = f(x)$ , where

$$f(x) = 3 \ln x + \frac{1}{x}, \quad x > 0.$$

The point  $P$  is a stationary point on  $C$ .

(a) Calculate the  $x$ -coordinate of  $P$ . (4)

(b) Show that the  $y$ -coordinate of  $P$  may be expressed in the form  $k - k \ln k$ , where  $k$  is a constant to be found. (2)

The point  $Q$  on  $C$  has  $x$ -coordinate 1.

(c) Find an equation for the normal to  $C$  at  $Q$ . (4)

The normal to  $C$  at  $Q$  meets  $C$  again at the point  $R$ .

(d) Show that the  $x$ -coordinate of  $R$

(i) satisfies the equation  $6 \ln x + x + \frac{2}{x} - 3 = 0$ ,

(ii) lies between 0.13 and 0.14. (4)

---

**END**