

4723 Core Mathematics 3

1 (i) Show correct process for composition of functions	M1 numerical or algebraic; the right way round
Obtain $(-3$ and hence) -23	A1 2
(ii) <u>Either</u> : State or imply $x^3 + 4 = 12$	B1
Attempt solution of equation involving x^3	M1 as far as $x = \dots$
Obtain 2	A1 3 and no other value
<u>Or</u> : Attempt expression for f^{-1}	M1 involving x or y ; involving cube root
Obtain $\sqrt[3]{x-4}$ or $\sqrt[3]{y-4}$	A1
Obtain 2	A1 (3) and no other value
<hr/>	
2 (i) Obtain correct first iterate 2.864	B1 or greater accuracy 2.864327...; condone 2 dp here and in working
Carry out correct iteration process	M1 to find at least 3 iterates in all
Obtain 2.877	A1 3 after at least 4 steps; answer required to exactly 3 dp
$[3 \rightarrow 2.864327 \rightarrow 2.878042 \rightarrow 2.876661 \rightarrow 2.876800]$	
(ii) State or imply $x = \sqrt[3]{31 - \frac{5}{2}x}$	B1
Attempt rearrangement of equation in x	M1 involving cubing and grouping non-zero terms on LHS
Obtain equation $2x^3 + 5x - 62 = 0$	A1 3 or equiv with integers
<hr/>	
3 (a) State correct equation involving $\cos \frac{1}{2}\alpha$	B1 such as $\cos \frac{1}{2}\alpha = \frac{1}{4}$ or $\frac{1}{\cos \frac{1}{2}\alpha} = 4$
Attempt to find value of α	M1 or ...
Obtain 151	A1 3 using correct order for the steps or greater accuracy; and no other values between 0 and 180
(b) State or imply $\cot \beta = \frac{1}{\tan \beta}$	B1
Rearrange to the form $\tan \beta = k$	M1 or equiv involving $\sin \beta$ only or $\cos \beta$ only; allow missing \pm
Obtain 69.3	A1
Obtain 111	A1 4 or greater accuracy; and no others between 0 and 180
<hr/>	
4 (i) Obtain derivative of form $kh^5(h^6 + 16)^n$	M1 any constant k ; any $n < \frac{1}{2}$; allow if -4 term retained
Obtain correct $3h^5(h^6 + 16)^{-\frac{1}{2}}$	A1 or (unsimplified) equiv; no -4 now
Substitute to obtain 10.7	A1 3 or greater accuracy or exact equiv
(ii) Attempt multn or divn using 8 and answer from (i) M1	
Attempt 8 divided by answer from (i)	M1
Obtain 0.75	A1 $\sqrt{3}$ or greater accuracy; allow 0.75 ± 0.01 ; following their answer from (i)

5 (a)	Obtain integral of form $k(3x + 7)^{10}$	M1	any constant k
	Obtain (unsimplified) $\frac{1}{10} \times \frac{1}{3} (3x + 7)^{10}$	A1	or equiv
	Obtain (simplified) $\frac{1}{30} (3x + 7)^{10} + c$	A1	3
(b)	State $\int \pi \left(\frac{1}{2\sqrt{x}}\right)^2 dx$	B1	or equiv involving x ; condone no dx
	Integrate to obtain $k \ln x$	M1	any constant k involving π or not; or equiv such as $k \ln 4x$ or $k \ln 2x$
	Obtain $\frac{1}{4}\pi \ln x$ or $\frac{1}{4} \ln x$ or $\frac{1}{4}\pi \ln 4x$ or $\frac{1}{4} \ln 4x$	A1	5
	Show use of the $\log a - \log b$ property	M1	not dependent on earlier marks
	Obtain $\frac{1}{4}\pi \ln 2$	A1	5 or similarly simplified equiv
<hr/>			
6 (i)	<u>Either:</u> Refer to translation and reflection State translation by 1 in negative x -direction	B1	in either order; allow clear equivs or equiv but now using correct terminology
	State reflection in x -axis	B1	3 using correct terminology
	<u>Or:</u> Refer to translation and reflection State reflection in y -axis State translation by 1 in positive x -direction	B1	in either order; allow clear equivs
		B1	(3) with order reflection then translation clearly intended
(ii)	Show sketch with attempt at reflection of 'negative' part in x -axis Show (more or less) correct sketch	M1	and curve for $0 < x < 1$ unchanged
		A1	2 with correct curvature
(iii)	Attempt correct process for finding at least one value	M1	as far as $x = \dots$; accept decimal equivs (degrees or radians) or expressions involving $\sin(\frac{1}{3}\pi)$
	Obtain $1 - \frac{1}{2}\sqrt{3}$	A1	or exact equiv
	Obtain $1 + \frac{1}{2}\sqrt{3}$	A1	3 or exact equiv; give A1A0 if extra incorrect solution(s) provided
<hr/>			
7 (i)	Attempt use of product rule for $x e^{2x}$	M1	obtaining $\dots + \dots$
	Obtain $e^{2x} + 2x e^{2x}$	A1	or equiv; maybe within QR attempt
	Attempt use of quotient rule	M1	with or without product rule
	Obtain unsimplified $\frac{(x+k)(e^{2x} + 2xe^{2x}) - xe^{2x}}{(x+k)^2}$	A1	
	Obtain $\frac{e^{2x}(2x^2 + 2kx + k)}{(x+k)^2}$	A1	5 AG; necessary detail required
(ii)	Attempt use of discriminant	M1	or equiv
	Obtain $4k^2 - 8k = 0$ or equiv and hence $k = 2$	A1	
	Attempt solution of $2x^2 + 2kx + k = 0$	M1	using their numerical value of k or solving in terms of k using correct formula
	Obtain $x = -1$	A1	
	Obtain $-e^{-2}$	A1	5 or exact equiv

8 (i)	State or imply $h = 1$ Attempt calculation involving attempts at y values Obtain $a(1 + 4 \times 2 + 2 \times 4 + 4 \times 8 + 2 \times 16 + 4 \times 32 + 64)$ A1 Obtain 91	B1 M1 A1 4	addition with each of coefficients 1, 2, 4 occurring at least once; involving at least 5 y values any constant a
(ii)	State $e^{x \ln 2}$ or $k = \ln 2$ Integrate e^{kx} to obtain $\frac{1}{k}e^{kx}$ Obtain $\frac{1}{\ln 2}(e^{6 \ln 2} - e^0)$ Simplify to obtain $\frac{63}{\ln 2}$	B1 M1 A1 A1 4	allow decimal equiv such as $e^{0.69x}$ any constant k or in terms of general k or exact equiv allow if simplification in part (iii)
(iii)	Equate answers to (i) and (ii) Obtain $\frac{63}{91}$ and hence $\frac{9}{13}$	M1 A1 2	provided $\ln 2$ involved other than in power of e AG ; necessary correct detail required

9 (i)	State at least one of $\cos \theta \cos 60 - \sin \theta \sin 60$ and $\cos \theta \cos 30 - \sin \theta \sin 30$ Attempt complete multiplication of identities of form $\pm \cos \cos \pm \sin \sin$ Use $\cos^2 \theta + \sin^2 \theta = 1$ and $2 \sin \theta \cos \theta = \sin 2\theta$ Obtain $\sqrt{3} - 2 \sin 2\theta$	B1 M1 M1 A1 4	with values $\frac{1}{2}\sqrt{3}$, $\frac{1}{2}$ involved AG ; necessary detail required
(ii)	Attempt use of 22.5 in right-hand side Obtain $\sqrt{3} - \sqrt{2}$	M1 A1 2	or exact equiv
(iii)	Obtain 10.7 Attempt correct process to find two angles Obtain 79.3	B1 M1 A1 3	or greater accuracy; allow ± 0.1 from values of 2θ between 0 and 180 or greater accuracy and no others between 0 and 90; allow ± 0.1
(iv)	Indicate or imply that critical values of $\sin 2\theta$ are -1 and 1 Obtain both of $k > \sqrt{3} + 2$, $k < \sqrt{3} - 2$ Obtain complete correct solution	M1 A1 A1 3	condoning decimal equivs, $\leq \geq$ signs now with exact values and unambiguously stated