

**ADVANCED GCE UNIT
MATHEMATICS**

Core Mathematics 3
MONDAY 11 JUNE 2007

4723/01

Afternoon

Time: 1 hour 30 minutes

Additional Materials: Answer Booklet (8 pages)
List of Formulae (MF1)

INSTRUCTIONS TO CANDIDATES

- Write your name, centre number and candidate number in the spaces provided on the answer booklet.
- Answer **all** the questions.
- Give non-exact numerical answers correct to 3 significant figures unless a different degree of accuracy is specified in the question or is clearly appropriate.
- You are permitted to use a graphical calculator in this paper.

INFORMATION FOR CANDIDATES

- The number of marks is given in brackets [] at the end of each question or part question.
- The total number of marks for this paper is 72.

ADVICE TO CANDIDATES

- Read each question carefully and make sure you know what you have to do before starting your answer.
- **You are reminded of the need for clear presentation in your answers.**

This document consists of **4** printed pages.

1 Differentiate each of the following with respect to x .

(i) $x^3(x+1)^5$ [2]

(ii) $\sqrt{3x^4+1}$ [3]

2 Solve the inequality $|4x-3| < |2x+1|$. [5]

3 The function f is defined for all non-negative values of x by

$$f(x) = 3 + \sqrt{x}.$$

(i) Evaluate $ff(169)$. [2]

(ii) Find an expression for $f^{-1}(x)$ in terms of x . [2]

(iii) On a single diagram sketch the graphs of $y = f(x)$ and $y = f^{-1}(x)$, indicating how the two graphs are related. [3]

4 The integral I is defined by

$$I = \int_0^{13} (2x+1)^{\frac{1}{3}} dx.$$

(i) Use integration to find the exact value of I . [4]

(ii) Use Simpson's rule with two strips to find an approximate value for I . Give your answer correct to 3 significant figures. [3]

5 A substance is decaying in such a way that its mass, m kg, at a time t years from now is given by the formula

$$m = 240e^{-0.04t}.$$

(i) Find the time taken for the substance to halve its mass. [3]

(ii) Find the value of t for which the mass is decreasing at a rate of 2.1 kg per year. [4]

6 (i) Given that $\int_0^a (6e^{2x} + x) dx = 42$, show that $a = \frac{1}{2} \ln(15 - \frac{1}{6}a^2)$. [5]


(ii) Use an iterative formula, based on the equation in part (i), to find the value of a correct to 3 decimal places. Use a starting value of 1 and show the result of each iteration. [4]

- 7 (i) Sketch the graph of $y = \sec x$ for $0 \leq x \leq 2\pi$. [2]
- (ii) Solve the equation $\sec x = 3$ for $0 \leq x \leq 2\pi$, giving the roots correct to 3 significant figures. [3]
- (iii) Solve the equation $\sec \theta = 5 \operatorname{cosec} \theta$ for $0 \leq \theta \leq 2\pi$, giving the roots correct to 3 significant figures. [4]

8 (i) Given that $y = \frac{4 \ln x - 3}{4 \ln x + 3}$, show that $\frac{dy}{dx} = \frac{24}{x(4 \ln x + 3)^2}$. [3]

- (ii) Find the exact value of the gradient of the curve $y = \frac{4 \ln x - 3}{4 \ln x + 3}$ at the point where it crosses the x -axis. [4]

(iii)



The diagram shows part of the curve with equation

$$y = \frac{2}{x^2(4 \ln x + 3)}.$$

The region shaded in the diagram is bounded by the curve and the lines $x = 1$, $x = e$ and $y = 0$. Find the exact volume of the solid produced when this shaded region is rotated completely about the x -axis. [4]

- 9 (i) Prove the identity

$$\tan(\theta + 60^\circ) \tan(\theta - 60^\circ) \equiv \frac{\tan^2 \theta - 3}{1 - 3 \tan^2 \theta}. \quad [4]$$

- (ii) Solve, for $0^\circ < \theta < 180^\circ$, the equation

$$\tan(\theta + 60^\circ) \tan(\theta - 60^\circ) = 4 \sec^2 \theta - 3,$$

giving your answers correct to the nearest 0.1° . [5]

- (iii) Show that, for all values of the constant k , the equation

$$\tan(\theta + 60^\circ) \tan(\theta - 60^\circ) = k^2$$

has two roots in the interval $0^\circ < \theta < 180^\circ$. [3]

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (OCR) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

OCR is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.