

**Mark Scheme 4751**  
**January 2007**


## Section A

1	$y = 2x + 4$	3	M1 for $m = 2$ stated [M0 if go on to use $m = -\frac{1}{2}$ ] or M1 for $y = 2x + k$ , $k \neq 7$ and M1indep for $y - 10 = m(x - 3)$ or $(3, 10)$ subst in $y = mx + c$ ; allow 3 for $y = 2x + k$ and $k = 4$	3
2	neg quadratic curve intercept $(0, 9)$ <u>through</u> $(3, 0)$ and $(-3, 0)$	1 1 1	condone $(0, 9)$ seen eg in table	3
3	$[a =] \frac{2c}{2-f}$ or $\frac{-2c}{f-2}$ as final answer	3	M1 for attempt to collect as and cs on different sides and M1 ft for $a(2-f)$ or dividing by $2-f$ ; allow M2 for $\frac{7c-5c}{2-f}$ etc	3
4	$f(2) = 3$ seen or used $2^3 + 2k + 5 = 3$ o.e. $k = -5$	M1 M1 B1	allow M1 for divn by $(x-2)$ with $x^2 + 2x + (k+4)$ or $x^2 + 2x - 1$ obtained alt: M1 for $(x-2)(x^2 + 2x - 1) + 3$ (may be seen in division) then M1dep (and B1) for $x^3 - 5x + 5$ alt divn of $x^3 + kx + 2$ by $x - 2$ with no rem.	3
5	375	3	allow $375x^4$ ; M1 for $5^2$ or 25 used or seen with $x^4$ and M1 for 15 or $\frac{6 \times 5}{2}$ oe eg $\frac{6!}{4!2!}$ or $1 \ 6 \ 15$ ... seen [ ${}^6C_4$ not sufft]	3
6	(i) 125  (ii) $\frac{9}{49}$ as final answer	2  2	M1 for $25^{\frac{1}{2}} = \sqrt{25}$ soi or for $\sqrt{25^3}$  M1 for $a^{-1} = \frac{1}{a}$ soi eg by $3/7$ or $3/49$	4
7	showing $a + b + c = 6$ o.e $bc = \frac{9^2 - 17}{16}$  =64/16 o.e. correctly obtained completion showing $abc = 6$ o.e.	1 M1  A1 A1	simple equiv fraction eg $192/32$ or $24/4$ correct expansion of numerator; may be unsimplified 4 term expansion; M0 if get no further than $(\sqrt{17})^2$ ; M0 if no evidence before $64/16$ o.e.  may be implicit in use of factors in completion	4

<b>8</b>	$b^2 - 4ac$ soi use of $b^2 - 4ac < 0$ $k^2 < 16$ [may be implied by $k < 4$ ] $-4 < k < 4$ or $k > -4$ and $k < 4$ isw	M1 M1 A1 A1	may be implied by $k^2 < 16$ deduct one mark in qn for $\leq$ instead of $<$ ; allow equalities earlier if final inequalities correct; condone $b$ instead of $k$ ; if M2 not earned, give SC2 for qn [or M1 SC1] for $k [=] 4$ and $-4$ as answer]	4
<b>9</b>	(i) $12a^5b^3$ as final answer  (ii) $\frac{(x+2)(x-2)}{(x-2)(x-3)}$ $\frac{x+2}{x-3}$ as final answer	2  M2  A1	1 for 2 'terms' correct in final answer  M1 for each of numerator or denom. correct or M1, M1 for correct factors seen separately	5
<b>10</b>	correct expansion of both brackets seen (may be unsimplified), or difference of squares used  $4m^2$ correctly obtained $[p =] [\pm] 2m$ cao	M2  A1 A1	M1 for one bracket expanded correctly; for M2, condone done together and lack of brackets round second expression if correct when we insert the pair of brackets	4

## Section B

<b>11</b>	<b>iA</b>	0.2 to 0.3 and 3.7 to 3.8	1+1	[tol. 1mm or 0.05 throughout qn]; if 0, allow M1 for drawing down lines at both values	2
	<b>iB</b>	$x + \frac{1}{x} = 4 - x$ their $y = 4 - x$ drawn	M1 M1	condone one error allow M2 for plotting positive branch of $y = 2x + 1/x$ [plots at (1,3) and (2,4.5) and above other graph] or for plot of $y = 2x^2 - 4x + 1$	4
		0.2 to 0.35 and 1.65 to 1.8	B2	1 each	2
	<b>ii</b>	$(0, \pm\sqrt{3})$	2	condone $y = \pm\sqrt{3}$ isw; 1 each or M1 for $1 + y^2 = 4$ or $y^2 = 3$ o.e.	4
	<b>iii</b>	centre (1, 0) radius 2 touches at (1, 2) [which is distance 2 from centre] all points on other branch > 2 from centre	1+1 1 1	allow seen in (ii) allow ft for both these marks for centre at (-1, 0), rad 2; allow 2 for good sketch or compass-drawn circle of rad 2 centre $(\pm 1, 0)$	4

12	i	<p>(3, 6)</p> <p>grad AB = <math>(8 - 4)/(7 - -1)</math> or 4/8 grad normal = -2 or ft</p> <p>perp bisector is <math>y - 6 = -2(x - 3)</math> or ft their grad. of normal (not AB) and/or midpoint correct step towards completion</p>	2	<p>1 each coord</p> <p>M1 indep obtained M1 for use of <math>m_1 m_2 = -1</math>; condone stated/used as -2 with no working only if 4/8 seen</p> <p>M1 or M1 for showing grad given line = -2 A1 and M1 for showing (3, 6) fits given line</p>	6
	ii	<p>Bisector crosses y axis at C (0, 12) seen or used AB crosses y axis at D (0, 4.5) seen or used</p> <p><math>\frac{1}{2} \times (12 - \text{their } 4.5) \times 3</math> (may be two triangles M1 each)</p> <p>45/4 o.e. without surds, isw</p>  <p><u>alt</u> allow integration used: <math>\int_0^3 (-2x + 12) dx [= 27]</math></p> <p>obtaining AB is <math>y - 8 = \text{their } \frac{1}{2}(x - 7)</math> oe [<math>y = \frac{1}{2}x + 4.5</math>] <math>\int_0^3 (\frac{1}{2}x + 4.5) dx</math> = 63/4 o.e. cao their area under CB – their area under AB = 45/4 o.e. cao</p>	<p>M1 may be implicit in their area calcn</p> <p>B2 M1 for 4 + their grad AB or for eqn AB is <math>y - 8 = \text{their } \frac{1}{2}(x - 7)</math> oe with coords of A or their M used M2 or M1 for <math>[MC]^2 = 3^2 + 6^2</math> or 45 or <math>[MD]^2 = 3^2 + 1.5^2</math> or 11.25 oe and M1 for <math>\frac{1}{2} \times \text{their } MC \times MD</math>; all ft their M A1</p> <p><u>MR</u>: AMC used not DMC: lose B2 for D but then allow ft M1 for <math>MC^2</math> or <math>MA^2 [= 4^2 + 2^2]</math> and M1 for <math>\frac{1}{2} \times MA \times MC</math> and A1 for 15</p> <p><u>MR</u>: intn used as D(0, 4) can score a max of M1, B0, M2 (eg M1 for their <math>DM = \sqrt{13}</math>), A0</p> <p>M1 condone poor notation</p> <p>M1 allow if seen, with correct line and limits seen/used as above</p> <p>M1 ft from their AB</p> <p>A1</p> <p>M1</p> <p>A1 allow only if at least some valid integration/area calculations for these trapezia seen if combined integration, so 63/4 not found separately, mark equivalently for Ms and allow A2 for final answer</p>	<p>6</p>	6
13	i	<p><math>x - 2</math> is factor soi attempt at divn by <math>x - 2</math> as far as <math>x^3 - 2x^2</math> seen in working <math>x^2 + 2x - 1</math> obtained attempt at quad formula or comp square <math>-1 \pm \sqrt{2}</math> as final answer</p>	<p>M1 eg may be implied by divn M1 or other factor (<math>x^2 \dots -1</math>) or (<math>x^2 + 2x \dots</math>)</p> <p>A1 or B3 www M1 ft their quadratic</p> <p>A2 A1 for <math>\frac{-2 \pm \sqrt{8}}{2}</math> seen; or B3 www</p>	<p>6</p>	6

ii	$f(x-3) = (x-3)^3 - 5(x-3) + 2$ $(x-3)(x^2 - 6x + 9)$ or other constructive attempt at expanding $(x-3)^3$ eg 1 3 3 1 soi $x^3 - 9x^2 + 27x - 27 - 5x + 15 [+2]$	B1 M1 A1 B1	or $(x-5)(x-2+\sqrt{2})(x-2-\sqrt{2})$ soi or ft from their (i) for attempt at multiplying out 2 brackets or valid attempt at multiplying all 3 alt: A2 for correct full unsimplified expansion or A1 for correct 2 bracket expansion eg $(x-5)(x^2 - 4x + 2)$	4
iii	5 $2 \pm \sqrt{2}$ or ft	B1 B1	condone factors here, not roots if B0 in this part, allow SC1 for their roots in (i) - 3	2