

## 4751 (C1) Introduction to Advanced Mathematics

1	$[a =]2c^2 - b$ www o.e.	3	M1 for each of 3 complete correct steps, ft from previous error if equivalent difficulty
2	$5x - 3 < 2x + 10$ $3x < 13$ $x < \frac{13}{3}$ o.e.	M1 M1 M1	condone '=' used for first two Ms M0 for just $5x - 3 < 2(x + 5)$ or $-13 < -3x$ or ft or ft; isw further simplification of $13/3$ ; M0 for just $x < 4.3$
3 (i)	(4, 0)	1	allow $y = 0, x = 4$ bod B1 for $x = 4$ but do not isw: 0 for (0, 4) seen 0 for (4, 0) and (0, 10) both given (choice) unless (4, 0) clearly identified as the $x$ -axis intercept
3 (ii)	$5x + 2(5 - x) = 20$ o.e.  (10/3, 5/3) www isw	M1 A2	for subst or for multn to make coeffs same and appropriate addn/subtn; condone one error  or A1 for $x = 10/3$ and A1 for $y = 5/3$ o.e. isw; condone 3.33 or better and 1.67 or better  A1 for (3.3, 1.7)
4 (i)	translation by $\begin{pmatrix} -4 \\ 0 \end{pmatrix}$ or 4 [units] to left	B1 B1	0 for shift/move or 4 units in negative $x$ direction o.e.
4 (ii)	sketch of parabola right way up and with minimum on negative $y$ -axis  min at (0, -4) and graph through -2 and 2 on $x$ -axis	B1 B1	mark intent for both marks  must be labelled or shown nearby
5 (i)	$\frac{1}{12}$ or $\pm \frac{1}{12}$	2	M1 for $\frac{1}{144^{\frac{1}{2}}}$ o.e. or for $\sqrt{144} = 12$ soi
5 (ii)	denominator = 18  numerator = $5 - \sqrt{7} + 4(5 + \sqrt{7})$ = $25 + 3\sqrt{7}$ as final answer	B1 M1 A1	B0 if 36 after addition  for M1, allow in separate fractions  allow B3 for $\frac{25 + 3\sqrt{7}}{18}$ as final answer www

6 (i)	cubic correct way up and with two turning pts  touching $x$ -axis at $-1$ , and through it at $2.5$ and no other intersections  $y$ - axis intersection at $-5$	<b>B1</b>  <b>B1</b>  <b>B1</b>	intns must be shown labelled or worked out nearby
6 (ii)	$2x^3 - x^2 - 8x - 5$	<b>2</b>	<b>B1</b> for 3 terms correct or <b>M1</b> for correct expansion of product of two of the given factors
7	attempt at $f(-3)$ $-27 + 18 - 15 + k = 6$  $k = 30$	<b>M1</b> <b>A1</b>  <b>A1</b>	or <b>M1</b> for long division by $(x + 3)$ as far as obtaining $x^2 - x$ and <b>A1</b> for obtaining remainder as $k - 24$ (but see below)  equating coefficients method: <b>M2</b> for $(x + 3)(x^2 - x + 8) [+6]$ o.e. (from inspection or division) eg <b>M2</b> for obtaining $x^2 - x + 8$ as quotient in division
8	$x^3 + 15x + \frac{75}{x} + \frac{125}{x^3}$ www isw or $x^3 + 15x + 75x^{-1} + 125x^{-3}$ www isw	<b>4</b>	<b>B1</b> for <b>both</b> of $x^3$ and $\frac{125}{x^3}$ or $125x^{-3}$ isw  and  <b>M1</b> for 1 3 3 1 soi; <b>A1</b> for <b>each</b> of $15x$ and $\frac{75}{x}$ or $75x^{-1}$ isw  <b>or</b> <b>SC2</b> for completely correct unsimplified answer

<p><b>9</b></p> <p><math>x^2 - 5x + 7 = 3x - 10</math></p> <p><math>x^2 - 8x + 17 [= 0]</math> o.e or <math>y^2 - 4y + 13 [= 0]</math> o.e</p> <p>use of <math>b^2 - 4ac</math> with numbers subst (condone one error in substitution) (may be in quadratic formula)</p> <p><math>b^2 - 4ac = 64 - 68</math> or <math>-4</math> cao [or <math>16 - 52</math> or <math>-36</math> if <math>y</math> used]</p> <p>[<math>&lt; 0</math>] so no [real] roots [so line and curve do not intersect]</p>		<p><b>M1</b></p> <p><b>M1</b></p> <p><b>M1</b></p> <p><b>A1</b></p> <p><b>A1</b></p>	<p>or attempt to subst <math>(y + 10)/3</math> for <math>x</math></p> <p>condone one error; allow <b>M1</b> for <math>x^2 - 8x = -17</math> [oe for <math>y</math>] only if they go on to completing square method</p> <p>or <math>(x - 4)^2 = 16 - 17</math> or <math>(x - 4)^2 + 1 = 0</math> (condone one error)</p> <p>or <math>(x - 4)^2 = -1</math> or <math>x = 4 \pm \sqrt{-1}</math> [or <math>(y - 2)^2 = -9</math> or <math>y = 2 \pm \sqrt{-9}</math>]</p> <p>or conclusion from comp. square; needs to be explicit correct conclusion and correct ft; allow '<math>&lt; 0</math> so no intersection' o.e.; allow '<math>-4</math> so no roots' etc</p> <p>allow A2 for full argument from sum of two squares = 0; A1 for weaker correct conclusion</p> <p>some may use the condition <math>b^2 &lt; 4ac</math> for no real roots; allow equivalent marks, with first A1 for <math>64 &lt; 68</math> o.e.</p>
<p><b>10 (i)</b></p> <p>grad CD = <math>\frac{5-3}{3-(-1)} = \frac{2}{4}</math> o.e. ] isw</p> <p>grad AB = <math>\frac{3-(-1)}{6-(-2)}</math> or <math>\frac{4}{8}</math> isw</p> <p>same gradient so parallel www</p>		<p><b>M1</b></p> <p><b>M1</b></p> <p><b>A1</b></p>	<p>NB needs to be obtained independently of grad AB</p> <p>must be explicit conclusion mentioning 'same gradient' or 'parallel'</p> <p>if M0, allow <b>B1</b> for 'parallel lines have same gradient' o.e.</p>
<p><b>10 (ii)</b></p> <p>[<math>BC^2 =</math>] <math>3^2 + 2^2</math> [<math>BC^2 =</math>] 13 showing <math>AD^2 = 1^2 + 4^2 [= 17]</math> [<math>\neq BC^2</math>] isw</p>		<p><b>M1</b></p> <p><b>A1</b></p> <p><b>A1</b></p>	<p>accept <math>(6 - 3)^2 + (3 - 5)^2</math> o.e. or [BC =] <math>\sqrt{13}</math> or [AD =] <math>\sqrt{17}</math></p> <p>or equivalent marks for finding AD or <math>AD^2</math> first</p> <p>alt method: showing <math>AC \neq BD</math> – mark equivalently</p>

10 (iii)	<p>[BD eqn is] <math>y = 3</math></p> <p>eqn of AC is <math>y - 5 = 6/5 \times (x - 3)</math> o.e [ <math>y = 1.2x + 1.4</math> o.e.]</p> <p>M is <math>(4/3, 3)</math> o.e. isw</p>	<p><b>M1</b></p> <p><b>M2</b></p> <p><b>A1</b></p>	<p>eg allow for 'at M, <math>y = 3</math>' or for 3 subst in eqn of AC</p> <p>or <b>M1</b> for grad AC = <math>6/5</math> o.e. (accept unsimplified) and <b>M1</b> for using their grad of AC with coords of A(-2, -1) or C (3, 5) in eqn of line or <b>M1</b> for 'stepping' method to reach M</p> <p>allow : at M, <math>x = 16/12</math> o.e. [eg <math>=4/3</math>] isw A0 for 1.3 without a fraction answer seen</p>
10 (iv)	<p>midpt of BD = <math>(5/2, 3)</math> or equivalent simplified form cao</p> <p>midpt AC = <math>(1/2, 2)</math> or equivalent simplified form cao or 'M is <math>2/3</math> of way from A to C'</p> <p>conclusion 'neither diagonal bisects the other'</p>	<p><b>M1</b></p> <p><b>M1</b></p> <p><b>A1</b></p>	<p>or showing <math>BM \neq MD</math> oe [<math>BM = 14/3, MD = 7/3</math>]</p> <p>or showing <math>AM \neq MC</math> or <math>AM^2 \neq MC^2</math></p> <p>in these methods A1 is dependent on coords of M having been obtained in part (iii) or in this part; the coordinates of M need not be correct; it is also dependent on midpts of both AC and BD attempted, at least one correct</p> <p>alt method: show that mid point of BD does not lie on AC (M1) and vice-versa (M1), A1 for both and conclusion</p>

<b>11 (i)</b>	centre $C' = (3, -2)$ radius 5	<b>1</b> <b>1</b>	0 for $\pm 5$ or $-5$
<b>11 (ii)</b>	showing $(6 - 3)^2 + (-6 + 2)^2 = 25$  showing that $\overrightarrow{AC'} = \overrightarrow{C'B} = \begin{pmatrix} -3 \\ 4 \end{pmatrix}$ o.e.	<b>B1</b>  <b>B2</b>	interim step needed  <b>or B1</b> each for two of: showing midpoint of $AB = (3, -2)$ ; showing $B(0, 2)$ is on circle; showing $AB = 10$  <b>or B2</b> for showing midpoint of $AB = (3, -2)$ and saying this is centre of circle  <b>or B1</b> for finding eqn of AB as $y = -4/3x + 2$ o.e. and <b>B1</b> for finding one of its intersections with the circle is $(0, 2)$  <b>or B1</b> for showing $C'B = 5$ and <b>B1</b> for showing $AB = 10$ or that $AC'$ and $BC'$ have the same gradient  <b>or B1</b> for showing that $AC'$ and $BC'$ have the same gradient and <b>B1</b> for showing that $B(0, 2)$ is on the circle
<b>11 (iii)</b>	grad $AC'$ or $AB = -4/3$ o.e.  grad $tgt = -1/\text{their } AC'$ grad  $y - (-6) = \text{their } m(x - 6)$ o.e.  $y = 0.75x - 10.5$ o.e. isw	<b>M1</b>  <b>M1</b>  <b>M1</b>  <b>A1</b>	or ft from their $C'$ , must be evaluated  may be seen in eqn for $tgt$ ; allow <b>M2</b> for $\text{grad } tgt = 3/4$ oe soi as first step  or <b>M1</b> for $y = \text{their } m \times x + c$ then subst $(6, -6)$  eg <b>A1</b> for $4y = 3x - 42$  allow <b>B4</b> for correct equation www isw
<b>11 (iv)</b>	centre $C$ is at $(12, -14)$ cao  circle is $(x - 12)^2 + (y + 14)^2 = 100$	<b>B2</b>  <b>B1</b>	<b>B1</b> for each coord  ft their $C$ if at least one coord correct

<b>12 (i)</b>	10	<b>1</b>	
<b>12 (ii)</b>	[x =] 5 or ft their (i) $\div 2$  ht = 5[m] cao	<b>1</b>  <b>1</b>	not necessarily ft from (i) eg they may start again with calculus to get $x = 5$
<b>12 (iii)</b>	$d = 7/2$ o.e.  [y =] $1/5 \times 3.5 \times (10 - 3.5)$ o.e. or ft  = $91/20$ o.e. cao isw	<b>M1</b>  <b>M1</b>  <b>A1</b>	or ft their (ii) $- 1.5$ or their (i) $\div 2 - 1.5$ o.e. or $7 - 1/5 \times 3.5^2$ or ft or showing $y - 4 = 11/20$ o.e. cao
<b>12 (iv)</b>	$4.5 = 1/5 \times x(10 - x)$ o.e.  $22.5 = x(10 - x)$ o.e.  $2x^2 - 20x + 45 [= 0]$ o.e. eg $x^2 - 10x + 22.5 [= 0]$ or $(x - 5)^2 = 2.5$  [x =] $\frac{20 \pm \sqrt{40}}{4}$ or $5 \pm \frac{1}{2}\sqrt{10}$ o.e.  width = $\sqrt{10}$ o.e. eg $2\sqrt{2.5}$ cao	<b>M1</b>  <b>M1</b>  <b>A1</b>  <b>M1</b>  <b>A1</b>	eg $4.5 = x(2 - 0.2x)$ etc cao; accept versions with fractional coefficients of $x^2$ , isw or $x - 5 = [\pm]\sqrt{2.5}$ o.e.; ft their quadratic eqn provided at least M1 gained already; condone one error in formula or substitution; need not be simplified or be real accept simple equivalents only