

Mathematics

Advanced GCE

Unit **4723**: Core Mathematics 3

Mark Scheme for January 2011

OCR (Oxford Cambridge and RSA) is a leading UK awarding body, providing a wide range of qualifications to meet the needs of pupils of all ages and abilities. OCR qualifications include AS/A Levels, Diplomas, GCSEs, OCR Nationals, Functional Skills, Key Skills, Entry Level qualifications, NVQs and vocational qualifications in areas such as IT, business, languages, teaching/training, administration and secretarial skills.

It is also responsible for developing new specifications to meet national requirements and the needs of students and teachers. OCR is a not-for-profit organisation; any surplus made is invested back into the establishment to help towards the development of qualifications and support which keep pace with the changing needs of today's society.

This mark scheme is published as an aid to teachers and students, to indicate the requirements of the examination. It shows the basis on which marks were awarded by Examiners. It does not indicate the details of the discussions which took place at an Examiners' meeting before marking commenced.

All Examiners are instructed that alternative correct answers and unexpected approaches in candidates' scripts must be given marks that fairly reflect the relevant knowledge and skills demonstrated.

Mark schemes should be read in conjunction with the published question papers and the Report on the Examination.

OCR will not enter into any discussion or correspondence in connection with this mark scheme.

© OCR 2011

Any enquiries about publications should be addressed to:

OCR Publications
PO Box 5050
Annesley
NOTTINGHAM
NG15 0DL

Telephone: 0870 770 6622
Facsimile: 01223 552610
E-mail: publications@ocr.org.uk

1	<u>Either:</u> Obtain $\frac{1}{3}a$	B1	condone $ x = \frac{1}{3}a$
	Attempt solution of linear eqn	M1	with signs of $3x$ and $5a$ different; allow M1 only if a given particular value and no recovery occurs; allow M1 only if a in terms of x attempted; allow M1 only if double inequality attempted but with no recovery to state actual values of x
	Obtain $-3a$	A1	3 as final answer
	<u>Or:</u> Obtain $9x^2 + 24ax + 16a^2 = 25a^2$	B1	
	Attempt solution of 3-term quad eqn	M1	as far as substitution into correct quadratic formula or correct factorisation of their quadratic; allow M1 only if a given particular value
	Obtain $-3a$ and $\frac{1}{3}a$	A1	(3) or equivs; as final answers; and no others
			3

2	Draw graph showing reflection in a horizontal axis	M1	
	Draw graph showing translation	M1	parallel to x -axis, in either direction; independent of first M1; not earned if curve still passes through O but ignore other coordinates given at this stage
	Draw (more or less) correct graph which must at least reach the negative x -axis, if not cross it, at left end of curve	A1	but ignoring no or wrong stretch in y -dir'n; condone graph existing only for $x < 0$; consider shape of curve and ignore coordinates given
State $(-5, 24)$ and $(-3, 0)$ wherever located	B1	4	or clearly implied by sketch; allow for coordinates whatever sketch looks like; allow if in solution with no sketch
			4

3	<u>Either:</u> State or imply $8\pi r$ as derivative	B1	or equiv
	Attempt to connect 12 and their derivative	M1	numerical or algebraic; using multiplication or division
	Obtain $8\pi \times 150 \times 12$ and hence 45000 or 14400π or 14000π	A1	3 or equiv; or greater accuracy (45239); condone absence of units or use of wrong units
	<u>Or:</u> Use $r = 12t$ to show $S = 576\pi t^2$	B1	
	Attempt $\frac{dS}{dt}$ and substitute for t	M1	
	Obtain $1152\pi \times \frac{150}{12}$ and hence 45000 or 14400π or 14000π	A1	(3) or equiv; or greater accuracy (45239); condone absence of units or use of wrong units
			3

4 (i)	Obtain $R = 25$ Attempt to find value of α	B1 M1	allow $\sqrt{625}$ or value rounding to 25 implied by correct answer or its complement; allow sin/cos muddles; allow use of radians for this mark; condone $\sin \alpha = 7$, $\cos \alpha = 24$ in the working
	Obtain 16.3°	A1	3 or greater accuracy 16.260...; must be degrees now; allow 16° here

(ii)	Show correct process for finding one answer Obtain $(28.69 - 16.26$ and hence) 12.4°	M1 A1	even if leading to answer outside 0 to 360 or greater accuracy 12.425... or anything rounding to 12.4
	Show correct process for finding second answer Obtain $(151.31 - 16.26$ and hence) 135° or 135.1°	M1 A1	even if further incorrect answers produced 4 or greater accuracy 135.054...; and no other between 0 and 360
	[SC: No working shown and 2 correct angles stated		- B1 only in part (ii)]
			7

5	Integrate to obtain form $k(3x - 2)^{\frac{1}{2}}$	M1	any non-zero constant k ; or equiv involving substitution
	Obtain correct $4(3x - 2)^{\frac{1}{2}}$	A1	or (unsimplified) equiv such as $\frac{6(3x - 2)^{\frac{1}{2}}}{3 \times \frac{1}{2}}$
	Apply limits and attempt solution for a	M1	assuming integral of form $k(3x - 2)^n$; taking solution as far as removal of root; with subtraction the right way round; if sub'n used, limits must be appropriate
	Obtain $a = 9$	A1	(this answer written down with no working scores 0/4 so far but all subsequent marks are available)
	State or imply formula $\int \frac{36\pi}{3x - 2} dx$	B1	or (unsimplified) equiv; condone absence of dx; allow B1 retroactively if π absent here but inserted later
	Integrate to obtain form $k \ln(3x - 2)$	*M1	any constant k including π or not; condone absence of brackets
	Obtain $12\pi \ln(3x - 2)$ or $12 \ln(3x - 2)$	A1√	following their integral of form $\int \frac{k}{3x - 2} dx$
	Apply limits the correct way round	M1	dep *M; use of limit 1 is implied by absence of second term; allow use of limit a
	Obtain $12\pi \ln 25$ (or $24\pi \ln 5$)	A1	9 or exact equiv but not with $\ln 1$ remaining; condone answers such as $\pi 12 \ln 25$ and $12 \ln 25\pi$
			9

6 (i)	Attempt use of quotient rule	M1	or equiv; allow numerator wrong way round but needs minus sign in numerator; for M1 condone 'minor' errors such as sign slips, absence of square in denominator, and absence of some brackets
	Obtain $\frac{3(x^3 - 4x^2 + 2) - (3x + 4)(3x^2 - 8x)}{(x^3 - 4x^2 + 2)^2}$	A1	or equiv; allow A1 if brackets absent from $3x + 4$ term or from $3x^2 - 8x$ term but not from both
	Equate numerator to 0 and attempt simplification	M1	at least as far as removing brackets, condoning sign or coeff slips; or equiv
	Obtain $-6x^3 + 32x + 6 = 0$ or equiv and hence $x = \sqrt[3]{\frac{16}{3}x + 1}$	A1	4 AG; necessary detail needed (i.e. at least one intermediate step) and following first derivative with correct numerator

(ii)	Obtain correct first iterate having used initial value 2.4	B1	showing at least 3 dp (2.398 or 2.399 or greater accuracy 2.39861...)
	Apply iterative process	M1	to obtain at least 3 iterates in all; implied by plausible, converging sequence of values; having started with any initial non-negative value
	Obtain at least 3 correct iterates from their starting point	A1	allowing recovery after error
	Obtain 2.398	A1	value required to exactly 3 dp
	Obtain -1.552	A1	5 value required to exactly 3 dp; allow if apparently obtained by substitution of 2.4; answers only with no iterates shown gets 0/5
	[2.4 → 2.3986103 → 2.3981808 → 2.3980480]		

7 (i)	State $\ln(x^2 + 8) = 8$ Attempt solution involving e^8 Obtain $\sqrt{e^8 - 8}$	B1 M1 A1	or equiv such as $x^2 + 8 = e^8$ by valid (exact) method at least as far as $x^2 = \dots$ 3 or exact equiv; and no other answer

(ii)	State f only State e^x or e^y Indicate domain is all real numbers	B1 B1 B1	3 or equiv; allow if g, or f and g, chosen however expressed

(iii)	Attempt use of chain rule Obtain $\frac{2 \ln x}{x}$ Obtain $6e^{-3}$	M1 A1 A1	3 whether applied to gf or fg; or equiv such as use of product rule on $(\ln x)(\ln x) + 8$ or equiv 3 or exact equiv but not including ln

(iv)	Attempt evaluation using y attempts Obn $k(\ln 24 + 4 \ln 12 + 2 \ln 8 + 4 \ln 12 + \ln 24)$ Use $k = \frac{2}{3}$ and obtain 20.3	M1 A1 A1	3 with coeffs 1, 4 and 2 occurring at least once each; whether fg or gf any constant k 3 or greater accuracy (20.26...) but must round to 20.3
[Note that use of Simpson's rule between 0 and 4 with two strips, coeffs 1, 4, 1, followed by doubling of result is equiv;			
SC: Use of Simpson's rule between 0 and 4 with four strips followed by doubling of result - allow 3/3 - answer is 20.2 (20.2327...)]			

- 8 (a) (i)** Draw at least two correctly shaped branches, one for $y > 0$, one for $y < 0$ M1 otherwise located anywhere including $x < 0$
 Draw four correct branches M1 now (more or less) correctly located;
 Draw (more or less) correct graph A1 **3** with some indication of horiz scale (perhaps only 4π indicated); with asymptotic behaviour shown (but not too fussy about branch drifting slightly away from asymptotic value nor about branch touching asymptote) but branches must not obviously cross asymptotic value; with -1 and 1 shown (or implied by presence of sine curve or by presence of only one of them on a reasonably accurate sketch); no need for vertical (dotted) lines drawn to indicate asymptotic values

- **(ii)** State expression of form $k\pi + \alpha$ or $k\pi - \alpha$ or $\alpha = k\pi + \beta$ or $\alpha = k\pi - \beta$ M1 any non-zero numerical value of k ; M0 if degrees used
 State $3\pi - \alpha$ A1 **2** or unsimplified equiv

- **(b) (i)** State $\frac{2 \tan \theta}{1 - \tan^2 \theta}$ B1 **1** or equiv such as $\frac{t + t}{1 - t \times t}$ or $\frac{2 \tan A}{1 - \tan^2 A}$

- **(ii)** State or imply $\tan \phi = \frac{1}{4}$ B1 or equiv such as $\frac{1}{\tan \phi} = 4$
 Attempt to evaluate $\tan 2\phi$ or $\cot 2\phi$ M1 perhaps within attempt at complete expression but using correct identity
 Obtain $\tan 2\phi = \frac{8}{15}$ or $\cot 2\phi = \frac{15}{8}$ A1 or (unsimplified) equiv; may be implied
 Attempt to evaluate value of $\tan 4\phi$ M1 perhaps within attempt at complete expression; condone only minor slip(s) in use of relevant identity
 Obtain $\frac{240}{161}$ A1 or (unsimplified) exact equiv; may be implied
 Obtain final answer $\frac{225}{322}$ A1 **6** or exact equiv

[SC – (use of calculator and little or no working)]

State or imply $\tan \phi = \frac{1}{4}$ B1; Obtain $\tan 2\phi = \frac{8}{15}$ B1; Obtain $\frac{225}{322}$ B1 (max 3/6)

State or imply $\tan \phi = \frac{1}{4}$ B1; Obtain $\frac{225}{322}$ B2 (max 3/6)

9 (i) (a)	Differentiate to obtain $k_1e^{2x} + k_2e^{-2x}$	M1	any constants k_1 and k_2 but derivative must be different from $f(x)$; condone presence of $+c$
	Obtain $2e^{2x} + 6e^{-2x}$	A1	or unsimplified equiv; no $+c$ now
	Refer to $e^{2x} > 0$ and $e^{-2x} > 0$ or to more general comment about exponential functions	A1	3 or equiv (which might be sketch of $y = f(x)$ with comment that gradient is positive or might be sketch of $y = f'(x)$ with comment that $y > 0$; AG

(b)	Differentiate to obtain $k_3e^{2x} + k_4e^{-2x}$	M1	any constants k_3 and k_4 but second derivative must be different from their first derivative; condone presence of $+c$
	Obtain $4e^{2x} - 12e^{-2x}$	A1	or unsimplified equiv; no $+c$ now
	Attempt solution of $f''(x) > 0$ or of $f(x) > 0$ or of corresponding eqn	M1	at least as far as term involving e^{4x} or e^{-4x}
	Obtain $x > \frac{1}{4}\ln 3$	A1	
	Confirm both give same result	B1	5 AG; necessary detail needed; either by solving the other or by observing that same inequality involved (just noting that $f''(x) = 4f(x)$ is sufficient)

(ii)	Differentiate to obtain $2e^{2x} - 2ke^{-2x}$	B1	or unsimplified equiv
	Attempt to find x -coordinate of stationary pt	M1	equating to 0 and reaching $e^{4x} = \dots$ or equiv
	Obtain $e^{4x} = k$ and hence $\frac{1}{4}\ln k$ or equiv	A1	or equiv such as $e^{2x} = \sqrt{k}$
	Substitute and attempt simplification	M1	using valid processes but allow if only limited progress [note that question can be successfully concluded (without actually finding x) by substitution of $e^{2x} = \sqrt{k}$ and $e^{-2x} = \frac{1}{\sqrt{k}}$]
	Obtain $g(x) \geq 2\sqrt{k}$ or $y \geq 2\sqrt{k}$	A1	5 or similarly simplified equiv with \geq not $>$

OCR (Oxford Cambridge and RSA Examinations)
1 Hills Road
Cambridge
CB1 2EU

OCR Customer Contact Centre

14 – 19 Qualifications (General)

Telephone: 01223 553998

Facsimile: 01223 552627

Email: general.qualifications@ocr.org.uk

www.ocr.org.uk

For staff training purposes and as part of our quality assurance programme your call may be recorded or monitored

Oxford Cambridge and RSA Examinations
is a Company Limited by Guarantee
Registered in England
Registered Office; 1 Hills Road, Cambridge, CB1 2EU
Registered Company Number: 3484466
OCR is an exempt Charity

OCR (Oxford Cambridge and RSA Examinations)
Head office
Telephone: 01223 552552
Facsimile: 01223 552553

