

# Mark Scheme (Results) January 2011

GCE

GCE Mechanics M1 (6677) Paper 1

Edexcel is one of the leading examining and awarding bodies in the UK and throughout the world. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers.

Through a network of UK and overseas offices, Edexcel's centres receive the support they need to help them deliver their education and training programmes to learners.

For further information, please call our GCE line on 0844 576 0025, our GCSE team on 0844 576 0027, or visit our website at [www.edexcel.com](http://www.edexcel.com).

If you have any subject specific questions about the content of this Mark Scheme that require the help of a subject specialist, you may find our [Ask The Expert](#) email service helpful.

Ask The Expert can be accessed online at the following link:

<http://www.edexcel.com/Aboutus/contact-us/>

January 2011

Publications Code UA026577

All the material in this publication is copyright  
© Edexcel Ltd 2011

## General Instructions for Marking

1. The total number of marks for the paper is 75.
2. The Edexcel Mathematics mark schemes use the following types of marks:
  - **M** marks: method marks are awarded for 'knowing a method and attempting to apply it', unless otherwise indicated.
  - **A** marks: Accuracy marks can only be awarded if the relevant method (M) marks have been earned.
  - **B** marks are unconditional accuracy marks (independent of M marks)
  - Marks should not be subdivided.
3. Abbreviations



These are some of the traditional marking abbreviations that will appear in the mark schemes.

- bod - benefit of doubt
- ft - follow through
- the symbol  $\checkmark$  will be used for correct ft
- cao - correct answer only
- cso - correct solution only. There must be no errors in this part of the question to obtain this mark
- isw - ignore subsequent working
- awrt - answers which round to
- SC: special case
- oe - or equivalent (and appropriate)
- dep - dependent
- indep - independent
- dp decimal places
- sf significant figures
- \* The answer is printed on the paper
- $\square$  The second mark is dependent on gaining the first mark



January 2011  
Mechanics M1 6677  
Mark Scheme

Question Number	Scheme	Marks
1. (a)	Conservation of momentum: $4m - 6 = m + 9$ $m = 5$	M1 A1 A1 (3)
(b)	Impulse = change in momentum $= 3 \times 3 - (3 \times -2) = 15$	M1 A1 (2) [5]



Question Number	Scheme	Marks
2.		
(a)	$-6.45 = u - 9.8 \times 0.75$ $0.9 = u$ **	M1 A1 A1 (3)
(b)	$0 = 0.81 - 2 \times 9.8 \times s$ $s = 0.041$ or $0.0413$	M1 A1 (2)
(c)	$h = -0.9 \times 0.75 + 4.9 \times 0.75^2$ $h = 2.1$ or $2.08$	M1 A1 A1 (3) [8]


Question Number	Scheme	Marks
3.	<p>(a)</p>  <p>Taking moments about B: <math>5 \times R_C = 20g \times 3</math>  <math>R_C = 12g</math> or <math>60g/5</math> or 118 or 120</p> <p>Resolving vertically: <math>R_C + R_B = 20g</math>  <math>R_B = 8g</math> or 78.4 or 78</p>	<p>M1A1 A1</p> <p>M1 A1</p> <p>(5)</p>
	<p>(b)</p>  <p>Resolving vertically: <math>50g = R + R</math></p> <p>Taking moments about B:</p> $5 \times 25g = 3 \times 20g + (6 - x) \times 30g$ $30x = 115$ $x = 3.8 \text{ or better or } 23/6 \text{ oe}$	<p>B1</p> <p>M1 A1 A1</p> <p>A1</p> <p>(5) [10]</p>

Question Number	Scheme	Marks
4.		
(a)	$\text{speed} = \sqrt{2^2 + (-5)^2}$ $= \sqrt{29} = 5.4 \text{ or better}$	M1 A1 (2)
(b)	$((7\mathbf{i} + 10\mathbf{j}) - (2\mathbf{i} - 5\mathbf{j}))/5$ $= (5\mathbf{i} + 15\mathbf{j})/5 = \mathbf{i} + 3\mathbf{j}$ $\mathbf{F} = m\mathbf{a} = 2(\mathbf{i} + 3\mathbf{j}) = 2\mathbf{i} + 6\mathbf{j}$	M1 A1 A1 DM1 A1ft (5)
(c)	$\mathbf{v} = \mathbf{u} + \mathbf{a}t = (2\mathbf{i} - 5\mathbf{j}) + (\mathbf{i} + 3\mathbf{j})t$ $(-5 + 3t)\mathbf{j}$ <p>Parallel to <math>\mathbf{i} \Rightarrow -5 + 3t = 0</math></p> $t = 5/3$	M1 A1 M1 A1 (4) [11]

Question Number	Scheme	Marks
5.		
(a)		
(i)	 <p>1<sup>st</sup> section correct</p> <p>2<sup>nd</sup> &amp; 3<sup>rd</sup> sections correct</p> <p>Numbers and v marked correctly on the axes.</p>	<p>B1</p> <p>B1</p> <p>DB1</p>
(ii)	 <p>1<sup>st</sup> section correct</p> <p>2<sup>nd</sup> section correct</p> <p>3<sup>rd</sup> section correct and no “extras” on the sketch</p>	<p>B1</p> <p>B1</p> <p>B1</p> <p>(6)</p>
(b)	$\frac{70 + 40}{2} \times v = 880$ $v = 880 \times \frac{2}{110} = 16$	<p>M1 A1</p> <p>DM1 A1</p> <p>(4)</p> <p>[10]</p>



Question Number	Scheme	Marks
6. (a)	 <p>Resolving perpendicular to the plane:  <math>S = 120\cos\alpha + 30\sin\alpha</math>  <math>= 114 \text{ *}</math></p>	M1 A1 A1 A1 (4)
(b)	 <p>Resolving perpendicular to the plane:  <math>R = 120\cos\alpha</math>  <math>= 96</math>  <math>F_{\max} = \frac{1}{2}R</math></p> <p>Resolving parallel to the plane:  In equilibrium: <math>P_{\max} = F_{\max} + 120\sin\alpha</math>  <math>= 48 + 72 = 120</math></p>	M1 A1 A1 M1  M1 A(2,1,0) A1 (8)
(c)	$30 + F = 120\sin\alpha \text{ OR } 30 - F = 120\sin\alpha$  So $F = 42\text{N}$ acting up the plane.	M1 A1  A1 (3) [15]

Question Number	Scheme	Marks
7.	<p>(a)</p>  <p> <math>\tan \theta = \frac{5}{12}</math>  <math>\sin \theta = \frac{5}{13}</math>  <math>\cos \theta = \frac{12}{13}</math> </p> <p>For A: <math>7g - T = 7a</math>  For B: parallel to plane <math>T - F - 3g \sin \theta = 3a</math>  perpendicular to plane <math>R = 3g \cos \theta</math>  <math>F = \mu R = 3g \cos \theta = 2g \cos \theta</math></p> <p>Eliminating <math>T</math>, <math>7g - F - 3g \sin \theta = 10a</math>  Equation in <math>g</math> and <math>a</math>: <math>7g - 2g \times \frac{12}{13} - 3g \frac{5}{13} = 7g - \frac{39}{13}g = 4g = 10a</math>  <math>a = \frac{2g}{5}</math> oe or 3.9 or 3.92</p>	<p>M1 A1  M1 A1  M1 A1  M1  DM1  DM1  A1  (10)</p>
(b)	<p>After 1 m,</p> $v^2 = u^2 + 2as, \quad v^2 = 0 + 2 \times \frac{2g}{5} \times 1$ $v = 2.8$	<p>M1  A1  (2)</p>
(c)	<p> <math>-(F + 3g \sin \theta) = 3a</math>  <math>\frac{2}{3} \times 3g \times \frac{12}{13} + 3g \times \frac{5}{13} = 3g = -3a, \quad a = -g</math>  <math>v = u + at, \quad 0 = 2.8 - 9.8t,</math>  <math>t = \frac{2}{9.8}</math> oe, 0.29. 0.286 </p>	<p>M1  A1  DM1  A1  (4)  [16]</p>



Further copies of this publication are available from  
Edexcel Publications, Adamsway, Mansfield, Notts, NG18 4FN

Telephone 01623 467467  
Fax 01623 450481

Email [publications@linneydirect.com](mailto:publications@linneydirect.com)

Order Code UA026577 January 2011

For more information on Edexcel qualifications, please visit [www.edexcel.com/quals](http://www.edexcel.com/quals)

Edexcel Limited. Registered in England and Wales no.4496750  
Registered Office: One90 High Holborn, London, WC1V 7BH