

Mathematics (MEI)

Advanced Subsidiary GCE

Unit **4751**: Introduction to Advanced Mathematics

Mark Scheme for January 2013

OCR (Oxford Cambridge and RSA) is a leading UK awarding body, providing a wide range of qualifications to meet the needs of candidates of all ages and abilities. OCR qualifications include AS/A Levels, Diplomas, GCSEs, Cambridge Nationals, Cambridge Technicals, Functional Skills, Key Skills, Entry Level qualifications, NVQs and vocational qualifications in areas such as IT, business, languages, teaching/training, administration and secretarial skills.

It is also responsible for developing new specifications to meet national requirements and the needs of students and teachers. OCR is a not-for-profit organisation; any surplus made is invested back into the establishment to help towards the development of qualifications and support, which keep pace with the changing needs of today's society.

This mark scheme is published as an aid to teachers and students, to indicate the requirements of the examination. It shows the basis on which marks were awarded by examiners. It does not indicate the details of the discussions which took place at an examiners' meeting before marking commenced.

All examiners are instructed that alternative correct answers and unexpected approaches in candidates' scripts must be given marks that fairly reflect the relevant knowledge and skills demonstrated.

Mark schemes should be read in conjunction with the published question papers and the report on the examination.

OCR will not enter into any discussion or correspondence in connection with this mark scheme.

© OCR 2013

Annotations

Annotation	Meaning
✓ and ✗	
BOD	Benefit of doubt
FT	Follow through
ISW	Ignore subsequent working
M0, M1	Method mark awarded 0, 1
A0, A1	Accuracy mark awarded 0, 1
B0, B1	Independent mark awarded 0, 1
SC	Special case
^	Omission sign
MR	Misread
Highlighting	
Other abbreviations in mark scheme	Meaning
E1	Mark for explaining
U1	Mark for correct units
G1	Mark for a correct feature on a graph
M1 dep*	Method mark dependent on a previous mark, indicated by *
cao	Correct answer only
oe	Or equivalent
rot	Rounded or truncated
soi	Seen or implied
www	Without wrong working

Subject-specific Marking Instructions

- a Annotations should be used whenever appropriate during your marking.

The A, M and B annotations must be used on your standardisation scripts for responses that are not awarded either 0 or full marks. It is vital that you annotate standardisation scripts fully to show how the marks have been awarded.

For subsequent marking you must make it clear how you have arrived at the mark you have awarded.

- b An element of professional judgement is required in the marking of any written paper. Remember that the mark scheme is designed to assist in marking incorrect solutions. Correct *solutions* leading to correct answers are awarded full marks but work must not be judged on the answer alone, and answers that are given in the question, especially, must be validly obtained; key steps in the working must always be looked at and anything unfamiliar must be investigated thoroughly.

Correct but unfamiliar or unexpected methods are often signalled by a correct result following an *apparently* incorrect method. Such work must be carefully assessed. When a candidate adopts a method which does not correspond to the mark scheme, award marks according to the spirit of the basic scheme; if you are in any doubt whatsoever (especially if several marks or candidates are involved) you should contact your Team Leader.

- c The following types of marks are available.

M

A suitable method has been selected and *applied* in a manner which shows that the method is essentially understood. Method marks are not usually lost for numerical errors, algebraic slips or errors in units. However, it is not usually sufficient for a candidate just to indicate an intention of using some method or just to quote a formula; the formula or idea must be applied to the specific problem in hand, eg by substituting the relevant quantities into the formula. In some cases the nature of the errors allowed for the award of an M mark may be specified.

A

Accuracy mark, awarded for a correct answer or intermediate step correctly obtained. Accuracy marks cannot be given unless the associated Method mark is earned (or implied). Therefore M0 A1 cannot ever be awarded.

B

Mark for a correct result or statement independent of Method marks.

E

A given result is to be established or a result has to be explained. This usually requires more working or explanation than the establishment of an unknown result.

Unless otherwise indicated, marks once gained cannot subsequently be lost, eg wrong working following a correct form of answer is ignored. Sometimes this is reinforced in the mark scheme by the abbreviation isw. However, this would not apply to a case where a candidate passes through the correct answer as part of a wrong argument.

- d When a part of a question has two or more 'method' steps, the M marks are in principle independent unless the scheme specifically says otherwise; and similarly where there are several B marks allocated. (The notation 'dep *' is used to indicate that a particular mark is dependent on an earlier, asterisked, mark in the scheme.) Of course, in practice it may happen that when a candidate has once gone wrong in a part of a question, the work from there on is worthless so that no more marks can sensibly be given. On the other hand, when two or more steps are successfully run together by the candidate, the earlier marks are implied and full credit must be given.
- e The abbreviation ft implies that the A or B mark indicated is allowed for work correctly following on from previously incorrect results. Otherwise, A and B marks are given for correct work only — differences in notation are of course permitted. A (accuracy) marks are not given for answers obtained from incorrect working. When A or B marks are awarded for work at an intermediate stage of a solution, there may be various alternatives that are equally acceptable. In such cases, exactly what is acceptable will be detailed in the mark scheme rationale. If this is not the case please consult your Team Leader.

Sometimes the answer to one part of a question is used in a later part of the same question. In this case, A marks will often be 'follow through'. In such cases you must ensure that you refer back to the answer of the previous part question even if this is not shown within the image zone. You may find it easier to mark follow through questions candidate-by-candidate rather than question-by-question.

- f Wrong or missing units in an answer should not lead to the loss of a mark unless the scheme specifically indicates otherwise. Candidates are expected to give numerical answers to an appropriate degree of accuracy, with 3 significant figures often being the norm. Small variations in the degree of accuracy to which an answer is given (e.g. 2 or 4 significant figures where 3 is expected) should not normally be penalised, while answers which are grossly over- or under-specified should normally result in the loss of a mark. The situation regarding any particular cases where the accuracy of the answer may be a marking issue should be detailed in the mark scheme rationale. If in doubt, contact your Team Leader.

g Rules for replaced work

If a candidate attempts a question more than once, and indicates which attempt he/she wishes to be marked, then examiners should do as the candidate requests.

If there are two or more attempts at a question which have not been crossed out, examiners should mark what appears to be the last (complete) attempt and ignore the others.

NB Follow these maths-specific instructions rather than those in the assessor handbook.

h For a *genuine* misreading (of numbers or symbols) which is such that the object and the difficulty of the question remain unaltered, mark according to the scheme but following through from the candidate's data. A penalty is then applied; 1 mark is generally appropriate, though this may differ for some units. This is achieved by withholding one A mark in the question.

Note that a miscopy of the candidate's own working is not a misread but an accuracy error.

Question		Answer	Marks	Guidance
1	(i)	$\frac{9}{25}$ or 0.36 isw	2 [2]	M1 for numerator or denominator correct or for squaring correctly or for inverting correctly M1 for eg $\frac{1}{\left(\frac{25}{9}\right)}$ or $\left(\frac{25}{9}\right)^{-1}$ or $\frac{25}{9}$ or for $\left(\frac{3}{5}\right)^2$ or $\frac{3}{5}$ M0 for just $\frac{1}{\left(\frac{5}{3}\right)^2}$
1	(ii)	27	2 [2]	eg M1 for 3^3 M0 for $81^3 = 531441$ (true but not helpful)
2		$4x^4y^{-3}$ or $\frac{4x^4}{y^3}$ as final answer	3 [3]	B1 each 'term'; or M1 for numerator = $64x^{15}y^3$ and M1 for denominator = $16x^{11}y^6$ B0 if obtained fortuitously mark B scheme or M scheme to advantage of candidate, but not a mixture of both schemes

Question		Answer	Marks	Guidance
3		<p>obtaining a correct relationship in any 3 of C, d, r and A</p> <p>or obtaining a correct relationship in k and no more than 2 other variables</p> <p>convincing argument leading to $k = 4$</p>	<p>M2</p> <p>A1</p> <p>[3]</p>	<p>may substitute into given relationship;</p> <p>or M1 for at least two of $A = \pi r^2$, $C = \pi d$, $C = 2\pi r$, $d = 2r$ or $r = \frac{d}{2}$ seen or used</p> <p>must be from general argument, not just substituting values for r or d; may start from given relationship and derive $k = 4$</p> <p>eg M2 for $Cd = 4\pi r^2$ or $\pi d^2 = k\pi r^2$ seen/obtained</p> <p>condone eg Area = πr^2; allow $A = \pi \left(\frac{d}{2}\right)^2$ to imply $A = \pi r^2$ and $r = \frac{d}{2}$ and so earn M1, if M2 not earned</p> <p>eg M1 only for eg $A = \pi r^2$ and $C = \pi d$ and so $k = 4$ with no further evidence</p>
4		<p>$(5x + 2)(x - 6)$</p> <p>boundary values -0.4 oe and 6 soi</p> <p>$-0.4 \leq x \leq 6$ oe</p>	<p>M1</p> <p>A1</p> <p>A2</p> <p>[4]</p>	<p>for factors giving at least two out of three terms correct when expanded and collected</p> <p>A0 for just $\frac{28 \pm \sqrt{1024}}{10}$</p> <p>may be separate inequalities; mark final answer</p> <p>A1 for one end correct eg $x \leq 6$ or for $-0.4 < x < 6$ oe</p> <p>or B1 for $a \leq x \leq b$ ft their boundary values</p> <p>or use of formula or completing the square with at most one error (comp square must reach $[5](x - a)^2 \leq b$ oe or $(5x - c)^2 \leq d$ oe stage) if correct: $5(x - 2.8)^2 \leq 51.2$ or $(x - 2.8)^2 \leq 10.24$ or $(5x - 14)^2 \leq 256$</p> <p>condone unsimplified but correct $\frac{28 - \sqrt{1024}}{10} \leq x \leq \frac{28 + \sqrt{1024}}{10}$ etc</p> <p>allow A1 for $-0.4 \leq 0 \leq 6$</p> <p>condone errors in the inequality signs during working towards final answer</p>

Question	Answer	Marks	Guidance
5	$4 + 2k + c = 0$ or $2^2 + 2k + c = 0$ $9 - 3k + c = 35$ correct method to eliminate one variable from their eqns $k = -6, c = 8$ or $[x^2 + kx + c =] (x - 2)(x - a)$ $-5 \times (-3 - a) = 35$ oe $a = 4$ $k = -6, c = 8$	B1 B1 M1 A1 or M1 M1 A1 A1 [4]	may be rearranged may be rearranged; the $(-3)^2$ must be evaluated / used as 9 eg subtraction or substitution for c ; condone one error from fully correct method, allowing recovery from slips or or $(x - 2)(x + b)$ condone -3^2 seen if used as 9 M0 for addition of eqns unless also multiplied appropriately if no errors and no method seen, allow correct answers to imply M1 provided B1B1 has been earned

Question		Answer	Marks	Guidance
6		<p>identifying term as $20(2x)^3\left(\frac{5}{x}\right)^3$ oe</p> <p>20 000</p>	<p>M3</p> <p>M1 for $[k](2x)^3\left(\frac{5}{x}\right)^3$ soi (eg in list or table), condoning lack of brackets</p> <p>and M1 for $k = 20$ or eg $\frac{6 \times 5 \times 4}{3 \times 2 \times 1}$ or for 1 6 15 20 15 6 1 seen (eg Pascal's triangle seen, even if no attempt at expansion)</p> <p>and M1 for selecting the appropriate term (eg may be implied by use of only $k = 20$, but this M1 is not dependent on the correct k used)</p> <p>A1</p> <p>or B4 for 20 000 obtained from multiplying out $\left(2x + \frac{5}{x}\right)^6$</p> <p>allow SC3 for 20000 as part of an expansion</p> <p>[4]</p>	<p>xs may be omitted; eg M3 for $20 \times 8 \times 125$</p> <p>first M1 not earned if elements added not multiplied; otherwise, if in list or table bod intent to multiply</p> <p>M0 for binomial coefficient if it still has factorial notation</p> <p>may be gained even if elements added</p>
7	(i)	$9\sqrt{3}$ www oe as final answer	<p>2</p> <p>[2]</p>	<p>M1 for $\sqrt{48} = 4\sqrt{3}$ or $\sqrt{75} = 5\sqrt{3}$ soi</p>
7	(ii)	$\frac{39 + 7\sqrt{5}}{44}$ www as final answer	<p>3</p> <p>[3]</p>	<p>M1 for attempt to multiply numerator and denominator by $7 - \sqrt{5}$</p> <p>B1 for each of numerator and denominator correct (must be simplified)</p> <p>condone $\frac{39}{44} + \frac{7\sqrt{5}}{44}$ for 3 marks</p> <p>eg M0B1 if denominator correctly rationalised to 44 but numerator not multiplied</p>

Question		Answer	Marks	Guidance
8		$5c + 9t = 2ac + at$ $5c - 2ac = at - 9t$ oe $c(5 - 2a) = at - 9t$ oe $[c =] \frac{at - 9t}{5 - 2a}$ or $\frac{t(a - 9)}{5 - 2a}$ oe as final answer	M1 M1 M1 M1 [4]	for correct expansion of brackets for correct collection of terms, ft eg after M0 for $5c + 9t = 2ac + t$ allow this M1 for $5c - 2ac = -8t$ oe for correctly factorising, ft; must be $c \times$ a two-term factor for correct division, ft their two-term factor for each M, ft previous errors if their eqn is of similar difficulty; may be earned before t terms collected treat as MR if t is the subject, with a penalty of 1 mark from those gained, marking similarly
9	(i)	sketch of cubic the right way up, with two tps their graph touching the x -axis at -2 and crossing it at 3 and no other places intersection of y -axis at -12	B1 B1 B1 [3]	No section to be ruled; no curving back; condone some curving out at ends but not approaching another turning point; condone some doubling (eg erased curves may continue to show); ignore position of turning points for this mark mark intent if 'daylight' between curve and axis at $x = -2$ if no graph but -12 marked on y -axis, or in table, allow this 3 rd mark
9	(ii)	-5 and 0	B2 [2]	B1 each; allow B2 for $-5, -5, 0$; or B1 for both correct with one extra value or for $(-5, 0)$ and $(0, 0)$ or SC1 for both of 1 and 6 if their graph wrong, allow -5 and 0 from starting again with eqn, or ft their graph with two intns with x -axis

Question		Answer	Marks	Guidance
10	(i)	<p>midpt of AB = $\left(\frac{1}{2}, \frac{5}{2}\right)$ oe www</p> <p>grad AB = $\frac{4-1}{3-(-2)}$ oe</p> <p>using gradient of AB to obtain grad perp bisector</p> <p>$y - 2.5 = \frac{-5}{3}(x - 0.5)$ oe</p>	<p>B2</p> <p>M1</p> <p>M1</p> <p>M1</p>	<p>allow unsimplified B1 for one coordinate correct</p> <p>must be obtained independently of given line; accept 3 and 5 correctly shown eg in a sketch, followed by 3/5</p> <p>M1 for rise/run = 3/5 etc</p> <p>M0 for just 3/5 with no evidence</p> <p>for use of $m_1 m_2 = -1$ soi or ft their gradient AB</p> <p>M0 for just $\frac{-5}{3}$ without AB grad found</p> <p>eg M1 for $y = \frac{-5}{3}x + c$ and subst of midpt; ft their gradient of perp bisector and midpt; M0 for just rearranging given equation</p> <p>if working shown, should come from $\left(\frac{3+(-2)}{2}, \frac{4+1}{2}\right)$ oe</p> <p>NB B0 for x coord. = $\frac{5}{2}$, (obtained from subtraction instead of addition)</p> <p>for those who find eqn of AB first, M0 for just $\frac{y-4}{1-4} = \frac{x-3}{-2-3}$ oe, but M1 for $y-4 = \frac{1-4}{-2-3}(x-3)$ oe</p> <p>ignore their going on to find the eqn of AB after finding grad AB</p> <p>this second M1 available for starting with given line = $\frac{-5}{3}$ and obtaining grad. of AB from it</p> <p>no ft for gradient of AB used</p>

Question		Answer	Marks	Guidance
		completion to given answer $3y + 5x = 10$, showing at least one interim step	M1 [6]	<p>condone a slight slip if they recover quickly and general steps are correct (eg sometimes a slip in working with the c in $y = \frac{-5}{3}x + c$)</p> <p>- condone $3y = -5x + c$ followed by substitution and consistent working)</p> <p>M0 if clearly 'fudging'</p> <p>NB answer given; mark process not answer; annotate if full marks not earned eg with a tick for each mark earned</p> <p>scores such as B2M0M0M1M1 are possible</p> <p>after B2, allow full marks for complete method of showing given line has gradient perp to AB (grad AB must be found independently at some stage) and passes through midpt of AB</p>
10	(ii)	$3y + 5(4y - 21) = 10$ $(-1, 5)$ or $y = 5, x = -1$ isw	M1 A2 [3]	<p>or other valid strategy for eliminating one variable attempted eg $\frac{-5}{3}x + \frac{10}{3} = \frac{x}{4} + \frac{21}{4}$;</p> <p>condone one error</p> <p>or eg $20y = 5x + 105$ and subtraction of two eqns attempted</p> <p>no ft from wrong perp bisector eqn, since given</p> <p>allow M1 for candidates who reach $y = 115/23$ and then make a worse attempt, thinking they have gone wrong</p> <p>NB M0A0 in this part for finding E using info from (iii) that implies E is midpt of CD</p>

Question		Answer	Marks	Guidance	
10	(iii)	$(x - a)^2 + (y - b)^2 = r^2$ seen or used	M1	or for $(x + 1)^2 + (y - 5)^2 = k$, or ft their E, where $k > 0$	<p>this M not earned for use of CE or DE or $\frac{1}{2}$ CD</p> <p>NB some cand's finding $AB^2 = 34$ then obtaining 17 erroneously so M0</p> <p>SC also earned if circle comes from C or D and E, but may recover and earn the second M1 later by using A or B</p>
		$1^2 + 4^2$ oe (may be unsimplified), from clear use of A or B	M1	for calculating AE or BE or their squares, or for subst coords of A or B into circle eqn to find r or r^2 , ft their E;	
		$(x + 1)^2 + (y - 5)^2 = 17$	A1	for eqn of circle centre E, through A and B; allow A1 for $r^2 = 17$ found after $(x + 1)^2 + (y - 5)^2 = r^2$ stated and second M1 clearly earned	
		showing midpt of CD = $(-1, 5)$	M1	if $(x + 1)^2 + (y - 5)^2 = 17$ appears without clear evidence of using A or B, allow the first M1 then M0 SC1	
		showing CE or DE = $\sqrt{17}$ oe or showing one of C and D on circle	M1	alt M1 for showing $CD^2 = 68$ oe allow to be earned earlier as an invalid attempt to find r	

Question	Answer	Marks	Guidance
		[5]	<p>showing that both C and D are on circle and commenting that E is on CD is enough for last M1M1; similarly showing $CD^2 = 68$ and both C and D are on circle oe earns last M1M1</p> <p>other methods exist, eg: may find eqn of circle with centre E and through C or D and then show that A and B and other of C/D are on this circle – the marks are then earned in a different order; award M1 for first fact shown and then final M1 for completing the argument;</p> <p>if part-marks earned, annotate with a tick for each mark earned beside where earned</p>
11	<p>(i)</p> $\left(x - \frac{5}{2}\right)^2 - \frac{1}{4} \text{ oe}$ $\left(\frac{5}{2}, -\frac{1}{4}\right) \text{ oe or ft}$	<p>B3</p> <p>B1</p> <p>[4]</p>	<p>B1 for $a = 5/2$ oe and M1 for $6 - \text{their } a^2$ soi;</p> <p>accept $x = 2.5, y = -0.25$ oe</p> <p>condone $\left(x - \frac{5}{2}\right)^2 - \frac{1}{4} \text{ oe} = 0$ condone omission of index –can earn all marks</p> <p>bod M1 for $6 - 4.25$ or $6 - 25/2$ etc, if bearing some relation to an attempt at $6 - \text{their } 2.5^2$; M0 for just 1.75 etc without further evidence</p> <p>condone starting again and finding using calculus</p>

Question		Answer	Marks	Guidance	
11	(ii)	(2, 0) and (3, 0)	B2	B1 each or B1 for both correct plus an extra or M1 for $(x - 2)(x - 3)$ or correct use of formula or for <i>their a</i> $\pm \sqrt{\textit{their b}}$ ft from (i)	condone not expressed as coordinates, for both x and y values; accept eg in table or marked on graph
		(0, 6) graph of quadratic the correct way up and crossing both axes	B1 B1	ignore label of their tp; condone stopping at y-axis	condone 'U' shape or slight curving back in/out; condone some doubling / feathering – deleted work sometimes still shows up in scoris; must not be ruled; condone fairly straight with clear attempt at curve at minimum; be reasonably generous on attempt at symmetry
			[4]		
11	(iii)	$x^2 - 5x + 6 = 2 - x$	M1	for attempt to equate or subtract eqns or attempt at rearrangement and elimination of x	accept calculus approach: $y' = 2x - 5$
		$x^2 - 4x + 4 [= 0]$	M1	for rearrangement to zero ft and collection of terms; condone one error; if using completing the square, need to get as far as $(x - k)^2 = c$, with at most one error [(x - 2) ² = 0 if correct]	use of $y' = -1$ M1

Question		Answer	Marks	Guidance
		$x = 2, [y = 0]$ ‘double root at $x = 2$ so tangent’ oe; www;	A1 A1 [4]	condone omission of $y = 0$ since already found in (ii) if they have eliminated x , $y = 0$ is not sufft for A1 – need to get $x = 2$ A0 for $x = 2$ and another root eg ‘only one point of contact, so tangent’; or showing $b^2 - 4ac = 0$, and concluding ‘so tangent’; www $x = 2$ A1 tgt is $y [- 0] = -(x - 2)$ and obtaining given line A1
12	(i)	$f(1) = 1 - 1 + 1 + 9 - 10 [= 0]$ attempt at division by $(x - 1)$ as far as $x^4 - x^3$ in working correctly obtaining $x^3 + x + 10$	B1 M1 A1 [3]	condone $1^4 - 1^3 + 1^2 + 9 - 10$ eg for inspection, M1 for two terms right and two wrong if M0 and this division / factorising is done in part (ii) or (iii), allow SC1 if correct cubic obtained there; attach the relevant part to (i) with a formal chain link if not already seen in the image zone for (i)

Question		Answer	Marks	Guidance	
12	(iii)	<p>attempted division of $x^3 + x + 10$ by $(x + 2)$ as far as $x^3 + 2x^2$ in working</p> <p>correctly obtaining $x^2 - 2x + 5$</p> <p>use of $b^2 - 4ac$ with $x^2 - 2x + 5$</p> <p>$b^2 - 4ac = 4 - 20 [= -16]$</p> <p>so only two real roots [of $f(x)$] [and hence no more linear factors]</p>	<p>M1</p> <p>A1</p> <p>M1</p> <p>A1</p> <p>A1</p> <p>[5]</p>	<p>or $x^3 - 3x^2 + 7x - 5$ by $(x - 1)$ as far as $x^3 - x^2$ in working</p> <p>or inspection with at least two terms of quadratic factor correct</p> <p>allow these first 2 marks if this has been done in (ii), even if not used here</p> <p>may be in attempt at formula (ignore rest of formula)</p> <p>may be in formula;</p> <p>or no real roots of $x^2 - 2x + 5 = 0$; allow this last mark if clear use of $x^2 - 2x + 5 = 0$, even if error in $b^2 - 4ac$, provided result negative, but no ft from wrong factor</p> <p>if last M1 not earned, allow SC1 for stating that the only factors of 5 are 1 and 5 and reasoning eg that $(x - 1)(x - 5)$ and $(x + 1)(x + 5)$ do not give $x^2 - 2x + 5$ [hence $x^2 - 2x + 5$ does not factorise]</p>	<p>alt method: allow M1 for attempted division of quartic by $x^2 + x - 2$ as far as $x^4 + x^3 - 2x^2$ in working, or inspection etc</p> <p>or completing square form attempted</p> <p>or attempt at calculus or symmetry to find min pt</p> <p>NB M0 for use of $b^2 - 4ac$ with cubic factor etc</p> <p>or $(x - 1)^2 + 4$</p> <p>or min = (1, 4)</p> <p>or $(x - 1)^2 + 4$ is always positive so no real roots [of $(x - 1)^2 + 4 = 0$] [and hence no linear factors]</p> <p>or similar conclusion from min pt</p>

OCR (Oxford Cambridge and RSA Examinations)
1 Hills Road
Cambridge
CB1 2EU

OCR Customer Contact Centre

Education and Learning

Telephone: 01223 553998

Facsimile: 01223 552627

Email: general.qualifications@ocr.org.uk

www.ocr.org.uk

For staff training purposes and as part of our quality assurance programme your call may be recorded or monitored

Oxford Cambridge and RSA Examinations
is a Company Limited by Guarantee
Registered in England
Registered Office; 1 Hills Road, Cambridge, CB1 2EU
Registered Company Number: 3484466
OCR is an exempt Charity

OCR (Oxford Cambridge and RSA Examinations)
Head office
Telephone: 01223 552552
Facsimile: 01223 552553

© OCR 2013

