

GCE

Mathematics (MEI)

Advanced Subsidiary GCE

Unit 4761: Mechanics 1

Mark Scheme for January 2013

OCR (Oxford Cambridge and RSA) is a leading UK awarding body, providing a wide range of qualifications to meet the needs of candidates of all ages and abilities. OCR qualifications include AS/A Levels, Diplomas, GCSEs, Cambridge Nationals, Cambridge Technicals, Functional Skills, Key Skills, Entry Level qualifications, NVQs and vocational qualifications in areas such as IT, business, languages, teaching/training, administration and secretarial skills.

It is also responsible for developing new specifications to meet national requirements and the needs of students and teachers. OCR is a not-for-profit organisation; any surplus made is invested back into the establishment to help towards the development of qualifications and support, which keep pace with the changing needs of today's society.

This mark scheme is published as an aid to teachers and students, to indicate the requirements of the examination. It shows the basis on which marks were awarded by examiners. It does not indicate the details of the discussions which took place at an examiners' meeting before marking commenced.

All examiners are instructed that alternative correct answers and unexpected approaches in candidates' scripts must be given marks that fairly reflect the relevant knowledge and skills demonstrated.

Mark schemes should be read in conjunction with the published question papers and the report on the examination.

OCR will not enter into any discussion or correspondence in connection with this mark scheme.

© OCR 2013

Annotations

Annotation in scoris	Meaning
✓ and ×	
BOD	Benefit of doubt
FT	Follow through
ISW	Ignore subsequent working
M0, M1	Method mark awarded 0, 1
A0, A1	Accuracy mark awarded 0, 1
B0, B1	Independent mark awarded 0, 1
SC	Special case
٨	Omission sign
MR	Misread
Highlighting	

Other abbreviations in mark scheme	Meaning
E1	Mark for explaining
U1	Mark for correct units
G1	Mark for a correct feature on a graph
M1 dep*	Method mark dependent on a previous mark, indicated by *
сао	Correct answer only
oe	Or equivalent
rot	Rounded or truncated
soi	Seen or implied
www	Without wrong working

Subject-specific Marking Instructions for GCE Mathematics (MEI) Mechanics strand

a Annotations should be used whenever appropriate during your marking.

The A, M and B annotations must be used on your standardisation scripts for responses that are not awarded either 0 or full marks. It is vital that you annotate standardisation scripts fully to show how the marks have been awarded.

For subsequent marking you must make it clear how you have arrived at the mark you have awarded.

b An element of professional judgement is required in the marking of any written paper. Remember that the mark scheme is designed to assist in marking incorrect solutions. Correct *solutions* leading to correct answers are awarded full marks but work must not be judged on the answer alone, and answers that are given in the question, especially, must be validly obtained; key steps in the working must always be looked at and anything unfamiliar must be investigated thoroughly.

Correct but unfamiliar or unexpected methods are often signalled by a correct result following an *apparently* incorrect method. Such work must be carefully assessed. When a candidate adopts a method which does not correspond to the mark scheme, award marks according to the spirit of the basic scheme; if you are in any doubt whatsoever (especially if several marks or candidates are involved) you should contact your Team Leader.

c The following types of marks are available.

Μ

A suitable method has been selected and *applied* in a manner which shows that the method is essentially understood. Method marks are not usually lost for numerical errors, algebraic slips or errors in units. However, it is not usually sufficient for a candidate just to indicate an intention of using some method or just to quote a formula; the formula or idea must be applied to the specific problem in hand, eg by substituting the relevant quantities into the formula. In some cases the nature of the errors allowed for the award of an M mark may be specified.

Α

Accuracy mark, awarded for a correct answer or intermediate step correctly obtained. Accuracy marks cannot be given unless the associated Method mark is earned (or implied). Therefore M0 A1 cannot ever be awarded.

В

Mark for a correct result or statement independent of Method marks.

Е

A given result is to be established or a result has to be explained. This usually requires more working or explanation than the establishment of an unknown result.

Unless otherwise indicated, marks once gained cannot subsequently be lost, eg wrong working following a correct form of answer is ignored. Sometimes this is reinforced in the mark scheme by the abbreviation isw. However, this would not apply to a case where a candidate passes through the correct answer as part of a wrong argument.

- d When a part of a question has two or more 'method' steps, the M marks are in principle independent unless the scheme specifically says otherwise; and similarly where there are several B marks allocated. (The notation 'dep *' is used to indicate that a particular mark is dependent on an earlier, asterisked, mark in the scheme.) Of course, in practice it may happen that when a candidate has once gone wrong in a part of a question, the work from there on is worthless so that no more marks can sensibly be given. On the other hand, when two or more steps are successfully run together by the candidate, the earlier marks are implied and full credit must be given.
- e The abbreviation ft implies that the A or B mark indicated is allowed for work correctly following on from previously incorrect results. Otherwise, A and B marks are given for correct work only – differences in notation are of course permitted. A (accuracy) marks are not given for answers obtained from incorrect working. When A or B marks are awarded for work at an intermediate stage of a solution, there may be various alternatives that are equally acceptable. In such cases, exactly what is acceptable will be detailed in the mark scheme rationale. If this is not the case please consult your Team Leader.

Sometimes the answer to one part of a question is used in a later part of the same question. In this case, A marks will often be 'follow through'. In such cases you must ensure that you refer back to the answer of the previous part question even if this is not shown within the image zone. You may find it easier to mark follow through questions candidate-by-candidate rather than question-by-question.

f Unless units are specifically requested, there is no penalty for wrong or missing units as long as the answer is numerically correct and expressed either in SI or in the units of the question. (eg lengths will be assumed to be in metres unless in a particular question all the lengths are in km, when this would be assumed to be the unspecified unit.)

We are usually quite flexible about the accuracy to which the final answer is expressed and we do not penalise over-specification.

When a value is given in the paper

Only accept an answer correct to at least as many significant figures as the given value. This rule should be applied to each case.

When a value is not given in the paper

Accept any answer that agrees with the correct value to 2 s.f.

ft should be used so that only one mark is lost for each distinct error made in the accuracy to which working is done or an answer given. Refer cases to your Team Leader where the same type of error (eg errors due to premature approximation leading to error) has been made in different questions or parts of questions.

Mark Scheme

There are some mistakes that might be repeated throughout a paper. If a candidate makes such a mistake, (eg uses a calculator in wrong angle mode) then you will need to check the candidate's script for repetitions of the mistake and consult your Team Leader about what penalty should be given.

There is no penalty for using a wrong value for g. E marks will be lost except when results agree to the accuracy required in the question.

g Rules for replaced work

If a candidate attempts a question more than once, and indicates which attempt he/she wishes to be marked, then examiners should do as the candidate requests.

If there are two or more attempts at a question which have not been crossed out, examiners should mark what appears to be the last (complete) attempt and ignore the others.

NB Follow these maths-specific instructions rather than those in the assessor handbook.

h For a *genuine* misreading (of numbers or symbols) which is such that the object and the difficulty of the question remain unaltered, mark according to the scheme but following through from the candidate's data. A penalty is then applied; 1 mark is generally appropriate, though this may differ for some units. This is achieved by withholding one A mark in the question.

Marks designated as cao may be awarded as long as there are no other errors. E marks are lost unless, by chance, the given results are established by equivalent working.

'Fresh starts' will not affect an earlier decision about a misread.

Note that a miscopy of the candidate's own working is not a misread but an accuracy error.

- i If a graphical calculator is used, some answers may be obtained with little or no working visible. Allow full marks for correct answers (provided, of course, that there is nothing in the wording of the question specifying that analytical methods are required). Where an answer is wrong but there is some evidence of method, allow appropriate method marks. Wrong answers with no supporting method score zero. If in doubt, consult your Team Leader.
- j If in any case the scheme operates with considerable unfairness consult your Team Leader.

4761

Question		n	Answer	Marks	Guidance
1	(i)		Normal reaction P 3kg 10 N 30° 3g	B1 B1 B1 [3]	3 marks –1 / error or omission Forces must have arrows and labels Accept "weight" and "friction"
1	(ii)		$R = 3g\cos 30^\circ = 25.46 = 25.5 \text{ (to 3 significant figures)}$	B1 [1]	Accept 25 or 26
1	(iii)		$P = 10 + 3g\sin 30^{\circ}$ $P = 24.7$	M1 A1 [2]	Correct elements must be present Cao
2	(i)		$\mathbf{v} = \mathbf{u} + \mathbf{a}t$ Velocity $\mathbf{v} = \begin{pmatrix} 2\\ 0 \end{pmatrix} + t \begin{pmatrix} -1\\ 1 \end{pmatrix} \left(= \begin{pmatrix} 2-t\\ t \end{pmatrix} \right)$ When $t = 8$, $\mathbf{v} = \begin{pmatrix} -6\\ 8 \end{pmatrix}$ speed $\sqrt{(-6)^2 + 8^2} = 10 \text{ m s}^{-1}$	M1 A1 A1 A1 [4]	May be implied by either of the next two answers but not the final answer. Evidence of use of vectors in question necessary. May be implied by the final answer Cao but condone no units Give SC2 for 10 without working

Question		n	Answer	Marks	Guidance
2	(ii)		$\mathbf{r} = \mathbf{r}_0 + \mathbf{u}t + \frac{1}{2}\mathbf{a}t^2$	M1	Use of correct equation with substitution. Condone omission of $\mathbf{r}_{0.}$ Or equivalent equation
			$\mathbf{r} = \begin{pmatrix} 0 \\ -2 \end{pmatrix} + \begin{pmatrix} 2 \\ 0 \end{pmatrix} \times 8 + \frac{1}{2} \times \begin{pmatrix} -1 \\ 1 \end{pmatrix} \times 8^2$	A1	Condone omission of \mathbf{r}_0 . Follow through for their value of \mathbf{v}
			$\mathbf{r} = \begin{pmatrix} -16\\ 30 \end{pmatrix}$	A1	Cao but may be implied by a correct final answer.
			Distance = 34 m	A1	Allow for 35.77 from $\mathbf{r} = \begin{pmatrix} -16 \\ 32 \end{pmatrix}$ and 37.57 from $\mathbf{r} = \begin{pmatrix} -16 \\ 34 \end{pmatrix}$
				[4]	
3	(i)		$s = ut + \frac{1}{2}at^2$	M1	Substitution required
			$7.2 = \frac{1}{2} \times a \times 6^2$	A1	
			$a = 0.4 \text{ ms}^{-2}$	A1	Cao
				[3]	
3	(ii)		F = ma	M1	Attempt at Newton's second law
				M1	Attempt at resolving both <i>S</i> and <i>T</i>
			$300\cos 30^\circ + 175\cos 15^\circ - R = 1000 \times 0.4$	A1	(Correct elements present and no extras); follow through for a
			$R = 28.8 \mathrm{N}$	A1	Cao
				[4]	
3	(iii)		The resistance perpendicular to the line of motion has been	B1	Allow
			ignored.		There is also a sideways resistance force
				[1]	

(Question		Answer	Marks	Guidance	
4	(i)		Either $s = \frac{1}{2}(u+v)t$ Take O as the origin.	M1	Use of one relevant equation, including substitution	
			$30 = \frac{1}{2} \times (u+9) \times 10$			
			u = -3	A1		
			v = u + at	M1	Use of a second relevant equation including substitution	
			9 = -3 + 10a			
			<i>a</i> = 1.2	A1		
			or $v = u + at \implies u + 10a = 9$	M1	Use of one relevant equation, including substitution	
			$s = ut + \frac{1}{2}at^2 \implies u + 5a = 3$	M1	Use of a second relevant equation including substitution	
			Solving simultaneously: $a = 1.2$	A1		
			<i>u</i> = -3	A1		
			$\mathbf{or} s = vt - \frac{1}{2}at^2$	M1	Use of one relevant equation, including substitution	
			$\Rightarrow a = 1.2$	A1		
			v = u + at	M1	Use of a second relevant equation including substitution	
			$\Rightarrow u = -3$	A1		
				[4]		
4	(ii)		Either $s = ut + \frac{1}{2}at^2$			
			Solving for P: $-5 = -3t + \frac{1}{2} \times 1.2t^2$	M1	Quadratic equation with $s = -5$	
			$0.6t^2 - 3t + 5 = 0$			
			Discriminant = $3^2 - 4 \times 0.6 \times 5 = -3$	M1	Considering the discriminant or equivalent	
			No real roots for $t \implies Particle is never at P)$	E1	Cao without wrong working in the whole question.	
	[]			

⁴⁷⁶¹

(Question		Answer	Marks	Guidance
			Or Find when $v = 0$	M1	
			$v = u + at$, $v = 0 \implies t = 2.5$		
			$s = ut + \frac{1}{2}at^2$ and $t = 2.5$	M1	Or use $v^2 = u^2 + 2as$
			\Rightarrow s = -3.75 > -5	E1	Cao without wrong working in the whole question. Comparison necessary
			Special areas when their $u > 0$ and their $a > 0$	SC1	"It is always going to the right"
			Special cases when their $u > 0$ and their $u > 0$	SC1	Demonstration that it is at –5 for two negative times.
				[3]	
5					
5	(1)		Vertical motion: $s = ut + \frac{1}{2}at^2$		
			At water: $-1.225 = 0 \times t + \frac{1}{2} \times (-9.8) \times t^2$	M1	Condone sign errors
			$\Rightarrow t = 0.5 \text{ s}$	A1	Signs must be consistent
				[2]	
5	(ii)		Horizontal component of velocity = 20 m s^{-1}	B1	
			Vertical component = $0.5 \times 9.8 = 4.9 \text{ m s}^{-1}$	B1	Follow through for "their $t \ge 9.8$ "
			Speed = $\sqrt{20^2 + 4.9^2} = 20.6$	M1	Use of Pythagoras on previous two answers
			$\tan \alpha = \frac{4.9}{20}$	M1	Use of an appropriate trig ratio with their figures for v . Must be explicit if final answer is incorrect.
			$\alpha = 13.8^{\circ}$	A1	Cao
				[5]	

Question		n	Answer	Marks	Guidance
6	(i)	(A)	Distance travelled = Area under the graph	M1	Attempt to find area
			$\frac{1}{2} \times 4 \times 8 + \frac{1}{2} \times 4 \times (8 + 12) + 4 \times 12$	M1	Splitting into suitable parts
			104 m	A1	Cao
					Allow all 3 marks for 104 without any working
6	(i)	(<i>B</i>)	Either		
			Working backwards from distance when $t = 12$	M1	
			$12 - \frac{(104 - 100)}{12}$	M1	Allow this mark for 0.33 Follow through from their total distance
			11.67 s	A1	Cao
[Or		
			Working forwards from when $t = 8$	M1	
			$8 + \frac{(100 - 56)}{12}$	M1	Allow this mark for 3.67 Follow through from their distance at time 8s
			11.67 s	A1	Cao
				[6]	
6	(ii)		Substituting $t = 8$ gives $v = \frac{5}{2} \times 8 - \frac{1}{8} \times 8^2 = 12$	B1	
				[1]	

Question		n	Answer	Marks	Guidance
6	(iii)		Distance = $\int_0^{12} \left(\frac{5t}{2} - \frac{t^2}{8} \right) dt$	M 1	Integrating v. Condone no limits.
			$\left[\frac{5t^2}{4} - \frac{t^3}{24}\right]_0^{12}$	A1	Condone no limits
			[180 - 72] (- $[0]$)	M1	Substituting $t = 12$
			108 m	A1	
				[4]	
6	(iv)		Model P: distance at $t = 11.35$ is 96.2	B1	Сао
			Model Q: distance at $t = 11.35$ is		
			$\left[\frac{5t^2}{4} - \frac{t^3}{24}\right]_0^{11.35} = 100.1$	M1	Substituting 11.35 in their expression from part (iii)
			Model Q places the runner closer	E1	Cao from correct previous working for both models
				[3]	
6	(v)		Model P: Greatest acceleration $\frac{8}{4} = 2 \text{ m s}^{-2}$	B1	
			Model Q: $a = \frac{dv}{dt} = \frac{5}{2} - \frac{t}{4}$	M1	Differentiating v
				A1	
			Model Q: Greatest acceleration is 2.5 m s^{-2}	B1	Award if correct answer seen
				[4]	

(Questio	n	Answer	Marks	Guidance
7	(i)	(A)	The pulley is smooth	B1	Award for "smooth" seen.
				[1]	
7	(i)	(B)	Horizontal equilibrium: $T \sin \theta = T \sin \phi$	M1	Attempt at horizontal equilibrium. Allow sin-cos interchange. The argument must be based on forces.
			$\Rightarrow heta = \phi$	E1	Do not allow if sin-cos interchange
				[2]	
7	(ii)		Call M the mid point of AB. AM = 1, AC=1.4, $\angle AMC = 90^{\circ}$	M1	Setting up triangle and use of trigonometry
			Pythagoras \Rightarrow MC = $\sqrt{1.4^2 - 1^2} = \sqrt{0.96}$		
			$\sqrt{0.96}$ $\sqrt{24}$	E1	
			$\cos\theta = \frac{\sqrt{3}}{1.4} = \frac{\sqrt{2}}{7}$		If decimals are matched, at least 3 figures must be given
				[2]	
7	(iii)		Vertical equilibrium	M1	Use of vertical equilibrium
			$2T\cos\theta = 50$	A1	Accept $T \cos \theta = 25$ as an equivalent statement
			T = 35.7 N	A1	Cao
				[3]	
7	(iv)		$1.2^2 + 1.6^2 = 2^2$		
			$\Rightarrow \angle ACB = 90^{\circ}$	B1	Use of Pythagoras, or equivalent
			$\cos\alpha = 0.6, \ \cos\beta = 0.8$	B1	Both No marks for sin-cos interchange
				[2]	

Question		n	Answer	Marks	Guidance
7	(v)		Either resolving horizontally and vertically		
			$T_1 \cos \alpha = T_2 \cos \beta$	M1	Attempt at horizontal equation. Allow consistent sin-cos interchange
			$T_1 \sin \alpha + T_2 \sin \beta = 50$	M1	Attempt at vertical equation. Allow consistent sin-cos interchange
			$0.6T_1 = 0.8T_2$ $0.8T_1 + 0.6T_2 = 50$	A1	Substitution in both equations. Dependent on both M marks. Cao
			Solving simultaneously	M1	Dependent on both the previous M marks
			$T_1 = 40, \ T_2 = 30$	A1	Cao
			Or resolving in the direction of the strings		
			Resolving in both directions	M1	A serious attempt to use this method. Allow sin-cos interchange
			$T_1 = 50\sin\alpha$	M1	
			$\Rightarrow T_1 = 50 \times 0.8 = 40$	A1	
			$T_2 = 50 \times \sin \beta$	M1	
			$\Rightarrow T_2 = 50 \times 0.6 = 30$	A1	
			Or triangle of forces		
			Use of a triangle of forces	M1	The triangle must be closed and have a right angle opposite the weight
			Labels	M1	The sides must be correctly annotated
			Angles	M1	The angles must be correctly annotated
			$T_1 = 50 \times 0.8 = 40$	A1	Cao Dependent of first M mark
			$T_2 = 50 \times 0.6 = 30$	A1	Cao Dependent of first M mark
				[5]	

Question		n	Answer	Marks	Guidance
7	(vi)		Attempt to find $\angle CAB$	M1	May be implied by the remaining answers
			Tension in AC is 50 N (it takes all the weight)	B1	
			Tension in BC is zero (it is slack)	B1	
				[3]	

OCR (Oxford Cambridge and RSA Examinations) 1 Hills Road Cambridge CB1 2EU

OCR Customer Contact Centre

Education and Learning

Telephone: 01223 553998 Facsimile: 01223 552627 Email: general.qualifications@ocr.org.uk

www.ocr.org.uk

For staff training purposes and as part of our quality assurance programme your call may be recorded or monitored

Oxford Cambridge and RSA Examinations is a Company Limited by Guarantee Registered in England Registered Office; 1 Hills Road, Cambridge, CB1 2EU Registered Company Number: 3484466 OCR is an exempt Charity

OCR (Oxford Cambridge and RSA Examinations) Head office Telephone: 01223 552552 Facsimile: 01223 552553





© OCR 2013