

Mark Scheme (Results)

January 2016

Pearson Edexcel International A Level in Statistics 1 (WST01) Paper 01

Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at <u>www.edexcel.com</u> or <u>www.btec.co.uk</u>. Alternatively, you can get in touch with us using the details on our contact us page at <u>www.edexcel.com/contactus</u>.

Pearson: helping people progress, everywhere

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

January 2016 Publications Code **IA043346** All the material in this publication is copyright © Pearson Education Ltd 2016

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

PEARSON EDEXCEL IAL MATHEMATICS

General Instructions for Marking

- 1. The total number of marks for the paper is 75
- 2. The Edexcel Mathematics mark schemes use the following types of marks:
- **M** marks: Method marks are awarded for `knowing a method and attempting to apply it', unless otherwise indicated.
- A marks: Accuracy marks can only be awarded if the relevant method (M) marks have been earned.
- **B** marks are unconditional accuracy marks (independent of M marks)
- Marks should not be subdivided.
- 3. Abbreviations

These are some of the traditional marking abbreviations that will appear in the mark schemes.

- bod benefit of doubt
- ft follow through
- the symbol $\sqrt{}$ will be used for correct ft
- cao correct answer only
- cso correct solution only. There must be no errors in this part of the question to obtain this mark
- isw ignore subsequent working
- awrt answers which round to
- SC: special case
- oe or equivalent (and appropriate)
- d... or dep dependent
- indep independent
- dp decimal places
- sf significant figures
- * The answer is printed on the paper or ag- answer given
- 4. All A marks are 'correct answer only' (cao.), unless shown, for example, as A1 ft to indicate that previous wrong working is to be followed through. After a misread however, the subsequent A marks affected are treated as A ft, but manifestly absurd answers should never be awarded A marks.

- 5. For misreading which does not alter the character of a question or materially simplify it, deduct two from any A or B marks gained, in that part of the question affected.
- 6. If a candidate makes more than one attempt at any question:
 - If all but one attempt is crossed out, mark the attempt which is NOT crossed out.
 - If either all attempts are crossed out or none are crossed out, mark all the attempts and score the highest single attempt.
- 7. Ignore wrong working or incorrect statements following a correct answer.

Special notes for marking Statistics exams

- If a candidate is "hedging their bets" e.g. give Attempt 1...Attempt 2...etc then please send to review.
- Any correct method should gain credit. If you cannot see how to apply the mark scheme but believe the method to be correct please send to review.

Jan2016 WST01 STATISTICS 1 International Mark Scheme

Question Scheme Marke		
Number	Scheme	Marks
1. (a)	$[F(5) =] \frac{5}{6}$	B1
		(1)
(b)	$\begin{bmatrix} E(X) = \end{bmatrix} -2 \times \frac{1}{4} + 1 \times \frac{1}{6} + 3 \times \frac{1}{3} + 4 \times \frac{1}{12} + 6 \times \frac{1}{6} \text{ or } \frac{1}{12} \left(-6 + 2 + 12 + 4 + 12 \right)$	M1
	= <u>2</u>	A1 (2)
(c)	$[E(X^{2}) =] (-2)^{2} \times \frac{1}{4} + 1 \times \frac{1}{6} + 3^{2} \times \frac{1}{3} + 4^{2} \times \frac{1}{12} + 6^{2} \times \frac{1}{6} $ (o.e.)	
		M1
	$\underline{\text{or}} \frac{1}{12} (12 + 2 + 36 + 16 + 72) \underline{\text{or}} \frac{138}{12} \underline{\text{or}} \frac{23}{2} (\text{o.e.})$	
	$[Var(X) =] "\frac{23}{2}" - "2"^2$	M1
	= <u>7.5</u> (o.e.)	A1 (3)
(d)(i)	[E(Y) = 7 - 2E(X) =] <u>3</u>	B1 (3)
(ii)	$[\operatorname{Var}(Y) =] (-2)^2 \operatorname{Var}(X) \underline{\text{or}} 4 \operatorname{Var}(X)$	M1
(***)	$=\underline{30}$	Al
(iii)	$7 - 2X \ge X \implies 7 \ge 5X$ so $X = 1$ or -2	M1 A1
	So $[P(Y > X) =] \frac{5}{12}$	A1
	12	(6)
(a)	NotesB1for $\frac{5}{6}$ or exact equivalents e.g. $\frac{10}{12}$ or $0.8\dot{3}$	
(a)	$\frac{1}{6}$ of exact equivalents e.g. $\frac{1}{12}$ or 0.05	
	For M1 in (b) & 1 st M1 in (c) take <u>full</u> method leading to answer: e.g. \div by 5 (or	
(b)	M1 for an attempt at $E(X)$ with at least 3 correct products seen. Answer of	nly M1A1
(c)	1 st M1 for an attempt at $E(X^2)$ with at least 3 correct products seen (even if 1	abelled Var(X))
	Condone -2^2 if it later becomes $+4$ but only 3 correct products are no	
	2^{nd} M1 for a correct numerical expression for Var(X) ft their E(X) and their E	(X^2)
	A1 for 7.5 or any exact equivalent e.g. $\frac{15}{2}$ Answer only M1M1A1	
(d)(i)	B1 for 3 only	
(ii)	M1 for correct use of $Var(aX+b)$ formula. If their value of $Var(X)$ even	if < 0
(iii)	A1 for 30 only M1 for attempt at solving a correct inequality as far as $a > bx$ (a and b be	oth >0)
(m)	1 st A1 for identifying $X = 1$ and -2 as the required values (or $Y = 11$ and 5) 2 nd A1 for $\frac{5}{12}$ or an exact equivalent	/
	[$y: 11 \ 5 \ 1 \ -1 \ -5$] Allow M1 in (ii) for attempt at $E(Y^2) = 39$ (≥ 3 corr	ect products)
ALT	and $Var(Y) = "39" - "3"^2$ ft their $E(Y)$	
	Allow M1 in (iii) for attempt at full set of Y values with corresponding X probs (\geq 3 c	correct)

Question Number	Sche	eme	Marks
Question Number	Scheme		Marks
2. (a)	[pass for] <u>30</u>	(labelled or 1 st answer)	B1 (1
(b)	[merit for] <u>46</u>	(labelled or 2 nd answer)	B1 (1
(c)	$[1.5(Q_3 - Q_1) = 1.5 \times 16 = 24]$ so $c = \underline{7}$	$\underline{0}$ and $d = \underline{6}$	B1, B1
(d)	<u>68, 72, 79</u>		B2/1/0 (2
(e)	5 < d therefore 5 is an outlier		(2 M1
	* 1 10 20 30 40	i * * 50 60 70 80	A1 A1
(f)	$\frac{1}{2} \times \left(\frac{1}{4}\right)^2 \times 3$	*********************	(3 M1M1
	$=\frac{3}{32}$		A1
			(3 [12 marks]
			0/ for (a)
(c) SC	$1^{\text{st}} B1$ for $c = 70$ $2^{\text{nd}} B1$ for $d =$	The correct numbers with % e.g. 30 6 (Allow B1B1 for unlabelled 70 f d = 70 or 6 and 70 in the wrong ord	followed by 6)
(d)	B2 for all 3 correct values (and no	extra value) (B1 for two correct)	
(e)	M1 for identifying or stating(e.g. or ending at 6 or 10 (May be impl 1 st A1 for only <u>one</u> outlier correctly m 2 nd A1 for a <u>single</u> lower whisker stop Condone 15 marked (e.g. dash	arked at 5 (whisker(s) must stop ab	<u>bove</u> 5)
		is a common incorrect answer and	
SC	Can award M0M1A0 if just $\frac{3}{6}$ A1 for $\frac{3}{32}$ or exact equivalent. All	$\frac{1}{4}$ (o.e.) is seen. ow 0.0937 or 0.0938 following a co	rrect expression.

Question Number	Scheme	Marks
3. (a)	$[S_{vs}] = 177.311 - \frac{36.8 \times 29}{8} = 43.911 = $ awrt <u>43.9</u> $[S_{ss}] = 209.72 - \frac{36.8^2}{8} = 40.44 = $ awrt <u>40.4</u>	A1 M1
	$[S_{ss}] = 209.72 - \frac{36.8^2}{8} = 40.44 = $ awrt <u>40.4</u>	Al
(b)	$r = \frac{"43.911"}{\sqrt{55.275 \times "40.44"}}$, = 0.92875 = awrt <u>0.929</u>	(3) M1, A1
(c)	<i>r</i> is close to 1 so there is support for the publisher's belief [if $1 > r/ 0.5$] (Allow "yes" because "strong corr." <u>but</u> "yes" & "positive corr." is B0)	(2) B1ft
(d)	$b = \frac{"43.911"}{55.275}, = 0.7944 = awrt 0.79$	(1) M1, A1
	$a = s - bv = 4.6 - 0.7944 \times 3.625 [= 1.720]$ $\underline{s = 1.72 + 0.794v}$	M1 A1 (4)
(e)	$\frac{y}{1000} = "1.72" + "0.794" \times \left(\frac{x+50}{200}\right)$ $y = 1920 + 3.97x$	M1 A1 A1ft
(f)	Gradient of textbooks is greater spend more advertising on textbooks	(3) B1ft dB1ft (2)
	Notes	[15 marks]
(a)	M1for one correct expressionFor correct answer with 1^{st} A1for $[S_{vs} =]$ awrt 43.9award M1 and the appro 2^{nd} A1for $[S_{ss} =]$ awrt 40.4Condone missing labels	priate A1
(b)	M1 for a correct expr' for <i>r</i> , ft their 43.911 (but not 177.311) and their 40.44 A1 for awrt 0.929 (correct ans only scores 2/2 and ans only of 0.93 scores 1	
(c)	B1ft for saying it <u>does</u> support the belief <u>or</u> a linear model/relationship is su and giving a suitable reason e.g. <u>strong</u> correlation [If $ r < 0.5$ allow "r close to 0" so "does <u>not</u> support" o.e.]	uitable
(d)	In (d) and (e) a correct answer with no working is awarded the M marks by implication 1^{st} M1 for a correct expression for <i>b</i> , ft their 43.911 and allow 3sf values to be used 1^{st} A1 for awrt 0.79 or allow an exact fraction from the 3sf values e.g. $\frac{439}{553}$	
	2 nd M1 for a correct method for <i>a</i> , ft their value of <i>b</i> NB $\overline{s} = 4.6 = \frac{3.6}{8}$ and 2 nd A1 for equation for <i>s</i> in terms of <i>v</i> with <i>a</i> = awrt 1.72 and <i>b</i> = awrt 0.79	0
(e)	M1 for correct sub. in their equation giving an equation in y and x. Allow 1^{st}A1 for $c = 1920$ (to 3 sf)	w 1 slip e.g. $\frac{y}{100}$
ALT	2^{nd} A1ft for $d = awrt 3.97$ or $5 \times (their b correct to 2 sig. figs.)$ Using coding formulae to get values for x, y requires a <u>full</u> method. Allow 1 slip but	correct $d \Rightarrow$ M1
(f)	1 st B1ft for a suitable reason based on gradients (o.e. in words e.g. rate of i	
ft	2^{nd} dB1ft for recommending spend more on advertising textbooks If gradient in (e) < 1.2 then a <u>comparison of grads</u> leading to spending on n	ovels is B1B1

Question Number	Scheme	Marks
4. (a)	0.7 Pass 0.3 Fail 0.5 Pass 0.5 Fail 0.3 Pass 0.5 Fail 0.3 Pass 0.1 Pass 0.9 Fail 0.9 Fail	B1 B1
(b)		(2) M1 A1
(c)	$\left[P(P_1 \cup P_2 \text{Pass}) = \right] \frac{0.7 + "0.3" \times 0.5}{(b)}, = \frac{0.85}{"0.9055"} = 0.938707 = \text{awrt } 0.939$	(2) M1, A1ft A1
(d)		(3) M1 dM1A1cso (3)
(e)	$p = \frac{2.2 \pm \sqrt{2.2^2 - 4 \times 1.15}}{2} \underline{\text{or Complete the sq:}} (p - 1.1)^2 - 1.1^2 + 1.15 = 0$ $= \frac{2.2 \pm 0.4898}{2} \underline{\text{or}} \frac{2.2 \pm \sqrt{0.24}}{2} \underline{\text{or}} 1.1 \pm \sqrt{0.06} \underline{\text{or}} (1.34), 0.855$	M1 A1
	$p = 0.85505102 \ p = 0.855$	A1 (3)
(a)	Notes1st B1for correctly placing 0.3 and 0.52nd B1for correctly placing 0.7, 0.1 and 0.9	
(b)	 Apart from (d), a correct answer with no incorrect working scores fu M1 for a correct expression (ft from their tree diagram) A1 for 0.9055 or exact equivalent e.g. ¹⁸¹¹/₂₀₀₀ Accept 0.906 <u>only</u> if correct 	
(c)	M1 for a correct ratio of probs ft their 0.3 and their answer to (b)[if < 1]. N A1ft for correct numerator and their part (b) on denominator A1 for awrt 0.939 or accept exact fraction eg $\frac{1700}{1811}$	um > Den M0
(d)	1^{st} M1for a correct expression for P(pass) in terms of $p[$ condone $p - (p - 2^{nd} dM1)$ 2^{nd} dM1dep. on 1^{st} M1 for expanding brackets and forming an equation in p Allow one slipAlcsocorrect processing leading to printed answer. No incorrect working	
(e)	1^{st} A1for correct expression and simplified square root or 1.34 and 0.855 2^{nd} A1for $p = 0.855$ only (penalise any extra value > 1) Correct ans only so	 cores 3/3
Ans. only		

Question Number	Scheme	Marks
5. (a)	$\left[P(H < 18) = \right] P\left(Z < \frac{18 - 22}{10} \right) = P(Z < -0.4)$	M1
	= 1 - 0.6554	dM1
	= 0.3446 or awrt <u>0.345</u>	A1 (3)
(b)		M1 A1
	P(H > 39) = P(Z > 1.7) = 1 - 0.9554 = 0.0446 $P(H > 50 + H = 20) P(H > 50) "0.0026"$	A1
	$P(H > 50 H > 39) = \frac{P(H > 50)}{P(H > 39)} \text{or} \frac{"0.0026"}{"0.0446"}$	M1
	$= 0.057 \sim 0.0585$	A1 (5)
(c)	$\frac{18 - \mu}{\sigma} = -0.8416 \qquad \frac{28 - \mu}{\sigma} = 1$	M1B1A1
	Solving: $\sigma = 1.8416\sigma$	M1
	$\sigma = \operatorname{awrt} 5.43$	A1
	$\mu = awrt 22.57$	A1 (6)
		[14 marks]
	Notes	
(a)	1 st M1 for standardising with 18, 22 and 10. Allow $\pm \frac{18-22}{10}$	
	2 nd dM1 dependent on 1 st M1 for $1 - p$ where $0.6A1 for 0.3446 or better or awrt 0.345. NB Calculator gives 0.3445783$	Ans only 3/3
(b)	1 st M1 for correct standardisation and $1 - q$ (where $q = 0.9$) for one of thes	se probs
	1^{st} A1 for 0.0026 or better (calc 0.0025551) or $1 - 0.9974$ (or better) 2^{nd} A1 for 0.0446 or better (calc 0.0445654) or $1 - 0.9554$ (or better)	
	2^{nd} M1 for a correct ratio of probability expressions or values	
	(ft their 0.0026 and 0.0446 but if num. > denom. then M0)	
	3^{rd} A1 for answer in the range 0.057~0.0585. No fractions but $\frac{13}{223}$ can score N	
Ans. only	Can score full marks for either awrt 0.0583 (tables) or awrt 0.0573 (c	alc) only
(c)	1 st M1 for attempt to standardise with μ , σ and 18 or 28 and set equal to a z v The z values should be in the range (0.8, 0.0) for "18" and (0.05, 1.05)	
	The z values should be in the range $(0.8, 0.9)$ for "18" and $(0.95, 1.05)$ B1 for using $z = 0.8416$ or better (allow <u>+</u>) Calculator gives 0.8416212	/
	1 st A1 for both equations with ± 1 and ± 0.84 or better	
SC	for $\frac{28-\mu}{\sigma} = \pm 0.8416$ and $\frac{18-\mu}{\sigma} = \pm 1$ award M1B1A0 (0.84 instead of 0.	8416 loses B1)
	2^{nd} M1 for solving their linear equations in $\mu \& \sigma$. Reducing to an equation i	
	Correct processes allow one sign slip 2^{nd} A1 for σ = awrt 5.43	
	2 rd A1 for σ = awrt 5.43 3 rd A1 for μ = awrt 22.57	
Calc	No $z = 0.8416$ or better seen: can award 6/6 for $\sigma = awrt 5.4300$ or 5.4301 and μ	= awrt 22.57
No	For σ = awrt 5.43 and μ = awrt 22.57 award M1B0A1M1A1A1 i.e. 5/6	
working	1 or $0 = awnt 5.45$ and $\mu = awnt 22.57$ awald with 0Anwith ATAT i.e. $3/0$	

Question Number	Scheme	Marks
6. (a)	$(\mu \text{ or } \overline{x} =) \frac{8360}{10} = \underline{836}$	B1
	$(\mu \text{ or } \overline{x} =) \frac{8360}{10} = \underline{836}$ $(\sigma =) \sqrt{\frac{\sum (x - \overline{x})^2}{10}} = \sqrt{6384} \text{ or } 4\sqrt{399}, = 79.89993} \text{ awrt } \underline{79.9}$	M1, A1
(b)	mean > median So <u>positive</u> (skew)	(3) B1 dB1 (2)
(c)	$\frac{776+896}{2} = 836$ which is the same as \overline{x} or one is 60 above \overline{x} , one 60 below	B1
	So <u>no change</u> in the mean	dB1 (2)
(d)	$(896-836)^2 = (776-836)^2 = 60^2 = 3600 < 6384$ the average of $\sum (x-\overline{x})^2$	B1
	$\underline{\text{Or}} \sum (x - \overline{x})^2 \to 63840 + 2 \times 60^2 = 71040 \text{ and } \frac{71040}{12} = 5920 < \frac{63840}{10}$	
	So standard deviation will <u>reduce</u>	dB1
		(2) [9 marks]
	Notes	
(a)	M1 for $\frac{63840}{10}$ with or without $$ (ignore labels) or $s^2 = \frac{63840}{9}$	
	NB $\sum x^2 = 7052800$ but must see at least $\sigma^2 = \frac{7052800}{10} - ("836")^2$ for N	/ 1
	A1 for awrt 79.9 Accept $s = awrt 84.2$ (84.2219). Correct answer only M	11A1
(b)	1 st B1 for a correct comparison of mean and median (allow just $836 > 815$)	
	May see $\frac{k(\text{mean} - \text{median})}{\sigma \text{ or } \sigma^2}$ (o.e.) if so just check sign of answer (provide	ed denom > 0)
SC	 2nd dB1 dependent on 1st B1 for positive (skew) only. Positive correlation is If their mean is < 815 award B0B1 for the comparison <u>and</u> statement of new 	
(c)	1 st B1 for a suitable calculation to show(or statement) that mean of these t	
	all 12) is the same e.g. new $\sum x = 8360 + 776 + 896 = 10032$, so mean =	$=\frac{10032}{12}=836$
SC	2 nd dB1 dependent on a suitable calculation or reason for stating "no change If they only say differences are the same (but not 1 above and 1 below no change then award B0B1	
(d)	1 st B1 for a suitable calculation showing 60 or 3600 and comparing with 79 respectively (must see some calculation here) or calculation of new variance (5920 vs 6384) or st. dev (76.9 vs 79.9	
	or calculation of new variance (5920 vs 6384) or st. dev (76.9 vs 79.9 2 nd dB1 dependent on 1 st B1 for stating s.d. "reduces" (o.e.)	")
Use of Σx^2	Send arguments based on Σx^2 to review	

Pearson Education Limited. Registered company number 872828 with its registered office at 80 Strand, London WC2R ORL