



1. Two particles  $P$  and  $Q$  have mass 0.4 kg and 0.6 kg respectively. The particles are initially at rest on a smooth horizontal table. Particle  $P$  is given an impulse of magnitude 3 N s in the direction  $PQ$ .

(a) Find the speed of  $P$  immediately before it collides with  $Q$ . **(3)**

Immediately after the collision between  $P$  and  $Q$ , the speed of  $Q$  is  $5\text{ m s}^{-1}$ .

(b) Show that immediately after the collision  $P$  is at rest. **(3)**

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---







4. A car is moving along a straight horizontal road. The speed of the car as it passes the point  $A$  is  $25\text{ m s}^{-1}$  and the car maintains this speed for  $30\text{ s}$ . The car then decelerates uniformly to a speed of  $10\text{ m s}^{-1}$ . The speed of  $10\text{ m s}^{-1}$  is then maintained until the car passes the point  $B$ . The time taken to travel from  $A$  to  $B$  is  $90\text{ s}$  and  $AB = 1410\text{ m}$ .

(a) Sketch, in the space below, a speed-time graph to show the motion of the car from  $A$  to  $B$ .

(2)

(b) Calculate the deceleration of the car as it decelerates from  $25\text{ m s}^{-1}$  to  $10\text{ m s}^{-1}$ .

(7)

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---


---

---





5.



**Figure 1**

Two forces **P** and **Q** act on a particle at a point *O*. The force **P** has magnitude 15 N and the force **Q** has magnitude *X* newtons. The angle between **P** and **Q** is  $150^\circ$ , as shown in Figure 1. The resultant of **P** and **Q** is **R**.

Given that the angle between **R** and **Q** is  $50^\circ$ , find

(a) the magnitude of **R**, (4)

(b) the value of *X*. (5)

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---


---

---

---



6.



**Figure 2**

A plank  $AB$  has mass  $12 \text{ kg}$  and length  $2.4 \text{ m}$ . A load of mass  $8 \text{ kg}$  is attached to the plank at the point  $C$ , where  $AC = 0.8 \text{ m}$ . The loaded plank is held in equilibrium, with  $AB$  horizontal, by two vertical ropes, one attached at  $A$  and the other attached at  $B$ , as shown in Figure 2. The plank is modelled as a uniform rod, the load as a particle and the ropes as light inextensible strings.

(a) Find the tension in the rope attached at  $B$ . (4)

The plank is now modelled as a non-uniform rod. With the new model, the tension in the rope attached at  $A$  is  $10 \text{ N}$  greater than the tension in the rope attached at  $B$ .

(b) Find the distance of the centre of mass of the plank from  $A$ . (6)

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---













