

Mathematics

Advanced GCE

Unit **4723**: Core Mathematics 3

Mark Scheme for June 2011

OCR (Oxford Cambridge and RSA) is a leading UK awarding body, providing a wide range of qualifications to meet the needs of pupils of all ages and abilities. OCR qualifications include AS/A Levels, Diplomas, GCSEs, OCR Nationals, Functional Skills, Key Skills, Entry Level qualifications, NVQs and vocational qualifications in areas such as IT, business, languages, teaching/training, administration and secretarial skills.

It is also responsible for developing new specifications to meet national requirements and the needs of students and teachers. OCR is a not-for-profit organisation; any surplus made is invested back into the establishment to help towards the development of qualifications and support which keep pace with the changing needs of today's society.

This mark scheme is published as an aid to teachers and students, to indicate the requirements of the examination. It shows the basis on which marks were awarded by Examiners. It does not indicate the details of the discussions which took place at an Examiners' meeting before marking commenced.

All Examiners are instructed that alternative correct answers and unexpected approaches in candidates' scripts must be given marks that fairly reflect the relevant knowledge and skills demonstrated.

Mark schemes should be read in conjunction with the published question papers and the Report on the Examination.

OCR will not enter into any discussion or correspondence in connection with this mark scheme.

© OCR 2011

Any enquiries about publications should be addressed to:

OCR Publications
PO Box 5050
Annesley
NOTTINGHAM
NG15 0DL

Telephone: 0870 770 6622
Facsimile: 01223 552610
E-mail: publications@ocr.org.uk

1 (i)	Obtain integral of form ke^{2x+1}	M1	any non-zero constant k different from 6; using substitution $u = 2x + 1$ to obtain ke^u earns M1 (but answer to be in terms of x)
	Obtain correct $3e^{2x+1}$	A1	or equiv such as $\frac{6}{2}e^{2x+1}$
(ii)	Obtain integral of form $k_1 \ln(2x+1)$	M1	any non-zero constant k_1 ; allow if brackets absent; $k_1 \ln u$ (after sub'n) earns M1
	Obtain correct $5\ln(2x+1)$	A1	or equiv such as $\frac{10}{2}\ln(2x+1)$; condone brackets rather than modulus signs but brackets or modulus signs must be present (so that $5\ln 2x+1$ earns A0)
	Include $\dots + c$ at least once	B1	5 anywhere in the whole of question 1; this mark available even if no marks awarded for integration

5

2	Apply one of the transformations correctly to their equation	B1	
	Obtain correct $-3\ln x + \ln 4$	B1	or equiv
	Show at least one logarithm property	M1	correctly applied to their equation of resulting curve (even if errors have been made earlier)
	Obtain $y = \ln(4x^{-3})$	A1	4 or equiv of required form; $\ln 4x^{-3}$ earns A1; correct answer only earns 4/4; condone absence of $y =$

4

3 (a)	State $14\sin \alpha \cos \alpha = 3\sin \alpha$	B1	or unsimplified equiv such as $7(2\sin \alpha \cos \alpha) = 3\sin \alpha$
	Attempt to find value of $\cos \alpha$	M1	by valid process; may be implied
	Obtain $\frac{3}{14}$	A1	3 exact answer required; ignore subsequent work to find angle

(b)	Attempt use of identity for $\cos 2\beta$	M1	of form $\pm 2\cos^2 \beta \pm 1$; initial use of $\cos^2 \beta - \sin^2 \beta$ needs attempt to express $\sin^2 \beta$ in terms of $\cos^2 \beta$ to earn M1
	Obtain $6\cos^2 \beta + 19\cos \beta + 10$	A1	or unsimplified equiv or equiv involving $\sec \beta$
	Attempt solution of 3-term quadratic eqn	M1	for $\cos \beta$ or (after adjustment) for $\sec \beta$
	Use $\sec \beta = \frac{1}{\cos \beta}$ at some stage	M1	or equiv
	Obtain $-\frac{3}{2}$	A1	5 or equiv; and (finally) no other answer

8

4	(i)	Draw sketch of $y = (x-2)^4$	*B1	touching positive x -axis and extending at least as far as the y -axis; no need for 2 or 16 to be marked; ignore wrong intercepts
		Draw straight line with positive gradient	*B1	at least in first quadrant and reaching positive y -axis; assess the two graphs independently of each other
		Indicate two roots	B1 3	AG; dep *B *B and two correct graphs which meet on the y -axis; indicated in words or by marks on sketch
		[SC: Draw sketch of $y = (x-2)^4 - x - 16$ and indicate the two roots : B1 (i.e. max 1 mark)]		
<hr/>				
	(ii)	State 0 or $x = 0$	B1 1	not merely for coordinates (0, 16)
<hr/>				
	(iii)	Obtain correct first iterate	B1	to at least 3 dp; any starting value (> -16)
		Show correct iteration process	M1	producing at least 3 iterates in all; may be implied by plausible converging values
		Obtain at least 3 correct iterates	A1	allowing recovery after error; iterates given to only 3 d.p. acceptable; values may be rounded or truncated
		Obtain 4.118	A1 4	answer required to exactly 3 dp; A0 here if number of iterates is not enough to justify 4.118; attempt consisting of answer only earns 0/4
	[0 \rightarrow 4 \rightarrow 4.114743 \rightarrow 4.117769 \rightarrow 4.117849; 1 \rightarrow 4.030543 \rightarrow 4.115549 \rightarrow 4.117790 \rightarrow 4.117849; 2 \rightarrow 4.059767 \rightarrow 4.116321 \rightarrow 4.117811 \rightarrow 4.117850; 3 \rightarrow 4.087798 \rightarrow 4.117060 \rightarrow 4.117830 \rightarrow 4.117850; 4 \rightarrow 4.114743 \rightarrow 4.117769 \rightarrow 4.117849 \rightarrow 4.117851; 5 \rightarrow 4.140695 \rightarrow 4.118452 \rightarrow 4.117867 \rightarrow 4.117851]			
8				

5	Attempt use of product rule	*M1	to produce $k_1 x \ln(4x-3) + \frac{k_2 x^2}{4x-3}$ form
	Obtain $2x \ln(4x-3)$	A1	
	Obtain $\dots + \frac{4x^2}{4x-3}$	A1	or equiv
	Attempt second use of product rule	*M1	
	Attempt use of quotient (or product) rule	*M1	allow numerator the wrong way round
	Obtain		
	$2 \ln(4x-3) + \frac{8x}{4x-3} + \frac{8x(4x-3) - 16x^2}{(4x-3)^2}$	A1	or equiv
	Substitute 2 into attempt at second deriv	M1	dep *M *M *M
	Obtain $2 \ln 5 + \frac{96}{25}$	A1 8	or exact equiv consisting of two terms
	8		

6 Method 1: (Differentiation; assume value $\frac{10}{3}$; eqn of tangent; through origin)

Differentiate to obtain $k(3x-5)^{-\frac{1}{2}}$	M1	any constant k
Obtain $\frac{3}{2}(3x-5)^{-\frac{1}{2}}$	A1	or equiv
Attempt to find equation of tangent at P and attempt to show tangent passing through origin	M1	assuming value $\frac{10}{3}$; or equiv
Obtain $y = \frac{3}{2\sqrt{5}}x$ and confirm that tangent passes through O	A1	AG; necessary detail needed

Method 2: (Differentiation; equate $\frac{y \text{ change}}{x \text{ change}}$ to deriv; solve for x)

Differentiate to obtain $k(3x-5)^{-\frac{1}{2}}$	M1	any constant k
Obtain $\frac{3}{2}(3x-5)^{-\frac{1}{2}}$	A1	or equiv
Equate $\frac{y \text{ change}}{x \text{ change}}$ to deriv and attempt solution	M1	
Obtain $\frac{\sqrt{3x-5}}{x} = \frac{3}{2}(3x-5)^{-\frac{1}{2}}$ and solve to obtain $\frac{10}{3}$ only	A1	

Method 3: (Differentiation; find x from $y = f'(x) \cdot x$ and $y = \sqrt{3x-5}$)

Differentiate to obtain $k(3x-5)^{-\frac{1}{2}}$	M1	any constant k
Obtain $\frac{3}{2}(3x-5)^{-\frac{1}{2}}$	A1	or equiv
State $y = \frac{3}{2}(3x-5)^{-\frac{1}{2}}x$, $y = \sqrt{3x-5}$, eliminate y and attempt solution	M1	condone this attempt at 'eqn of tangent'
Obtain $\frac{10}{3}$ only	A1	

Method 4: (No differentiation; general line through origin to meet curve at one point only)

Eliminate y from equations $y = kx$ and

$y = \sqrt{3x-5}$ and attempt formation of quadratic eqn	M1	
Obtain $k^2x^2 - 3x + 5 = 0$	A1	or equiv
Equate discriminant to zero to find k	M1	
Obtain $k = \frac{3}{2\sqrt{5}}$ or equiv and confirm $x = \frac{10}{3}$	A1	

Method 5: (No differentiation; use coords of P to find eqn of OP ; confirm meets curve once)

Use coordinates $(\frac{10}{3}, \sqrt{5})$ to obtain $y = \frac{3\sqrt{5}}{10}x$ or equiv as equation of OP	B1	
Eliminate y from this eqn and eqn of curve and attempt quadratic eqn	M1	should be $9x^2 - 60x + 100 = 0$ or equiv
Attempt solution or attempt discriminant	M1	
Confirm $\frac{10}{3}$ only or discriminant = 0	A1	

Either:

Integrate to obtain $k(3x-5)^{\frac{3}{2}}$	*M1	any constant k
Obtain correct $\frac{2}{9}(3x-5)^{\frac{3}{2}}$	A1	
Apply limits $\frac{5}{3}$ and $\frac{10}{3}$	M1	dep *M; the right way round
Make sound attempt at triangle area and calculate (triangle area) minus (their area under curve)	M1	or equiv
Obtain $\frac{10}{6}\sqrt{5} - \frac{10}{9}\sqrt{5}$ and hence $\frac{5}{9}\sqrt{5}$	A1	9 or exact equiv involving single term
<u>Or:</u>		
Arrange to $x = \dots$ and integrate to obtain $k_1y^3 + k_2y$ form	*M1	
Obtain $\frac{1}{9}y^3 + \frac{5}{3}y$	A1	
Apply limits 0 and $\sqrt{5}$	M1	dep *M; the right way round
Make sound attempt at triangle area and calculate (their area from integration) minus (triangle area)	M1	
Obtain $\frac{20}{9}\sqrt{5} - \frac{5}{3}\sqrt{5}$ and hence $\frac{5}{9}\sqrt{5}$	A1	(9) or exact equiv involving single term

9

- 7 (i) Either: Attempt solution of at least one linear eq'n of form $ax + b = 12$ M1
- Obtain $\frac{1}{3}$ A2 **3** and (finally) no other answer
- Or: Attempt solution of 3-term quadratic eq'n obtained by squaring attempt at $g(x+2)$ on LHS and squaring 12 or -12 on RHS M1
- Obtain $\frac{1}{3}$ A2 **(3)** and (finally) no other answer
-
- (ii) Either: Obtain $3(3x+5)+5$ for h B1
- Attempt to find inverse function M1 of function of form $ax + b$
- Obtain $\frac{1}{9}(x-20)$ A1 **3** or equiv in terms of x
- Or: State or imply g^{-1} is $\frac{1}{3}(x-5)$ B1
- Attempt composition of g^{-1} with g^{-1} M1
- Obtain $\frac{1}{9}(x-5) - \frac{5}{3}$ A1 **(3)** or more simplified equiv in terms of x
-
- (iii) State $x \leq 0$ B2 **2** give B1 for answer $x < 0$

8

8	(i)	Differentiate to obtain form $ke^{-0.014t}$	M1	any constant k different from 400
		Obtain $5.6e^{-0.014t}$ or $-5.6e^{-0.014t}$	A1	or (unsimplified) equiv
		Obtain 4.9 or -4.9 or 4.87 or -4.87	A1	3 but not greater accuracy; allow if final statement seems contradictory; answer only earns 0/3 – differentiation is needed

	(ii)	<u>Either:</u> State or imply $M_2 = 75e^{kt}$	B1	or equiv
		Attempt to find formula for M_2	M1	
		Obtain $M_2 = 75e^{0.047t}$	A1	or equiv such as $75e^{(\frac{1}{10}\ln\frac{8}{5})t}$
		Equate masses and attempt rearrangement	M1	as far as equation with e appearing once
		Obtain $e^{0.061t} = \frac{16}{3}$	A1	5 or equiv of required form which might involve 5.33 or greater accuracy on RHS; final two marks might be earned in part iii
		<u>Or:</u> State or imply $M_2 = 75 \times r^{0.1t}$	B1	for positive value r
		Obtain $75 \times 1.6^{0.1t}$	B1	
		Attempt to find M_2 in terms of e	M1	
		Equate masses and attempt rearrangement	M1	
		Obtain $e^{0.061t} = \frac{16}{3}$	A1	5 or equiv of required form which might involve 5.33 or greater accuracy on RHS; final two marks might be earned in part iii

	(iii)	Attempt solution involving logarithm		
		of any equation of form $e^{mt} = c_1$	M1	whether the conclusion of part ii or not
		Obtain 27.4	A1	2 or greater accuracy 27.4422...; correct answer only earns both marks

10

9 (i) Use at least one identity correctly Attempt use of relevant identities in single rational expression	B1	angle-sum or angle-difference identity
	M1	not earned if identities used in expression where step equiv to $\frac{A+B+C}{D+E+F} = \frac{A}{D} + \frac{B}{E} + \frac{C}{F}$ or similar has been carried out; condone (for M1A1) if signs of identities apparently switched (so that, for example, denominator appears as $\cos \theta \cos \alpha - \sin \theta \sin \alpha +$ $3 \cos \theta + \cos \theta \cos \alpha + \sin \theta \sin \alpha$)
Obtain $\frac{2 \sin \theta \cos \alpha + 3 \sin \theta}{2 \cos \theta \cos \alpha + 3 \cos \theta}$	A1	or equiv but with the other two terms from each of num'r and den'r absent
Attempt factorisation of num'r and den'r	M1	
Obtain $\frac{\sin \theta}{\cos \theta}$ and hence $\tan \theta$	A1	5 AG; necessary detail needed

(ii) State or imply form $k \tan 150^\circ$	M1	obtained without any wrong method seen
State or imply $\frac{4}{3} \tan 150^\circ$	A1	or equiv such as $\frac{12 \sin 150^\circ}{9 \cos 150^\circ}$
Obtain $-\frac{4}{9}\sqrt{3}$	A1	3 or exact equiv (such as $-\frac{4}{3\sqrt{3}}$); correct answer only earns 3/3

(iii) State or imply $\tan 6\theta = k$	B1	
State $\frac{1}{6} \tan^{-1} k$	B1	
Attempt second value of θ	M1	using $6\theta = \tan^{-1} k + (\text{multiple of } 180)$
Obtain $\frac{1}{6} \tan^{-1} k + 30^\circ$	A1	4 and no other value
12		

OCR (Oxford Cambridge and RSA Examinations)
1 Hills Road
Cambridge
CB1 2EU

OCR Customer Contact Centre

14 – 19 Qualifications (General)

Telephone: 01223 553998

Facsimile: 01223 552627

Email: general.qualifications@ocr.org.uk

www.ocr.org.uk

For staff training purposes and as part of our quality assurance programme your call may be recorded or monitored

Oxford Cambridge and RSA Examinations
is a Company Limited by Guarantee
Registered in England
Registered Office; 1 Hills Road, Cambridge, CB1 2EU
Registered Company Number: 3484466
OCR is an exempt Charity

OCR (Oxford Cambridge and RSA Examinations)
Head office
Telephone: 01223 552552
Facsimile: 01223 552553

