

Mark Scheme (Results)

Summer 2014

Pearson Edexcel GCE in Core Mathematics 3
(6665_01)

Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at www.edexcel.com or www.btec.co.uk. Alternatively, you can get in touch with us using the details on our contact us page at www.edexcel.com/contactus.

Pearson: helping people progress, everywhere

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

Summer 2014

Publications Code UA038461

All the material in this publication is copyright

© Pearson Education Ltd 2014

General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

PEARSON EDEXCEL GCE MATHEMATICS

General Instructions for Marking

1. The total number of marks for the paper is 75
2. The Edexcel Mathematics mark schemes use the following types of marks:
 - **M** marks: Method marks are awarded for 'knowing a method and attempting to apply it', unless otherwise indicated.
 - **A** marks: Accuracy marks can only be awarded if the relevant method (M) marks have been earned.
 - **B** marks are unconditional accuracy marks (independent of M marks)
 - Marks should not be subdivided.

3. Abbreviations

These are some of the traditional marking abbreviations that will appear in the mark schemes.

- bod – benefit of doubt
 - ft – follow through
 - the symbol \checkmark will be used for correct ft
 - cao – correct answer only
 - cso - correct solution only. There must be no errors in this part of the question to obtain this mark
 - isw – ignore subsequent working
 - awrt – answers which round to
 - SC: special case
 - oe – or equivalent (and appropriate)
 - d... or dep – dependent
 - indep – independent
 - dp decimal places
 - sf significant figures
 - * The answer is printed on the paper or ag- answer given
 - \square or d... The second mark is dependent on gaining the first mark
4. All A marks are 'correct answer only' (cao.), unless shown, for example, as A1 ft to indicate that previous wrong working is to be followed through. After a misread however, the subsequent A marks affected are treated as A ft, but manifestly absurd answers should never be awarded A marks.

5. For misreading which does not alter the character of a question or materially simplify it, deduct two from any A or B marks gained, in that part of the question affected.
6. If a candidate makes more than one attempt at any question:
 - If all but one attempt is crossed out, mark the attempt which is NOT crossed out.
 - If either all attempts are crossed out or none are crossed out, mark all the attempts and score the highest single attempt.
7. Ignore wrong working or incorrect statements following a correct answer.

General Principles for Core Mathematics Marking

(But note that specific mark schemes may sometimes override these general principles).

Method mark for solving 3 term quadratic:

1. Factorisation

$(x^2 + bx + c) = (x + p)(x + q)$, where $|pq| = |c|$, leading to $x = \dots$

$(ax^2 + bx + c) = (mx + p)(nx + q)$, where $|pq| = |c|$ and $|mn| = |a|$, leading to $x = \dots$

2. Formula

Attempt to use the correct formula (with values for a, b and c).

3. Completing the square

Solving $x^2 + bx + c = 0$: $\left(x \pm \frac{b}{2}\right)^2 \pm q \pm c = 0$, $q \neq 0$, leading to $x = \dots$

Method marks for differentiation and integration:

1. Differentiation

Power of at least one term decreased by 1. ($x^n \rightarrow x^{n-1}$)

2. Integration

Power of at least one term increased by 1. ($x^n \rightarrow x^{n+1}$)

Use of a formula

Where a method involves using a formula that has been learnt, the advice given in recent examiners' reports is that the formula should be quoted first.

Normal marking procedure is as follows:

Method mark for quoting a correct formula and attempting to use it, even if there are small errors in the substitution of values.

Where the formula is not quoted, the method mark can be gained by implication from correct working with values, but may be lost if there is any mistake in the working.

Exact answers

Examiners' reports have emphasised that where, for example, an exact answer is asked for, or working with surds is clearly required, marks will normally be lost if the candidate resorts to using rounded decimals.

Question Number	Scheme	Marks
<p>1.(a)</p>	$f(x) = \frac{4x+1}{x-2}, \quad x > 2$ <p>Applies $\frac{vu' - uv'}{v^2}$ to get $\frac{(x-2) \times 4 - (4x+1) \times 1}{(x-2)^2}$</p> $= \frac{-9}{(x-2)^2}$	<p>M1A1</p> <p>A1*</p> <p>(3)</p>
<p>(b)</p>	$\frac{-9}{(x-2)^2} = -1 \Rightarrow x = ..$ <p>(5,7)</p>	<p>M1</p> <p>A1,A1</p> <p>(3)</p> <p>6 marks</p>
<p>Alt 1.(a)</p>	$f(x) = \frac{4x+1}{x-2} = 4 + \frac{9}{x-2}$ <p>Applies chain rule to get $f'(x) = A(x-2)^{-2}$</p> $= -9(x-2)^{-2} = \frac{-9}{(x-2)^2}$	<p>M1</p> <p>A1, A1*</p> <p>(3)</p>

(a)

M1 Applies the quotient rule to $f(x) = \frac{4x+1}{x-2}$ with $u = 4x+1$ and $v = x-2$. If the rule is quoted it must be

correct. It may be implied by their $u = 4x+1, v = x-2, u' = \dots, v' = \dots$ followed by $\frac{vu' - uv'}{v^2}$.

If neither quoted nor implied only accept expressions of the form $\frac{(x-2) \times A - (4x+1) \times B}{(x-2)^2}$ $A, B > 0$

allowing for a sign slip inside the brackets.

Condone missing brackets for the method mark but not the final answer mark.

Alternatively they could apply the product rule with $u = 4x+1$ and $v = (x-2)^{-1}$. If the rule is quoted

it must be correct. It may be implied by their $u = 4x+1, v = (x-2)^{-1}, u' = \dots, v' = \dots$ followed by $vu' + uv'$.

If it is neither quoted nor implied only accept expressions of the form/ or equivalent to the form

$$(x-2)^{-1} \times C + (4x+1) \times D(x-2)^{-2}$$

A third alternative is to use the Chain rule. For this to score there must have been some attempt to

divide first to achieve $f(x) = \frac{4x+1}{x-2} = \dots + \frac{\dots}{x-2}$ before applying the chain rule to get

$$f'(x) = A(x-2)^{-2}$$

A1 A correct and unsimplified form of the answer.

Accept $\frac{(x-2) \times 4 - (4x+1) \times 1}{(x-2)^2}$ from the quotient rule

Accept $\frac{4x-8-4x-1}{(x-2)^2}$ from the quotient rule even if the brackets were missing in line 1

Accept $(x-2)^{-1} \times 4 + (4x+1) \times -1(x-2)^{-2}$ or equivalent from the product rule

Accept $9 \times -1(x-2)^{-2}$ from the chain rule

A1* Proceeds to achieve the given answer $= \frac{-9}{(x-2)^2}$. Accept $-9(x-2)^{-2}$

All aspects must be correct including the bracketing.

If they differentiated using the product rule the intermediate lines must be seen.

$$\text{Eg. } (x-2)^{-1} \times 4 + (4x+1) \times -1(x-2)^{-2} = \frac{4}{(x-2)} - \frac{4x+1}{(x-2)^2} = \frac{4(x-2) - (4x+1)}{(x-2)^2} = \frac{-9}{(x-2)^2}$$

(b)

M1 Sets $\frac{-9}{(x-2)^2} = -1$ and proceeds to $x = \dots$

The minimum expectation is that they multiply by $(x-2)^2$ and then either, divide by -1 before square rooting or multiply out before solving a 3TQ equation.

A correct answer of $x = 5$ would also score this mark following $\frac{-9}{(x-2)^2} = -1$ as long as no incorrect

work is seen.

A1 $x = 5$

A1 (5, 7) or $x = 5, y = 7$. Ignore any reference to $x = -1$ (and $y = 1$). Do not accept 21/3 for 7

If there is an extra solution, $x > 2$, then withhold this final mark.

Question Number	Scheme	Marks
<p>2.(a)</p> <p>(b)</p>	$2 \ln(2x+1) - 10 = 0 \Rightarrow \ln(2x+1) = 5 \Rightarrow 2x+1 = e^5 \Rightarrow x = ..$ $\Rightarrow x = \frac{e^5 - 1}{2}$ $3^x e^{4x} = e^7 \Rightarrow \ln(3^x e^{4x}) = \ln e^7$ $\ln 3^x + \ln e^{4x} = \ln e^7 \Rightarrow x \ln 3 + 4x \ln e = 7 \ln e$ $x(\ln 3 + 4) = 7 \Rightarrow x = ...$ $x = \frac{7}{(\ln 3 + 4)}$ <p style="text-align: right;">oe</p>	<p>M1</p> <p>A1</p> <p>(2)</p> <p>M1,M1</p> <p>dM1</p> <p>A1</p> <p>(4)</p> <p>6 marks</p>
<p>Alt 1</p> <p>2(b)</p>	$3^x e^{4x} = e^7 \Rightarrow 3^x = \frac{e^7}{e^{4x}}$ $3^x = e^{7-4x} \Rightarrow x \ln 3 = (7-4x) \ln e$ $x(\ln 3 + 4) = 7 \Rightarrow x = ...$ $x = \frac{7}{(\ln 3 + 4)}$	<p>M1,M1</p> <p>dM1</p> <p>A1</p> <p>(4)</p>
<p>Alt 2</p> <p>2(b)</p> <p>Using logs</p>	$3^x e^{4x} = e^7 \Rightarrow \log(3^x e^{4x}) = \log e^7$ $\log 3^x + \log e^{4x} = \log e^7 \Rightarrow x \log 3 + 4x \log e = 7 \log e$ $x(\log 3 + 4 \log e) = 7 \log e \Rightarrow x = ...$ $x = \frac{7 \log e}{(\log 3 + 4 \log e)}$	<p>M1, M1</p> <p>dM1</p> <p>A1</p> <p>(4)</p>
<p>Alt 3</p> <p>2(b)</p> <p>Using \log_3</p>	$3^x e^{4x} = e^7 \Rightarrow 3^x = \frac{e^7}{e^{4x}}$ $3^x = e^{7-4x} \Rightarrow x = (7-4x) \log_3 e$ $x(1+4 \log_3 e) = 7 \log_3 e \Rightarrow x = ...$ $x = \frac{7 \log_3 e}{(1+4 \log_3 e)}$	<p>M1,M1</p> <p>dM1</p> <p>A1</p> <p>(4)</p>
<p>Alt 4</p> <p>2(b)</p> <p>Using $3^x = e^{x \ln 3}$</p>	$3^x e^{4x} = e^7 \Rightarrow e^{x \ln 3} e^{4x} = e^7$ $\Rightarrow e^{x \ln 3 + 4x} = e^7, \Rightarrow x \ln 3 + 4x = 7$ $x(\ln 3 + 4) = 7 \Rightarrow x = ...$ $x = \frac{7}{(\ln 3 + 4)}$	<p>M1,M1</p> <p>dM1 A1</p> <p>(4)</p>

(a)

M1 Proceeds from $2\ln(2x+1) - 10 = 0$ to $\ln(2x+1) = 5$ before taking exp's to achieve x in terms of e^5
Accept for M1 $2\ln(2x+1) - 10 = 0 \Rightarrow \ln(2x+1) = 5 \Rightarrow x = f(e^5)$

Alternatively they could use the power law before taking exp's to achieve x in terms of $\sqrt{e^{10}}$
 $2\ln(2x+1) = 10 \Rightarrow \ln(2x+1)^2 = 10 \Rightarrow (2x+1)^2 = e^{10} \Rightarrow x = g(\sqrt{e^{10}})$

A1 cso. Accept $x = \frac{e^5 - 1}{2}$ or other exact simplified alternatives such as $x = \frac{e^5}{2} - \frac{1}{2}$. Remember to isw.

The decimal answer of 73.7 will score M1A0 unless the exact answer has also been given.

The answer $\frac{\sqrt{e^{10}} - 1}{2}$ does not score this mark unless simplified. $x = \frac{\pm e^5 - 1}{2}$ is M1A0

(b)

M1 Takes ln's or logs of both sides and applies the addition law.

$\ln(3^x e^{4x}) = \ln 3^x + \ln e^{4x}$ or $\ln(3^x e^{4x}) = \ln 3^x + 4x$ is evidence for the addition law

If the e^{4x} was 'moved' over to the right hand side score for either e^{7-4x} or the subtraction law.

$\ln \frac{e^7}{e^{4x}} = \ln e^7 - \ln e^{4x}$ or $3^x e^{4x} = e^7 \Rightarrow 3^x = \frac{e^7}{e^{4x}} \Rightarrow 3^x = e^{7-4x}$ is evidence of the subtraction law

M1 Uses the power law of logs (seen at least once in a term with x as the index Eg $3^x, e^{4x}$ or e^{7-4x}).

$\ln 3^x + \ln e^{4x} = \ln e^7 \Rightarrow x \ln 3 + 4x \ln e = 7 \ln e$ is an example after the addition law

$3^x = e^{7-4x} \Rightarrow x \log 3 = (7 - 4x) \log e$ is an example after the subtraction law.

It is possible to score M0M1 by applying the power law after an incorrect addition/subtraction law

For example $3^x e^{4x} = e^7 \Rightarrow \ln(3^x) \times \ln(e^{4x}) = \ln e^7 \Rightarrow x \ln 3 \times 4x \ln e = 7 \ln e$

dM1 This is dependent upon **both** previous M's. Collects/factorises out term in x and proceeds to $x =$.
Condone sign slips for this mark. An unsimplified answer can score this mark.

A1 If the candidate has taken ln's then they must use $\ln e = 1$ and achieve $x = \frac{7}{(\ln 3 + 4)}$ or equivalent.

If the candidate has taken log's they must be writing log as oppose to ln and achieve

$x = \frac{7 \log e}{(\log 3 + 4 \log e)}$ or other exact equivalents such as $x = \frac{7 \log e}{\log 3e^4}$.

Question Number	Scheme	Marks
3.(a)	$x = 8 \frac{\pi}{8} \tan \left(2 \times \frac{\pi}{8} \right) = \pi$	B1* (1)
(b)	$\frac{dx}{dy} = 8 \tan 2y + 16y \sec^2(2y)$ $\text{At } P \frac{dx}{dy} = 8 \tan 2 \frac{\pi}{8} + 16 \frac{\pi}{8} \sec^2 \left(2 \times \frac{\pi}{8} \right) = \{8 + 4\pi\}$ $\frac{y - \frac{\pi}{8}}{x - \pi} = \frac{1}{8 + 4\pi}, \text{ accept } y - \frac{\pi}{8} = 0.049(x - \pi)$ $\Rightarrow (8 + 4\pi)y = x + \frac{\pi^2}{2}$	M1A1A1 M1 M1A1 A1 (7) (8 marks)

(a)

B1* Either sub $y = \frac{\pi}{8}$ into $x = 8y \tan(2y) \Rightarrow x = 8 \times \frac{\pi}{8} \tan \left(2 \times \frac{\pi}{8} \right) = \pi$

Or sub $x = \pi$, $y = \frac{\pi}{8}$ into $x = 8y \tan(2y) \Rightarrow \pi = 8 \times \frac{\pi}{8} \tan \left(2 \times \frac{\pi}{8} \right) = \pi \times 1 = \pi$

This is a proof and therefore an expectation that at least one intermediate line must be seen, including a term in tangent.

Accept as a minimum $y = \frac{\pi}{8} \Rightarrow x = \pi \tan \left(\frac{\pi}{4} \right) = \pi$

Or $\pi = \pi \times \tan \left(\frac{\pi}{4} \right) = \pi$ ✓

This is a given answer however, and as such there can be no errors.

(b)

M1 Applies the product rule to $8y \tan 2y$ achieving $A \tan 2y + B y \sec^2(2y)$

A1 One term correct. Either $8 \tan 2y$ or $+16y \sec^2(2y)$. There is no requirement for $\frac{dx}{dy} =$

A1 Both lhs and rhs correct. $\frac{dx}{dy} = 8 \tan 2y + 16y \sec^2(2y)$

It is an intermediate line and the expression does not need to be simplified.

Accept $\frac{dx}{dy} = \tan 2y \times 8 + 8y \times 2 \sec^2(2y)$ or $\frac{dy}{dx} = \frac{1}{\tan 2y \times 8 + 8y \times 2 \sec^2(2y)}$ or using implicit

differentiation $1 = \tan 2y \times 8 \frac{dy}{dx} + 8y \times 2 \sec^2(2y) \frac{dy}{dx}$

M1 For fully substituting $y = \frac{\pi}{8}$ into their $\frac{dx}{dy}$ or $\frac{dy}{dx}$ to find a 'numerical' value

Accept $\frac{dx}{dy} =$ awrt 20.6 or $\frac{dy}{dx} =$ awrt 0.05 as evidence

M1 For a correct attempt at an equation of the tangent at the point $\left(\pi, \frac{\pi}{8}\right)$.

The gradient must be an inverted numerical value of their $\frac{dx}{dy}$

$$\text{Look for } \frac{y - \frac{\pi}{8}}{x - \pi} = \frac{1}{\text{numerical } \frac{dx}{dy}},$$

Watch for negative reciprocals which is M0

If the form $y = mx + c$ is used it must be a full method to find a 'numerical' value to c .

A1 A correct equation of the tangent.

Accept $\frac{y - \frac{\pi}{8}}{x - \pi} = \frac{1}{8 + 4\pi}$ or if $y = mx + c$ is used accept $m = \frac{1}{8 + 4\pi}$ and $c = \frac{\pi}{8} - \frac{\pi}{8 + 4\pi}$

Watch for answers like this which are correct $x - \pi = (8 + 4\pi) \left(y - \frac{\pi}{8}\right)$



Accept the decimal answers awrt 2sf $y = 0.049x + 0.24$, awrt 2sf $21y = x + 4.9$, $\frac{y - 0.39}{x - 3.1} = 0.049$

Accept a mixture of decimals and π 's for example $20.6 \left(y - \frac{\pi}{8}\right) = x - \pi$

A1 Correct answer and solution only. $(8 + 4\pi)y = x + \frac{\pi^2}{2}$

Accept exact alternatives such as $4(2 + \pi)y = x + 0.5\pi^2$ and because the question does not ask for a and b to be simplified in the form $ay = x + b$, accept versions like

$$(8 + 4\pi)y = x + \frac{\pi}{8}(8 + 4\pi) - \pi \text{ and } (8 + 4\pi)y = x + (8 + 4\pi) \left(\frac{\pi}{8} - \frac{\pi}{8 + 4\pi}\right)$$

Question Number	Scheme	Marks
4.(a)		<p>'W' Shape B1 (0, 11) and (6, 1) B1</p> <p>(2)</p>
(b)		<p>'V' shape B1 (-6,1) B1 (0,25) B1</p> <p>(3)</p>
(c)	<p>One of $a = 2$ or $b = 6$</p> <p>$a = 2$ and $b = 6$</p>	<p>B1 B1</p> <p>(2)</p> <p>(7 marks)</p>

(a)

B1 A W shape in any position. The arms of the W do not need to be symmetrical but the two bottom points must appear to be at the same height. Do not accept rounded W's.

A correct sketch of $y = f(|x|)$ would score this mark.

B1 A W shape in quadrants 1 and 2 sitting on the x axis with $P' = (0,11)$ **and** $Q' = (6,1)$. It is not necessary to see them labelled. Accept 11 being marked on the y axis for P' . Condone $P' = (11,0)$ marked on the correct axis, but $Q' = (1,6)$ is B0


(b)

B1 Score for a V shape in any position on the grid. The arms of the V do not need to be symmetrical. Do not accept rounded or upside down V's for this mark.

B1 $Q' = (-6, 1)$. It does not need to be labelled but it must correspond to the minimum point on the curve and be in the correct quadrant.

B1 $P' = (0, 25)$. It does not need to be labelled but it must correspond to the y intercept and the line must cross the axis. Accept 25 marked on the correct axis. Condone $P' = (25,0)$ marked on the positive y axis.

Special case: A candidate who mistakenly sketches $y = -2f(x) + 3$ or $y = -2f(-x) + 3$ will arrive at one of the following. They can be awarded SC B1B0B0



(c)

B1 Either states $a = 2$ **or** $b = 6$.

This can be implied (if there are no stated answers given) by the candidate writing that $y = \dots|x - 6| - 1$ or $y = 2|x - \dots| - 1$. If they are both stated and written, the stated answer takes precedence.

B1 States both $a = 2$ **and** $b = 6$

This can be implied by the candidate stating that $y = 2|x - 6| - 1$

If they are both stated and written, the stated answer takes precedence.

Question Number	Scheme	Marks
5.(a)	$x^2 + x - 6 = (x+3)(x-2)$ $\frac{x}{x+3} + \frac{3(2x+1)}{(x+3)(x-2)} = \frac{x(x-2) + 3(2x+1)}{(x+3)(x-2)}$ $= \frac{x^2 + 4x + 3}{(x+3)(x-2)}$ $= \frac{\cancel{(x+3)}(x+1)}{\cancel{(x+3)}(x-2)}$ $= \frac{(x+1)}{(x-2)} \quad \text{cso}$	B1 M1 A1 A1* (4)
(b)	One end either $(y) > 1, (y) \geq 1$ or $(y) < 4, (y) \leq 4$ $1 < y < 4$	B1 B1 (2)
(c)	Attempt to set Either $g(x) = x$ or $g(x) = g^{-1}(x)$ or $g^{-1}(x) = x$ or $g^2(x) = x$ $\frac{(x+1)}{(x-2)} = x \quad \frac{x+1}{x-2} = \frac{2x+1}{x-1} \quad \frac{2x+1}{x-1} = x \quad \frac{\frac{x+1}{x-2} + 1}{\frac{x+1}{x-2} - 2} = x$ $x^2 - 3x - 1 = 0 \Rightarrow x = \dots$ $a = \frac{3 + \sqrt{13}}{2} \text{ oe } (1.5 + \sqrt{3.25}) \quad \text{cso}$	M1 A1, dM1 A1 (4) (10 marks)

(a)

B1 $x^2 + x - 6 = (x+3)(x-2)$ This can occur anywhere in the solution.

M1 For combining the two fractions with a common denominator. The denominator must be correct for their fractions and at least one numerator must have been adapted. Accept as separate fractions. Condone missing brackets.

$$\text{Accept } \frac{x}{x+3} + \frac{3(2x+1)}{x^2+x-6} = \frac{x(x^2+x-6) + 3(2x+1)(x+3)}{(x+3)(x^2+x-6)}$$

$$\text{Condone } \frac{x}{x+3} + \frac{3(2x+1)}{(x+3)(x-2)} = \frac{x \times x - 2}{(x+3)(x-2)} + \frac{3(2x+1)}{(x+3)(x-2)}$$

A1 A correct intermediate form of $\frac{\text{simplified quadratic}}{\text{simplified quadratic}}$

$$\text{Accept } \frac{x^2+4x+3}{(x+3)(x-2)}, \frac{x^2+4x+3}{x^2+x-6}, \text{ OR } \frac{x^3+7x^2+15x+9}{(x+3)(x^2+x-6)} \rightarrow \frac{(x+1)(x+3)(x+3)}{(x+3)(x^2+x-6)}$$

As in question one they can score this mark having 'invisible' brackets on line 1.

A1* Further factorises and cancels (which may be implied) to complete the proof to reach the given

answer = $\frac{(x+1)}{(x-2)}$. All aspects including bracketing must be correct. If a cubic is formed then it needs to be correct.

(b)

B1 States either end of the range. Accept either $y < 4, y \leq 4$ or $y > 1, y \geq 1$ with or without the y's.

B1 Correct range. Accept $1 < y < 4, 1 < g < 4, y > 1$ and $y < 4, (1, 4), 1 < \text{Range} < 4$, even $1 < f < 4$, Do not accept $1 < x < 4, 1 < y \leq 4, [1, 4)$ etc.

Special case, allow B1B0 for $1 < x < 4$

(c)

M1 Attempting to set $g(x) = x, g^{-1}(x) = x$ or $g(x) = g^{-1}(x)$ or $g^2(x) = x$.

If $g^{-1}(x)$ has been used then a full attempt must have been made to make x the subject of the formula. A full attempt would involve cross multiplying, collecting terms, factorising and ending with division.

As a result, it must be in the form $g^{-1}(x) = \frac{\pm 2x \pm 1}{\pm x \pm 1}$

$$\text{Accept as evidence } \frac{(x+1)}{(x-2)} = x \text{ OR } \frac{x+1}{x-2} = \frac{\pm 2x \pm 1}{\pm x \pm 1} \text{ OR } \frac{\pm 2x \pm 1}{\pm x \pm 1} = x \text{ OR } \frac{\frac{x+1}{x-2} + 1}{\frac{x+1}{x-2} - 2} = x$$

A1 $x^2 - 3x - 1 = 0$ or exact equivalent. The $=0$ may be implied by subsequent work.

dM1 For solving a 3TQ=0. It is dependent upon the first M being scored. Do not accept a method using factors unless it clearly factorises. Allow the answer written down awrt 3.30 (from a graphical calculator).

A1 a or $x = \frac{3 + \sqrt{13}}{2}$. Ignore any reference to $\frac{3 - \sqrt{13}}{2}$

Withhold this mark if additional values are given for $x, x > 3$

Question Number	Scheme	Marks
6.(a)	$y_{2,1} = -0.224 \quad , \quad y_{2,2} = (+)0.546$ <p>Change of sign $\Rightarrow Q$ lies between</p>	M1 A1 (2)
(b)	<p>At R $\frac{dy}{dx} = -2x \sin\left(\frac{1}{2}x^2\right) + 3x^2 - 3$</p> $-2x \sin\left(\frac{1}{2}x^2\right) + 3x^2 - 3 = 0 \Rightarrow x = \sqrt{1 + \frac{2}{3}x \sin\left(\frac{1}{2}x^2\right)}$	M1A1 cs0 M1A1* (4)
(c)	$x_1 = \sqrt{1 + \frac{2}{3} \times 1.3 \sin\left(\frac{1}{2} \times 1.3^2\right)}$ <p>$x_1 = \text{awrt } 1.284 \quad x_2 = \text{awrt } 1.276$</p>	M1 A1 (2) (8 marks)

(a)

M1 Sub both $x = 2.1$ and $x = 2.2$ into y and achieve at least one correct to 1 sig fig
In radians $y_{2.1} = \text{awrt } -0.2$ $y_{2.2} = \text{awrt/truncating to } 0.5$

In degrees $y_{2.1} = \text{awrt } 3$ $y_{2.2} = \text{awrt } 4$

A1 Both values correct to 1 sf with a reason and a minimal conclusion.

$y_{2.1} = \text{awrt } -0.2$ $y_{2.2} = \text{awrt/truncating to } 0.5$

Accept change of sign, positive and negative, $y_{2.1} \times y_{2.2} = -1$ as reasons and hence root, Q lies between 2.1 and 2.2 , QED as a minimal conclusion.

Accept a smaller interval spanning the root of 2.131528 , say 2.13 and 2.14 , but the A1 can only be scored when the candidate refers back to the question, stating that as root lies between 2.13 and 2.14 it lies between 2.1 and 2.2

(b)

M1 Differentiating to get $\frac{dy}{dx} = \dots \sin\left(\frac{1}{2}x^2\right) + 3x^2 - 3$ where \dots is a constant, or a linear function in x .

A1 $\frac{dy}{dx} = -2x \sin\left(\frac{1}{2}x^2\right) + 3x^2 - 3$

M1 Sets their $\frac{dy}{dx} = 0$ and proceeds to make the x of their $3x^2$ the subject of the formula

Alternatively they could state $\frac{dy}{dx} = 0$ and write a line such as

$2x \sin\left(\frac{1}{2}x^2\right) = 3x^2 - 3$, before making the x of $3x^2$ the subject of the formula

A1* Correct given solution. $x = \sqrt{1 + \frac{2}{3}x \sin\left(\frac{1}{2}x^2\right)}$

Watch for missing x 's in their formula

(c)

M1 Subs $x = 1.3$ into the iterative formula to find at least x_1 .

This can be implied by $x_1 = \text{awrt } 1.3$ (not just 1.3)

or $x_1 = \sqrt{1 + \frac{2}{3} \times 1.3 \sin\left(\frac{1}{2} \times 1.3^2\right)}$ or $x_1 = \text{awrt } 1.006$ (degrees)

A1 Both answers correct (awrt 3 decimal places). The subscripts are not important. Mark as the first and second values seen. $x_1 = \text{awrt } 1.284$ $x_2 = \text{awrt } 1.276$

Question Number	Scheme	Marks
7.(a)	$\begin{aligned} \operatorname{cosec} 2x + \cot 2x &= \frac{1}{\sin 2x} + \frac{\cos 2x}{\sin 2x} \\ &= \frac{1 + \cos 2x}{\sin 2x} \\ &= \frac{1 + 2\cos^2 x - 1}{2\sin x \cos x} \\ &= \frac{2\cos^2 x}{2\sin x \cos x} \\ &= \frac{\cos x}{\sin x} = \cot x \end{aligned}$	M1 M1 M1 A1 A1*
(b)	$\begin{aligned} \operatorname{cosec}(4\theta + 10^\circ) + \cot(4\theta + 10^\circ) &= \sqrt{3} \\ \cot(2\theta \pm \dots) &= \sqrt{3} \\ 2\theta \pm \dots &= 30^\circ \Rightarrow \theta = 12.5^\circ \\ 2\theta \pm \dots &= 180 + PV^\circ \Rightarrow \theta = \dots^\circ \\ \theta &= 102.5^\circ \end{aligned}$	(5) M1 dM1, A1 dM1 A1 (5) (10 marks)

(a)

M1 Writing $\operatorname{cosec} 2x = \frac{1}{\sin 2x}$ **and** $\cot 2x = \frac{\cos 2x}{\sin 2x}$ or $\frac{1}{\tan 2x}$

M1 Writing the lhs as a single fraction $\frac{a+b}{c}$. The denominator must be correct for their terms.

M1 Uses the appropriate double angle formulae/trig identities to produce a fraction in a form containing no addition or subtraction signs. A form $\frac{p \times q}{s \times t}$ or similar

A1 A correct intermediate line. Accept $\frac{2 \cos^2 x}{2 \sin x \cos x}$ or $\frac{2 \sin x \cos x}{2 \sin x \cos x \tan x}$ or similar

This cannot be scored if errors have been made

A1* Completes the proof by cancelling and using either $\frac{\cos x}{\sin x} = \cot x$ or

$$\frac{1}{\tan x} = \cot x$$

The cancelling could be implied by seeing $\frac{2 \cos x \cos x}{2 \sin x \cos x} = \cot x$

The proof cannot rely on expressions like $\cot = \frac{\cos}{\sin}$ (with missing x 's) for the

final A1

(b)

M1 Attempt to use the solution to part (a) with $2x = 4\theta + 10 \Rightarrow$ to write or imply $\cot(2\theta \pm \dots^\circ) = \sqrt{3}$

Watch for attempts which start $\cot \alpha = \sqrt{3}$. The method mark here is not scored until the α has been replaced by $2\theta \pm \dots^\circ$

Accept a solution from $\cot(2x \pm \dots^\circ) = \sqrt{3}$ where θ has been replaced by another variable.

dM1 Proceeds from the previous method and uses $\tan \dots = \frac{1}{\cot \dots}$ and

$$\arctan\left(\frac{1}{\sqrt{3}}\right) = 30^\circ \text{ to solve } 2\theta \pm \dots^\circ = 30^\circ \Rightarrow \theta = \dots$$

A1 $\theta = 12.5^\circ$ or exact equivalent. Condone answers such as $x = 12.5^\circ$

dM1 This mark is for the correct method to find a second solution to θ . It is dependent upon the first M only.

Accept $2\theta \pm \dots = 180 + PV^\circ \Rightarrow \theta = \dots^\circ$

A1 $\theta = 102.5^\circ$ or exact equivalent. Condone answers such as $x = 102.5^\circ$

Ignore any solutions outside the range. This mark is withheld for any extra solutions within the range.

If radians appear they could just lose the answer marks. So for example

$$2\theta \pm \dots = \frac{\pi}{6} (0.524) \Rightarrow \theta = \dots \text{ is M1dM1A0 followed by}$$

$$2\theta \pm \dots = \pi + \frac{\pi}{6} \Rightarrow \theta = \dots \text{ dM1A0}$$

Special case 1: For candidates in (b) who solve $\cot(4\theta \pm \dots^\circ) = \sqrt{3}$ the mark scheme is severe, so we are awarding a special case solution, scoring 00011.

$$\cot(4\theta + \beta^\circ) = \sqrt{3} \Rightarrow 4\theta + \beta = 30^\circ \Rightarrow \theta = \dots \text{ is M0M0A0 where } \beta = 5^\circ \text{ or } 10^\circ$$

$$\Rightarrow 4\theta + \beta = 210^\circ \Rightarrow \theta = \dots \text{ can score M1A1 Special case.}$$

$$\text{If } \beta = 5^\circ, \theta = 51.25 \text{ If } \beta = 10^\circ, \theta = 50$$

Special case 2: Just answers in (b) **with no working** scores 1 1 0 0 0 for 12.5 and 102.5

$$\text{BUT } \cot(2\theta \pm 5^\circ) = \sqrt{3} \Rightarrow \theta = 12.5^\circ, 102.5^\circ \text{ scores all available marks.}$$

Question Number	Scheme	Marks
7.(a)Alt 1	$\operatorname{cosec} 2x + \cot 2x = \frac{1}{\sin 2x} + \frac{1}{\tan 2x}$ $= \frac{1}{2 \sin x \cos x} + \frac{1 - \tan^2 x}{2 \tan x}$	1 ST M1
	$= \frac{\tan x + (1 - \tan^2 x) \sin x \cos x}{2 \sin x \cos x \tan x} \quad \text{or} \quad = \frac{2 \tan x + 2(1 - \tan^2 x) \sin x \cos x}{4 \sin x \cos x \tan x}$ $= \frac{\tan x + \sin x \cos x - \tan^2 x \sin x \cos x}{2 \sin x \cos x \tan x}$ $= \frac{\tan x + \sin x \cos x - \tan x \sin^2 x}{2 \sin x \cos x \tan x}$ $= \frac{\tan x(1 - \sin^2 x) + \sin x \cos x}{2 \sin x \cos x \tan x}$ $= \frac{\tan x \cos^2 x + \sin x \cos x}{2 \sin x \cos x \tan x}$ $= \frac{\sin x \cos x + \sin x \cos x}{2 \sin x \cos x \tan x}$ $= \frac{2 \sin x \cos x}{2 \sin x \cos x \tan x} \quad \text{oe}$ $= \frac{1}{\tan x} = \cot x$	2 nd M1
7.(a)Alt 2	<p>Example of how main scheme could work in a roundabout route</p> $\operatorname{cosec} 2x + \cot 2x = \cot x \Leftrightarrow \frac{1}{\sin 2x} + \frac{1}{\tan 2x} = \frac{1}{\tan x}$ $\Leftrightarrow \tan 2x \tan x + \sin 2x \tan x = \sin 2x \tan 2x$ $\Leftrightarrow \frac{2 \tan x}{1 - \tan^2 x} \times \tan x + 2 \sin x \cos x \times \frac{\sin x}{\cos x} = 2 \sin x \cos x \times \frac{2 \tan x}{1 - \tan^2 x}$	3 rd M1A1 A1* (5)
		1 st M1
		2 nd M1

Question Number	Scheme	Marks
	$\Leftrightarrow \frac{2 \tan^2 x}{1 - \tan^2 x} + 2 \sin^2 x = \frac{4 \sin^2 x}{1 - \tan^2 x}$ $\times(1 - \tan^2 x) \Leftrightarrow 2 \tan^2 x + 2 \sin^2 x(1 - \tan^2 x) = 4 \sin^2 x$ $\Leftrightarrow 2 \tan^2 x - 2 \sin^2 x \tan^2 x = 2 \sin^2 x$ $\Leftrightarrow 2 \tan^2 x(1 - \sin^2 x) = 2 \sin^2 x$ $\div 2 \tan^2 x \Leftrightarrow 1 - \sin^2 x = \cos^2 x$ <p>As this is true, initial statement is true</p>	<p>3rd M1 A1 A1*</p> <p>(5)</p>

Question Number	Scheme	Marks
8.(a)	$P = \frac{800e^0}{1+3e^0} = \frac{800}{1+3} = 200$	M1,A1 (2)
(b)	$250 = \frac{800e^{0.1t}}{1+3e^{0.1t}}$ $250(1+3e^{0.1t}) = 800e^{0.1t} \Rightarrow 50e^{0.1t} = 250, \Rightarrow e^{0.1t} = 5$ $t = \frac{1}{0.1} \ln(5)$ $t = 10 \ln(5)$	M1,A1 M1 A1 (4)
(c)	$P = \frac{800e^{0.1t}}{1+3e^{0.1t}} \Rightarrow \frac{dP}{dt} = \frac{(1+3e^{0.1t}) \times 800 \times 0.1e^{0.1t} - 800e^{0.1t} \times 3 \times 0.1e^{0.1t}}{(1+3e^{0.1t})^2}$ At $t=10$ $\frac{dP}{dt} = \frac{(1+3e) \times 80e - 240e^2}{(1+3e)^2} = \frac{80e}{(1+3e)^2}$	M1,A1 M1,A1 (4)
(d)	$P = \frac{800e^{0.1t}}{1+3e^{0.1t}} = \frac{800}{e^{-0.1t} + 3} \Rightarrow P_{\max} = \frac{800}{3} = 266.67$. Hence P cannot be 270	B1 (1) (11 marks)

(a)

M1 Sub $t = 0$ into P **and** use $e^0 = 1$ in at least one of the two cases. Accept $P = \frac{800}{1+3}$ as evidence

A1 200. Accept this for both marks as long as no incorrect working is seen.

(b)

M1 Sub $P=250$ into $P = \frac{800e^{0.1t}}{1+3e^{0.1t}}$, cross multiply, collect terms in $e^{0.1t}$ **and** proceed

to $Ae^{0.1t} = B$

Condone bracketing issues and slips in arithmetic.

If they divide terms by $e^{0.1t}$ you should expect to see $Ce^{-0.1t} = D$

A1 $e^{0.1t} = 5$ or $e^{-0.1t} = 0.2$

M1 Dependent upon gaining $e^{0.1t} = E$, for taking \ln 's of both sides and proceeding to $t = \dots$

Accept $e^{0.1t} = E \Rightarrow 0.1t = \ln E \Rightarrow t = \dots$. It could be implied by $t = \text{awrt } 16.1$

A1 $t = 10 \ln(5)$

Accept exact equivalents of this as long as a and b are integers. Eg. $t = 5 \ln(25)$ is fine.

(c)

M1 Scored for a full application of the quotient rule and knowing that

$$\frac{d}{dt} e^{0.1t} = ke^{0.1t} \text{ and NOT } kte^{0.1t}$$

If the rule is quoted it must be correct.

It may be implied by their $u = 800e^{0.1t}, v = 1 + 3e^{0.1t}, u' = pe^{0.1t}, v' = qe^{0.1t}$

followed by $\frac{vu' - uv'}{v^2}$.

If it is neither quoted nor implied only accept expressions of the form

$$\frac{(1 + 3e^{0.1t}) \times pe^{0.1t} - 800e^{0.1t} \times qe^{0.1t}}{(1 + 3e^{0.1t})^2}$$

Condone missing brackets.

You may see the chain or product rule applied to

For applying the product rule see question 1 but still insist on $\frac{d}{dt} e^{0.1t} = ke^{0.1t}$

For the chain rule look for

$$P = \frac{800e^{0.1t}}{1 + 3e^{0.1t}} = \frac{800}{e^{-0.1t} + 3} \Rightarrow \frac{dP}{dt} = 800 \times (e^{-0.1t} + 3)^{-2} \times -0.1e^{-0.1t}$$

A1 A correct unsimplified answer to

$$\frac{dP}{dt} = \frac{(1 + 3e^{0.1t}) \times 800 \times 0.1e^{0.1t} - 800e^{0.1t} \times 3 \times 0.1e^{0.1t}}{(1 + 3e^{0.1t})^2}$$

M1 For substituting $t = 10$ into their $\frac{dP}{dt}$, NOT P

Accept numerical answers for this. 2.59 is the numerical value if $\frac{dP}{dt}$ was correct

A1 $\frac{dP}{dt} = \frac{80e}{(1 + 3e)^2}$ or equivalent such as $\frac{dP}{dt} = 80e(1 + 3e)^{-2}, \frac{80e}{1 + 6e + 9e^2}$

Note that candidates who substitute $t = 10$ before differentiation will score 0 marks

(d)

B1 Accept solutions from substituting $P=270$ and showing that you get an unsolvable equation

Eg. $270 = \frac{800e^{0.1t}}{1 + 3e^{0.1t}} \Rightarrow -27 = e^{0.1t} \Rightarrow 0.1t = \ln(-27)$ which has no answers.

Eg. $270 = \frac{800e^{0.1t}}{1 + 3e^{0.1t}} \Rightarrow -27 = e^{0.1t} \Rightarrow e^{0.1t} / e^x$ is never negative

Accept solutions where it implies the max value is 266.6 or 267. For example accept sight of $\frac{800}{3}$, with a comment 'so it cannot reach 270', or a large value of t ($t > 99$) being substituted in to get 266.6 or 267 with a similar statement, or a graph drawn with an asymptote marked at 266.6 or 267

Do not accept exp's cannot be negative or you cannot ln a negative number without numerical evidence.

Look for both a statement and a comment

Question Number	Scheme	Marks
9.(a)	$R = \sqrt{20}$ $\tan \alpha = \frac{4}{2} \Rightarrow \alpha = \text{awrt } 1.107$	B1 M1A1 (3)
(b)(i)	$'4 + 5R^2' = 104$	B1ft
(ii)	$3\theta - '1.107' = \frac{\pi}{2} \Rightarrow \theta = \text{awrt } 0.89$	M1A1 (3)
(c)(i)	4	B1
(ii)	$3\theta - '1.107' = 2\pi \Rightarrow \theta = \text{awrt } 2.46$	M1A1 (3) (9 marks)

(a)

B1 Accept $R = \sqrt{20}$ or $2\sqrt{5}$ or awrt 4.47

Do not accept $R = \pm\sqrt{20}$

This could be scored in parts (b) or (c) as long as you are certain it is R

M1 for sight of $\tan \alpha = \pm\frac{4}{2}$, $\tan \alpha = \pm\frac{2}{4}$. Condone $\sin \alpha = 4$, $\cos \alpha = 2 \Rightarrow \tan \alpha = \frac{4}{2}$

If R is found first only accept $\sin \alpha = \pm\frac{4}{R}$, $\cos \alpha = \pm\frac{2}{R}$

A1 $\alpha =$ awrt 1.107. The degrees equivalent 63.4° is A0.

If a candidate does all the question in degrees they will lose just this mark.

(b)(i)

B1ft Either 104 or if R was incorrect allow for the numerical value of their ' $4 + 5R^2$ '.
Allow a tolerance of 1 dp on decimal R 's.

(b)(ii)

M1 Using $3\theta \pm$ their '1.107' = $\frac{\pi}{2} \Rightarrow \theta = ..$

Accept $3\theta \pm$ their '1.107' = $(2n+1)\frac{\pi}{2} \Rightarrow \theta = ..$ where n is an integer

Allow slips on the lhs with an extra bracket such as

$3(\theta \pm$ their '1.107') = $\frac{\pi}{2} \Rightarrow \theta = ..$

The degree equivalent is acceptable $3\theta -$ their ' 63.4° ' = $90^\circ \Rightarrow \theta =$

Do not allow mixed units in this question

A1 awrt 0.89 radians or 51.1° . Do not allow multiple solutions for this mark.

(c)(i)

B1 4

(c)(ii)

M1 Using $3\theta \pm$ their '1.107' = $2\pi \Rightarrow \theta = ..$

Accept $3\theta \pm$ their '1.107' = $n\pi \Rightarrow \theta = ..$ where n is an integer, including 0

Allow slips on the lhs with an extra bracket such as

$3(\theta \pm$ their '1.107') = $2\pi \Rightarrow \theta = ..$

The degree equivalent is acceptable $3\theta -$ their ' 63.4° ' = $360^\circ \Rightarrow \theta =$ but

Do not allow mixed units in this question

A1 $\theta =$ awrt 2.46 radians or 141.1° Do not allow multiple solutions for this mark.

Pearson Education Limited. Registered company number 872828
with its registered office at Edinburgh Gate, Harlow, Essex CM20 2JE