








3.



**Figure 1**

A particle of mass 2 kg is suspended from a horizontal ceiling by two light inextensible strings,  $PR$  and  $QR$ . The particle hangs at  $R$  in equilibrium, with the strings in a vertical plane. The string  $PR$  is inclined at  $55^\circ$  to the horizontal and the string  $QR$  is inclined at  $35^\circ$  to the horizontal, as shown in Figure 1.

Find

- (i) the tension in the string  $PR$ ,
- (ii) the tension in the string  $QR$ .

**(7)**

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

**Question 3 continued**


[Lined area for writing answer]

**(Total 7 marks)**

**Q3**



4.



**Figure 2**

A lift of mass 200 kg is being lowered into a mineshaft by a vertical cable attached to the top of the lift. A crate of mass 55 kg is on the floor inside the lift, as shown in Figure 2. The lift descends vertically with constant acceleration. There is a constant upwards resistance of magnitude 150 N on the lift. The crate experiences a constant normal reaction of magnitude 473 N from the floor of the lift.

(a) Find the acceleration of the lift.

(3)

(b) Find the magnitude of the force exerted on the lift by the cable.

(4)

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---



**Question 4 continued**

Multiple horizontal lines for writing.



P 4 4 8 3 5 A 0 9 2 8

5.




Figure 3

A beam  $AB$  has length 5 m and mass 25 kg. The beam is suspended in equilibrium in a horizontal position by two vertical ropes. One rope is attached to the beam at  $A$  and the other rope is attached to the point  $C$  on the beam where  $CB = 0.5$  m, as shown in Figure 3. A particle  $P$  of mass 60 kg is attached to the beam at  $B$  and the beam remains in equilibrium in a horizontal position. The beam is modelled as a uniform rod and the ropes are modelled as light strings.

(a) Find

- (i) the tension in the rope attached to the beam at  $A$ ,
- (ii) the tension in the rope attached to the beam at  $C$ .

(6)

Particle  $P$  is removed and replaced by a particle  $Q$  of mass  $M$  kg at  $B$ . Given that the beam remains in equilibrium in a horizontal position,

(b) find

- (i) the greatest possible value of  $M$ ,
- (ii) the greatest possible tension in the rope attached to the beam at  $C$ .

(6)

---

---

---

---

---

---

---

---

---

---











7. A train travels along a straight horizontal track between two stations,  $A$  and  $B$ . The train starts from rest at  $A$  and moves with constant acceleration  $0.5 \text{ m s}^{-2}$  until it reaches a speed of  $V \text{ m s}^{-1}$ , ( $V < 50$ ). The train then travels at this constant speed before it moves with constant deceleration  $0.25 \text{ m s}^{-2}$  until it comes to rest at  $B$ .

(a) Sketch in the space below a speed-time graph for the motion of the train between the two stations  $A$  and  $B$ .

(2)

The total time for the journey from  $A$  to  $B$  is 5 minutes.

(b) Find, in terms of  $V$ , the length of time, in seconds, for which the train is

(i) accelerating,

(ii) decelerating,

(iii) moving with constant speed.

(5)

Given that the distance between the two stations  $A$  and  $B$  is 6.3 km,

(c) find the value of  $V$ .

(6)







