

Possible C3 questions from past papers P1—P3

Source of the original question is given in brackets, e.g. [P2 January 2001 Question 1]; a question which has been edited is indicated with an asterisk, e.g. [P3 January 2003 Question 8*].

1. The function f , defined for $x \in \mathbb{R}, x > 0$, is such that

$$f'(x) = x^2 - 2 + \frac{1}{x^2}.$$

(a) Find the value of $f''(x)$ at $x = 4$. (3)

(b) Given that $f(3) = 0$, find $f(x)$. (4)

(c) Prove that f is an increasing function. (3)

[P1 June 2001 Question 5]

2. The curve C has equation $y = 2e^x + 3x^2 + 2$. The point A with coordinates $(0, 4)$ lies on C . Find the equation of the tangent to C at A . (5)

[P2 June 2001 Question 1]

3. The root of the equation $f(x) = 0$, where

$$f(x) = x + \ln 2x - 4$$

is to be estimated using the iterative formula $x_{n+1} = 4 - \ln 2x_n$, with $x_0 = 2.4$.

(a) Showing your values of x_1, x_2, x_3, \dots , obtain the value, to 3 decimal places, of the root. (4)

(b) By considering the change of sign of $f(x)$ in a suitable interval, justify the accuracy of your answer to part (a). (2)

[P2 June 2001 Question 2]

4. (i) Prove, by counter-example, that the statement

$$\text{“}\sec(A + B) \equiv \sec A + \sec B, \text{ for all } A \text{ and } B\text{”}$$

is false. (2)

(ii) Prove that

$$\tan \theta + \cot \theta \equiv 2 \operatorname{cosec} 2\theta, \quad \theta \neq \frac{n\pi}{2}, \quad n \in \mathbb{Z}. \quad (5)$$

[P2 June 2001 Question 4]

5. The function f is given by

$$f : x \mapsto \frac{x}{x^2 - 1} - \frac{1}{x + 1}, \quad x > 1.$$

(a) Show that $f(x) = \frac{1}{(x-1)(x+1)}$. (3)

(b) Find the range of f . (2)

The function g is given by

$$g : x \mapsto \frac{2}{x}, \quad x > 0.$$

(c) Solve $gf(x) = 70$. (4)

[P2 June 2001 Question 7]

6. (a) Express $2 \cos \theta + 5 \sin \theta$ in the form $R \cos (\theta - \alpha)$, where $R > 0$ and $0 < \alpha < \frac{\pi}{2}$.
Give the values of R and α to 3 significant figures. (3)

(b) Find the maximum and minimum values of $2 \cos \theta + 5 \sin \theta$ and the smallest possible value of θ for which the maximum occurs. (2)

The temperature T °C, of an unheated building is modelled using the equation

$$T = 15 + 2 \cos \left(\frac{\pi t}{12} \right) + 5 \sin \left(\frac{\pi t}{12} \right), \quad 0 \leq t < 24,$$

where t hours is the number of hours after 1200.

(c) Calculate the maximum temperature predicted by this model and the value of t when this maximum occurs. (4)

(d) Calculate, to the nearest half hour, the times when the temperature is predicted to be 12 °C. (6)

[P2 June 2001 Question 9]

7. The function f is defined by

$$f: x \mapsto |2x - a|, \quad x \in \mathbb{R},$$

where a is a positive constant.

(a) Sketch the graph of $y = f(x)$, showing the coordinates of the points where the graph cuts the axes. (2)

(b) On a separate diagram, sketch the graph of $y = f(2x)$, showing the coordinates of the points where the graph cuts the axes. (2)

(c) Given that a solution of the equation $f(x) = \frac{1}{2}x$ is $x = 4$, find the two possible values of a . (4)

[P2 January 2002 Question 3]

8. (a) Prove that

$$\frac{1 - \cos 2\theta}{\sin 2\theta} \equiv \tan \theta, \quad \theta \neq \frac{n\pi}{2}, \quad n \in \mathbb{Z}. \quad (3)$$

(b) Solve, giving exact answers in terms of π ,

$$2(1 - \cos 2\theta) = \tan \theta, \quad 0 < \theta < \pi. \quad (6)$$

[P2 January 2002 Question 6]

9.

Figure 2




Figure 2 shows part of the curve C with equation $y = f(x)$, where

$$f(x) = 0.5e^x - x^2.$$

The curve C cuts the y -axis at A and there is a minimum at the point B .

(a) Find an equation of the tangent to C at A . (4)

The x -coordinate of B is approximately 2.15. A more exact estimate is to be made of this coordinate using iterations $x_{n+1} = \ln g(x_n)$.

(b) Show that a possible form for $g(x)$ is $g(x) = 4x$. (3)

(c) Using $x_{n+1} = \ln 4x_n$, with $x_0 = 2.15$, calculate x_1 , x_2 and x_3 . Give the value of x_3 to 4 decimal places. (2)

[P2 January 2002 Question 7]

10.
$$f(x) = \frac{2}{x-1} - \frac{6}{(x-1)(2x+1)}, \quad x > 1.$$

(a) Prove that $f(x) = \frac{4}{2x+1}$. (4)

(b) Find the range of f . (2)

(c) Find $f^{-1}(x)$. (3)

(d) Find the range of $f^{-1}(x)$. (1)

[P2 January 2002 Question 8]

11. Use the derivatives of $\sin x$ and $\cos x$ to prove that the derivative of $\tan x$ is $\sec^2 x$. (4)

[P3 January 2002 Question 2]

12. Express $\frac{3}{x^2 + 2x} + \frac{x-4}{x^2 - 4}$ as a single fraction in its simplest form. (7)

[P2 June 2002 Question 2]

13.

Figure 1




Figure 1 shows a sketch of the curve with equation $y = f(x)$, where

$$f(x) = 10 + \ln(3x) - \frac{1}{2}e^x, \quad 0.1 \leq x \leq 3.3.$$

Given that $f(k) = 0$,

(a) show, by calculation, that $3.1 < k < 3.2$. (2)

(b) Find $f'(x)$. (3)

The tangent to the graph at $x = 1$ intersects the y -axis at the point P .

(c) (i) Find an equation of this tangent.

(ii) Find the exact y -coordinate of P , giving your answer in the form $a + \ln b$. (5)

[P2 June 2002 Question 6]

14. $f(x) = x^2 - 2x - 3, x \in \mathbb{R}, x \geq 1.$
- (a) Find the range of f . (1)
- (b) Write down the domain and range of f^{-1} . (2)
- (c) Sketch the graph of f^{-1} , indicating clearly the coordinates of any point at which the graph intersects the coordinate axes. (4)
- Given that $g(x) = |x - 4|, x \in \mathbb{R},$
- (d) find an expression for $gf(x)$. (2)
- (e) Solve $gf(x) = 8$. (5)

[P2 June 2002 Question 8]

15. Express $\frac{y+3}{(y+1)(y+2)} - \frac{y+1}{(y+2)(y+3)}$ as a single fraction in its simplest form. (5)

[P2 November 2002 Question 1]

16. (a) Express $1.5 \sin 2x + 2 \cos 2x$ in the form $R \sin (2x + \alpha)$, where $R > 0$ and $0 < \alpha < \frac{1}{2}\pi$, giving your values of R and α to 3 decimal places where appropriate. (4)
- (b) Express $3 \sin x \cos x + 4 \cos^2 x$ in the form $a \cos 2x + b \sin 2x + c$, where a, b and c are constants to be found. (2)
- (c) Hence, using your answer to part (a), deduce the maximum value of $3 \sin x \cos x + 4 \cos^2 x$. (2)

[P2 November 2002 Question 3]

17. The curve C with equation $y = p + qe^x$, where p and q are constants, passes through the point $(0, 2)$. At the point $P(\ln 2, p + 2q)$ on C , the gradient is 5.
- (a) Find the value of p and the value of q . (5)
- The normal to C at P crosses the x -axis at L and the y -axis at M .
- (b) Show that the area of $\triangle OLM$, where O is the origin, is approximately 53.8. (5)

[P2 November 2002 Question 5]

18.

Figure 1




Figure 1 shows a sketch of the curve with equation $y = e^{-x} - 1$.

(a) Copy Fig. 1 and on the same axes sketch the graph of $y = \frac{1}{2} |x - 1|$. Show the coordinates of the points where the graph meets the axes. (2)

The x -coordinate of the point of intersection of the graph is α .

(b) Show that $x = \alpha$ is a root of the equation $x + 2e^{-x} - 3 = 0$. (3)

(c) Show that $-1 < \alpha < 0$. (2)

The iterative formula $x_{n+1} = -\ln[\frac{1}{2}(3 - x_n)]$ is used to solve the equation $x + 2e^{-x} - 3 = 0$.

(d) Starting with $x_0 = -1$, find the values of x_1 and x_2 . (2)

(e) Show that, to 2 decimal places, $\alpha = -0.58$. (2)

[P2 November 2002 Question 6]

19. The function f is defined by $f: x \mapsto \frac{3x-1}{x-3}, x \in \mathbb{R}, x \neq 3$.

(a) Prove that $f^{-1}(x) = f(x)$ for all $x \in \mathbb{R}, x \neq 3$. (3)

(b) Hence find, in terms of k , $ff(k)$, where $x \neq 3$. (2)

Figure 3




Figure 3 shows a sketch of the one-one function g , defined over the domain $-2 \leq x \leq 2$.

(c) Find the value of $fg(-2)$. (3)

(d) Sketch the graph of the inverse function g^{-1} and state its domain. (3)

The function h is defined by $h: x \mapsto 2g(x-1)$.

(e) Sketch the graph of the function h and state its range. (3)

[P2 November 2002 Question 8]

20. Express $\frac{x}{(x+1)(x+3)} + \frac{x+12}{x^2-9}$ as a single fraction in its simplest form. (6)

[P2 January 2003 Question 1]

21. (a) Sketch the graph of $y = |2x + a|$, $a > 0$, showing the coordinates of the points where the graph meets the coordinate axes. (2)

- (b) On the same axes, sketch the graph of $y = \frac{1}{x}$. (1)

- (c) Explain how your graphs show that there is only one solution of the equation

$$x|2x + a| - 1 = 0. \quad (1)$$

- (d) Find, using algebra, the value of x for which $x|2x + 1| - 1 = 0$. (3)

[P2 January 2003 Question 3]

22. The curve with equation $y = \ln 3x$ crosses the x -axis at the point $P(p, 0)$.

- (a) Sketch the graph of $y = \ln 3x$, showing the exact value of p . (2)

The normal to the curve at the point Q , with x -coordinate q , passes through the origin.

- (b) Show that $x = q$ is a solution of the equation $x^2 + \ln 3x = 0$. (4)

- (c) Show that the equation in part (b) can be rearranged in the form $x = \frac{1}{3}e^{-x^2}$. (2)

- (d) Use the iteration formula $x_{n+1} = \frac{1}{3}e^{-x_n^2}$, with $x_0 = \frac{1}{3}$, to find x_1, x_2, x_3 and x_4 . Hence write down, to 3 decimal places, an approximation for q . (3)

[P2 January 2003 Question 6]

23. (a) Express $\sin x + \sqrt{3} \cos x$ in the form $R \sin(x + \alpha)$, where $R > 0$ and $0 < \alpha < 90^\circ$. (4)

- (b) Show that the equation $\sec x + \sqrt{3} \operatorname{cosec} x = 4$ can be written in the form

$$\sin x + \sqrt{3} \cos x = 2 \sin 2x. \quad (3)$$

- (c) Deduce from parts (a) and (b) that $\sec x + \sqrt{3} \operatorname{cosec} x = 4$ can be written in the form

$$\sin 2x - \sin(x + 60^\circ) = 0. \quad (1)$$

[P2 January 2003 Question 7*]

24.

Figure 3




Figure 3 shows a sketch of the curve with equation $y = f(x)$, $x \geq 0$. The curve meets the coordinate axes at the points $(0, c)$ and $(d, 0)$.

In separate diagrams sketch the curve with equation

(a) $y = f^{-1}(x)$, (2)

(b) $y = 3f(2x)$. (3)

Indicate clearly on each sketch the coordinates, in terms of c or d , of any point where the curve meets the coordinate axes.

Given that f is defined by

$$f : x \mapsto 3(2^{-x}) - 1, \quad x \in \mathbb{R}, \quad x \geq 0,$$

(c) state

(i) the value of c ,

(ii) the range of f . (3)

(d) Find the value of d , giving your answer to 3 decimal places. (3)

The function g is defined by

$$g : x \rightarrow \log_2 x, \quad x \in \mathbb{R}, \quad x \geq 1.$$

(e) Find $fg(x)$, giving your answer in its simplest form. (3)

[P2 January 2003 Question 8]

25. (a) Simplify $\frac{x^2 + 4x + 3}{x^2 + x}$. (2)

(b) Find the value of x for which $\log_2(x^2 + 4x + 3) - \log_2(x^2 + x) = 4$. (4)

[P2 June 2003 Question 1]

26. The functions f and g are defined by

$$f: x \mapsto x^2 - 2x + 3, x \in \mathbb{R}, 0 \leq x \leq 4,$$

$$g: x \mapsto \lambda x^2 + 1, \text{ where } \lambda \text{ is a constant, } x \in \mathbb{R}.$$

(a) Find the range of f . (3)

(b) Given that $gf(2) = 16$, find the value of λ . (3)

[P2 June 2003 Question 2]

27.

Figure 1




Figure 1 shows a sketch of the curve with equation $y = f(x)$, $-1 \leq x \leq 3$. The curve touches the x -axis at the origin O , crosses the x -axis at the point $A(2, 0)$ and has a maximum at the point $B(\frac{4}{3}, 1)$.

In separate diagrams, show a sketch of the curve with equation

(a) $y = f(x + 1)$, (3)

(b) $y = |f(x)|$, (3)

(c) $y = f(|x|)$, (4)

marking on each sketch the coordinates of points at which the curve

(i) has a turning point,

(ii) meets the x -axis.

[P2 June 2003 Question 4]

28. (a) Sketch, on the same set of axes, the graphs of

$$y = 2 - e^{-x} \text{ and } y = \sqrt{x}. \quad (3)$$

[It is not necessary to find the coordinates of any points of intersection with the axes.]

Given that $f(x) = e^{-x} + \sqrt{x} - 2$, $x \geq 0$,

- (b) explain how your graphs show that the equation $f(x) = 0$ has only one solution, (1)

- (c) show that the solution of $f(x) = 0$ lies between $x = 3$ and $x = 4$. (2)

The iterative formula $x_{n+1} = (2 - e^{-x_n})^2$ is used to solve the equation $f(x) = 0$.

- (d) Taking $x_0 = 4$, write down the values of x_1 , x_2 , x_3 and x_4 , and hence find an approximation to the solution of $f(x) = 0$, giving your answer to 3 decimal places. (4)

[P2 June 2003 Question 5]

- 28a. (i) Given that $\cos(x + 30)^\circ = 3 \cos(x - 30)^\circ$, prove that $\tan x^\circ = -\frac{\sqrt{3}}{2}$. (5)

- (ii) (a) Prove that $\frac{1 - \cos 2\theta}{\sin 2\theta} \equiv \tan \theta$. (3)

- (b) Verify that $\theta = 180^\circ$ is a solution of the equation $\sin 2\theta = 2 - 2 \cos 2\theta$. (1)

- (c) Using the result in part (a), or otherwise, find the other two solutions, $0 < \theta < 360^\circ$, of the equation using $\sin 2\theta = 2 - 2 \cos 2\theta$. (4)

[P2 June 2003 Question 8]

29. (a) Express as a fraction in its simplest form

$$\frac{2}{x-3} + \frac{13}{x^2 + 4x - 21}. \quad (3)$$

- (b) Hence solve

$$\frac{2}{x-3} + \frac{13}{x^2 + 4x - 21} = 1. \quad (3)$$

[P2 November 2003 Question 1]

30. Prove that

$$\frac{1 - \tan^2 \theta}{1 + \tan^2 \theta} \equiv \cos 2\theta. \quad (4)$$

[P2 November 2003 Question 5*]

31. The functions f and g are defined by

$$f: x \mapsto |x - a| + a, \quad x \in \mathbb{R},$$

$$g: x \mapsto 4x + a, \quad x \in \mathbb{R}.$$

where a is a positive constant.

- (a) On the same diagram, sketch the graphs of f and g , showing clearly the coordinates of any points at which your graphs meet the axes. (5)
- (b) Use algebra to find, in terms of a , the coordinates of the point at which the graphs of f and g intersect. (3)
- (c) Find an expression for $fg(x)$. (2)
- (d) Solve, for x in terms of a , the equation

$$fg(x) = 3a. \quad (3)$$

[P2 November 2003 Question 7]

32. The curve C has equation $y = f(x)$, where

$$f(x) = 3 \ln x + \frac{1}{x}, \quad x > 0.$$

The point P is a stationary point on C .

- (a) Calculate the x -coordinate of P . (4)

- (b) Show that the y -coordinate of P may be expressed in the form $k - k \ln k$, where k is a constant to be found. (2)

The point Q on C has x -coordinate 1.

- (c) Find an equation for the normal to C at Q . (4)

The normal to C at Q meets C again at the point R .

- (d) Show that the x -coordinate of R

(i) satisfies the equation $6 \ln x + x + \frac{2}{x} - 3 = 0$,

- (ii) lies between 0.13 and 0.14. (4)

[P2 November 2003 Question 8]

33. The function f is given by $f: x \mapsto 2 + \frac{3}{x+2}$, $x \in \mathbb{R}$, $x \neq -2$.

- (a) Express $2 + \frac{3}{x+2}$ as a single fraction. (1)

- (b) Find an expression for $f^{-1}(x)$. (3)

- (c) Write down the domain of f^{-1} . (1)

[P2 January 2004 Question 1]

34. The function f is even and has domain \mathbb{R} . For $x \geq 0$, $f(x) = x^2 - 4ax$, where a is a positive constant.

(a) In the space below, sketch the curve with equation $y = f(x)$, showing the coordinates of all the points at which the curve meets the axes. (3)

(b) Find, in terms of a , the value of $f(2a)$ and the value of $f(-2a)$. (2)

Given that $a = 3$,

(c) use algebra to find the values of x for which $f(x) = 45$. (4)

[P2 January 2004 Question 4]

35. Given that $y = \log_a x$, $x > 0$, where a is a positive constant,

(a) (i) express x in terms of a and y , (1)

(ii) deduce that $\ln x = y \ln a$. (1)

(b) Show that $\frac{dy}{dx} = \frac{1}{x \ln a}$. (2)

The curve C has equation $y = \log_{10} x$, $x > 0$. The point A on C has x -coordinate 10. Using the result in part (b),

(c) find an equation for the tangent to C at A . (4)

The tangent to C at A crosses the x -axis at the point B .

(d) Find the exact x -coordinate of B . (2)

[P2 January 2004 Question 5]

36. (i) (a) Express $(12 \cos \theta - 5 \sin \theta)$ in the form $R \cos (\theta + \alpha)$, where $R > 0$ and $0 < \alpha < 90^\circ$. (4)

(b) Hence solve the equation

$$12 \cos \theta - 5 \sin \theta = 4,$$

for $0 < \theta < 90^\circ$, giving your answer to 1 decimal place. (3)

(ii) Solve

$$8 \cot \theta - 3 \tan \theta = 2,$$

for $0 < \theta < 90^\circ$, giving your answer to 1 decimal place. (5)

[P2 January 2004 Question 8]

37. Express as a single fraction in its simplest form

$$\frac{x^2 - 8x + 15}{x^2 - 9} \times \frac{2x^2 + 6x}{(x - 5)^2}.$$

(4)

[P2 June 2004 Question 1]

38. (i) Given that $\sin x = \frac{3}{5}$, use an appropriate double angle formula to find the exact value of $\sec 2x$. (4)

(ii) Prove that

$$\cot 2x + \operatorname{cosec} 2x \equiv \cot x, \quad \left(x \neq \frac{n\pi}{2}, n \in \mathbb{Z} \right).$$

(4)

[P2 June 2004 Question 2]

39.

$$f(x) = x^3 + x^2 - 4x - 1.$$

The equation $f(x) = 0$ has only one positive root, α .

(a) Show that $f(x) = 0$ can be rearranged as

$$x = \sqrt{\left(\frac{4x+1}{x+1}\right)}, x \neq -1. \quad (2)$$

The iterative formula $x_{n+1} = \sqrt{\left(\frac{4x_n+1}{x_n+1}\right)}$ is used to find an approximation to α .

(b) Taking $x_1 = 1$, find, to 2 decimal places, the values of x_2 , x_3 and x_4 . (3)

(c) By choosing values of x in a suitable interval, prove that $\alpha = 1.70$, *correct* to 2 decimal places. (3)

(d) Write down a value of x_1 for which the iteration formula $x_{n+1} = \sqrt{\left(\frac{4x_n+1}{x_n+1}\right)}$ does *not* produce a valid value for x_2 .

Justify your answer.

(2)

[P2 June 2004 Question 5]

40.
$$f(x) = x + \frac{e^x}{5}, \quad x \in \mathbb{R}.$$

(a) Find $f'(x)$. (2)

The curve C , with equation $y = f(x)$, crosses the y -axis at the point A .

(b) Find an equation for the tangent to C at A . (3)

(c) Complete the table, giving the values of $\sqrt{x + \frac{e^x}{5}}$ to 2 decimal places.

x	0	0.5	1	1.5	2
$\sqrt{x + \frac{e^x}{5}}$	0.45	0.91			

(2)

[P2 June 2004 Question 7*]

41. The function f is given by

$$f: x \mapsto \ln(3x - 6), \quad x \in \mathbb{R}, \quad x > 2.$$

(a) Find $f^{-1}(x)$. (3)

(b) Write down the domain of f^{-1} and the range of f^{-1} . (2)

(c) Find, to 3 significant figures, the value of x for which $f(x) = 3$. (2)

The function g is given by

$$g: x \mapsto \ln|3x - 6|, \quad x \in \mathbb{R}, \quad x \neq 2.$$

(d) Sketch the graph of $y = g(x)$. (3)

(e) Find the exact coordinates of all the points at which the graph of $y = g(x)$ meets the coordinate axes. (3)

[P2 June 2004 Question 8]