

**ADVANCED SUBSIDIARY GCE
MATHEMATICS**

4725/01

Further Pure Mathematics 1

FRIDAY 11 JANUARY 2008

Morning

Time: 1 hour 30 minutes

Additional materials: Answer Booklet (8 pages)
List of Formulae (MF1)

INSTRUCTIONS TO CANDIDATES

- Write your name, centre number and candidate number in the spaces provided on the answer booklet.
- Read each question carefully and make sure you know what you have to do before starting your answer.
- Answer **all** the questions.
- Give non-exact numerical answers correct to 3 significant figures unless a different degree of accuracy is specified in the question or is clearly appropriate.
- You are permitted to use a graphical calculator in this paper.

INFORMATION FOR CANDIDATES

- The number of marks is given in brackets [] at the end of each question or part question.
- The total number of marks for this paper is 72.
- **You are reminded of the need for clear presentation in your answers.**

This document consists of **4** printed pages.

- 1** The transformation S is a shear with the y -axis invariant (i.e. a shear parallel to the y -axis). It is given that the image of the point $(1, 1)$ is the point $(1, 0)$.

(i) Draw a diagram showing the image of the unit square under the transformation S . [2]

(ii) Write down the matrix that represents S . [2]

- 2** Given that $\sum_{r=1}^n (ar^2 + b) \equiv n(2n^2 + 3n - 2)$, find the values of the constants a and b . [5]

- 3** The cubic equation $2x^3 - 3x^2 + 24x + 7 = 0$ has roots α , β and γ .

(i) Use the substitution $x = \frac{1}{u}$ to find a cubic equation in u with integer coefficients. [2]

(ii) Hence, or otherwise, find the value of $\frac{1}{\alpha\beta} + \frac{1}{\beta\gamma} + \frac{1}{\gamma\alpha}$. [2]

- 4** The complex number $3 - 4i$ is denoted by z . Giving your answers in the form $x + iy$, and showing clearly how you obtain them, find

(i) $2z + 5z^*$, [2]

(ii) $(z - i)^2$, [3]

(iii) $\frac{3}{z}$. [3]

- 5** The matrices \mathbf{A} , \mathbf{B} and \mathbf{C} are given by $\mathbf{A} = \begin{pmatrix} 3 \\ 1 \\ 2 \end{pmatrix}$, $\mathbf{B} = \begin{pmatrix} 4 \\ 0 \\ 3 \end{pmatrix}$ and $\mathbf{C} = \begin{pmatrix} 2 & 4 & -1 \end{pmatrix}$. Find

(i) $\mathbf{A} - 4\mathbf{B}$, [2]

(ii) \mathbf{BC} , [4]

(iii) \mathbf{CA} . [2]

- 6** The loci C_1 and C_2 are given by

$$|z| = |z - 4i| \quad \text{and} \quad \arg z = \frac{1}{6}\pi$$

respectively.

(i) Sketch, on a single Argand diagram, the loci C_1 and C_2 . [5]

(ii) Hence find, in the form $x + iy$, the complex number represented by the point of intersection of C_1 and C_2 . [3]

7 The matrix \mathbf{A} is given by $\mathbf{A} = \begin{pmatrix} a & 3 \\ -2 & 1 \end{pmatrix}$.

(i) Given that \mathbf{A} is singular, find a . [2]

(ii) Given instead that \mathbf{A} is non-singular, find \mathbf{A}^{-1} and hence solve the simultaneous equations

$$\begin{aligned} ax + 3y &= 1, \\ -2x + y &= -1. \end{aligned} \quad [5]$$

8 The sequence u_1, u_2, u_3, \dots is defined by $u_1 = 1$ and $u_{n+1} = u_n + 2n + 1$.

(i) Show that $u_4 = 16$. [2]

(ii) Hence suggest an expression for u_n . [1]

(iii) Use induction to prove that your answer to part (ii) is correct. [4]

9 (i) Show that $\alpha^3 + \beta^3 = (\alpha + \beta)^3 - 3\alpha\beta(\alpha + \beta)$. [2]

(ii) The quadratic equation $x^2 - 5x + 7 = 0$ has roots α and β . Find a quadratic equation with roots α^3 and β^3 . [6]

10 (i) Show that $\frac{2}{r} - \frac{1}{r+1} - \frac{1}{r+2} = \frac{3r+4}{r(r+1)(r+2)}$. [2]

(ii) Hence find an expression, in terms of n , for

$$\sum_{r=1}^n \frac{3r+4}{r(r+1)(r+2)}. \quad [6]$$

(iii) Hence write down the value of $\sum_{r=1}^{\infty} \frac{3r+4}{r(r+1)(r+2)}$. [1]

(iv) Given that $\sum_{r=N+1}^{\infty} \frac{3r+4}{r(r+1)(r+2)} = \frac{7}{10}$, find the value of N . [4]

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (OCR) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

OCR is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.