

ADVANCED SUBSIDIARY GCE MATHEMATICS

Further Pure Mathematics 1

4725

Candidates answer on the Answer Booklet

OCR Supplied Materials:

- 8 page Answer Booklet
- List of Formulae (MF1)

Other Materials Required: None

Thursday 15 January 2009 Morning

Duration: 1 hour 30 minutes



INSTRUCTIONS TO CANDIDATES

- Write your name clearly in capital letters, your Centre Number and Candidate Number in the spaces provided on the Answer Booklet.
- Use black ink. Pencil may be used for graphs and diagrams only.
- Read each question carefully and make sure that you know what you have to do before starting your answer.
- Answer all the questions.
- Do **not** write in the bar codes.
- Give non-exact numerical answers correct to 3 significant figures unless a different degree of accuracy is specified in the question or is clearly appropriate.
- You are permitted to use a graphical calculator in this paper.

INFORMATION FOR CANDIDATES

- The number of marks is given in brackets [] at the end of each question or part question.
- You are reminded of the need for clear presentation in your answers.
- The total number of marks for this paper is 72.
- This document consists of 4 pages. Any blank pages are indicated.

Express $\frac{2+3i}{5-i}$ in the form x + iy, showing clearly how you obtain your answer. 1 [4]

2 The matrix **A** is given by
$$\mathbf{A} = \begin{pmatrix} 2 & 0 \\ a & 5 \end{pmatrix}$$
. Find
(i) \mathbf{A}^{-1} , [2]

(ii)
$$2\mathbf{A} - \begin{pmatrix} 1 & 2 \\ 0 & 4 \end{pmatrix}$$
. [2]

- Find $\sum_{r=1}^{n} (4r^3 + 6r^2 + 2r)$, expressing your answer in a fully factorised form. 3 [6]
- 4 Given that A and B are 2×2 non-singular matrices and I is the 2×2 identity matrix, simplify

$$\mathbf{B}(\mathbf{A}\mathbf{B})^{-1}\mathbf{A} - \mathbf{I}.$$
 [4]

5 By using the determinant of an appropriate matrix, or otherwise, find the value of k for which the simultaneous equations

$$2x - y + z = 7,3y + z = 4,x + ky + kz = 5,$$

do not have a unique solution for x, y and z.

- (i) The transformation P is represented by the matrix $\begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$. Give a geometrical description of 6 transformation P. [2]
 - (ii) The transformation Q is represented by the matrix $\begin{pmatrix} 0 & -1 \\ -1 & 0 \end{pmatrix}$. Give a geometrical description of transformation Q. [2]
 - (iii) The transformation R is equivalent to transformation P followed by transformation Q. Find the matrix that represents R. [2]
 - (iv) Give a geometrical description of the single transformation that is represented by your answer to part (iii). [3]
- It is given that $u_n = 13^n + 6^{n-1}$, where *n* is a positive integer. 7
 - (i) Show that $u_n + u_{n+1} = 14 \times 13^n + 7 \times 6^{n-1}$. [3]
 - (ii) Prove by induction that u_n is a multiple of 7.

4725 Jan09

[5]

[4]

8 (i) Show that $(\alpha - \beta)^2 \equiv (\alpha + \beta)^2 - 4\alpha\beta$. [2]

The quadratic equation $x^2 - 6kx + k^2 = 0$, where k is a positive constant, has roots α and β , with $\alpha > \beta$.

- (ii) Show that $\alpha \beta = 4\sqrt{2k}$. [4]
- (iii) Hence find a quadratic equation with roots $\alpha + 1$ and $\beta 1$. [4]
- 9 (i) Show that $\frac{1}{2r-3} \frac{1}{2r+1} = \frac{4}{4r^2 4r 3}$. [2]
 - (ii) Hence find an expression, in terms of *n*, for

$$\sum_{r=2}^{n} \frac{4}{4r^2 - 4r - 3}.$$
 [6]

(iii) Show that
$$\sum_{r=2}^{\infty} \frac{4}{4r^2 - 4r - 3} = \frac{4}{3}$$
. [1]

- 10 (i) Use an algebraic method to find the square roots of the complex number $2 + i\sqrt{5}$. Give your answers in the form x + iy, where x and y are exact real numbers. [6]
 - (ii) Hence find, in the form x + iy where x and y are exact real numbers, the roots of the equation

$$z^4 - 4z^2 + 9 = 0.$$
 [4]

- (iii) Show, on an Argand diagram, the roots of the equation in part (ii). [1]
- (iv) Given that α is the root of the equation in part (ii) such that $0 < \arg \alpha < \frac{1}{2}\pi$, sketch on the same Argand diagram the locus given by $|z \alpha| = |z|$. [3]



Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (OCR) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

4

OCR is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.