




Mark Scheme 4753
June 2006

1 $3x-2 =x$ $\Rightarrow 3x-2=x \Rightarrow 2x=2 \Rightarrow x=1$ or $2-3x=x \Rightarrow 2=4x \Rightarrow x=\frac{1}{2}$ or $(3x-2)^2=x^2$ $\Rightarrow 8x^2-12x+4=0 \Rightarrow 2x^2-3x+1=0$ $\Rightarrow (x-1)(2x-1)=0,$ $\Rightarrow x=1, \frac{1}{2}$	B1 M1 A1 M1 A1 A1 [3]	$x=1$ solving correct quadratic
2 let $u=x$, $dv/dx=\sin 2x \Rightarrow v=-\frac{1}{2}\cos 2x$ $\Rightarrow \int_0^{\pi/6} x \sin 2x dx = \left[x \cdot -\frac{1}{2} \cos 2x \right]_0^{\pi/6} + \int_0^{\pi/6} \frac{1}{2} \cos 2x \cdot 1 \cdot dx$ $= \frac{\pi}{6} \cdot -\frac{1}{2} \cos \frac{\pi}{3} - 0 + \left[\frac{1}{4} \sin 2x \right]_0^{\pi/6}$ $= -\frac{\pi}{24} + \frac{\sqrt{3}}{8}$ $= \frac{3\sqrt{3}-\pi}{24} *$	M1 A1 B1ft M1 B1 E1 [6]	parts with $u=x$, $dv/dx=\sin 2x$... + $\left[\frac{1}{4} \sin 2x \right]_0^{\pi/6}$ substituting limits $\cos \pi/3 = \frac{1}{2}$, $\sin \pi/3 = \sqrt{3}/2$ soi www
3 (i) $x-1=\sin y$ $\Rightarrow x=1+\sin y$ $\Rightarrow dx/dy=\cos y$ (ii) When $x=1.5$, $y=\arcsin(0.5)=\pi/6$ $\frac{dy}{dx}=\frac{1}{\cos y}$ $=\frac{1}{\cos \pi/6}$ $=2/\sqrt{3}$	M1 A1 E1 M1 A1 M1 A1 [7]	www condone 30° or 0.52 or better or $\frac{dy}{dx}=\frac{1}{\sqrt{1-(x-1)^2}}$ or equivalent, but must be exact
4(i) $V=\pi h^2-\frac{1}{3}\pi h^3$ $\Rightarrow \frac{dV}{dh}=2\pi h-\pi h^2$ (ii) $\frac{dV}{dt}=0.02$ $\frac{dV}{dt}=\frac{dV}{dh} \cdot \frac{dh}{dt}$ $\Rightarrow \frac{dh}{dt}=\frac{0.02}{dV/dh}=\frac{0.02}{2\pi h-\pi h^2}$ When $h=0.4$, $\Rightarrow \frac{dh}{dt}=\frac{0.02}{0.8\pi-0.16\pi}=0.0099 \text{ m/min}$	M1 A1 B1 M1 M1dep A1cao [6]	expanding brackets (correctly) or product rule oe soi $\frac{dV}{dt}=\frac{dV}{dh} \cdot \frac{dh}{dt}$ oe substituting $h=0.4$ into their $\frac{dV}{dh}$ and $\frac{dV}{dt}=0.02$ 0.01 or better or $1/32\pi$

5(i) $a^2 + b^2 = (2t)^2 + (t^2 - 1)^2$ $= 4t^2 + t^4 - 2t^2 + 1$ $= t^4 + 2t^2 + 1$ $= (t^2 + 1)^2 = c^2$ (ii) $c = \sqrt{(20^2 + 21^2)} = 29$ For example: $2t = 20 \Rightarrow t = 10$ $\Rightarrow t^2 - 1 = 99$ which is not consistent with 21	M1 M1 E1 B1 M1 E1 [6]	substituting for a, b and c in terms of t Expanding brackets correctly www Attempt to find t Any valid argument or E2 'none of 20, 21, 29 differ by two'.
6 (i)  (ii) $\frac{M}{M_0} = e^{-0.000121 \times 5730} = e^{-0.6933...} \approx \frac{1}{2}$ (iii) $\frac{M}{M_0} = e^{-kT} = \frac{1}{2}$ $\Rightarrow \ln \frac{1}{2} = -kT$ $\Rightarrow \ln 2 = kT$ $\Rightarrow T = \frac{\ln 2}{k} *$ (iv) $T = \frac{\ln 2}{2.88 \times 10^{-5}} \approx 24000$ years	B1 B1 M1 E1 M1 M1 E1 B1 [8]	Correct shape Passes through $(0, M_0)$ substituting $k = -0.00121$ and $t = 5730$ into equation (or ln eqn) showing that $M \approx \frac{1}{2} M_0$ substituting $M/M_0 = \frac{1}{2}$ into equation (oe) taking lns correctly 24 000 or better

Section B

7(i) $x = 1$	B1 [1]	
(ii) $\frac{dy}{dx} = \frac{(x-1)2x - (x^2+3).1}{(x-1)^2}$ $= \frac{2x^2 - 2x - x^2 - 3}{(x-1)^2}$ $= \frac{x^2 - 2x - 3}{(x-1)^2}$ $dy/dx = 0$ when $x^2 - 2x - 3 = 0$ $\Rightarrow (x-3)(x+1) = 0$ $\Rightarrow x = 3 \text{ or } -1$ When $x = 3$, $y = (9+3)/2 = 6$ So P is (3, 6)	M1 A1 M1 M1 A1 B1ft [6]	Quotient rule correct expression their numerator = 0 solving quadratic by any valid method $x = 3$ from correct working $y = 6$
(iii) Area = $\int_2^3 \frac{x^2+3}{x-1} dx$ $u = x - 1 \Rightarrow du/dx = 1, du = dx$ When $x = 2$, $u = 1$; when $x = 3$, $u = 2$ $= \int_1^2 \frac{(u+1)^2+3}{u} du$ $= \int_1^2 \frac{u^2+2u+4}{u} du$ $= \int_1^2 (u+2+\frac{4}{u}) du *$ $= \left[\frac{1}{2}u^2 + 2u + 4\ln u \right]_1^2$ $= (2+4+4\ln 2) - (\frac{1}{2}+2+4\ln 1)$ $= 3\frac{1}{2} + 4\ln 2$	M1 B1 B1 E1 B1 M1 A1cao [7]	Correct integral and limits Limits changed, and substituting $dx = du$ substituting $\frac{(u+1)^2+3}{u}$ www [$\frac{1}{2}u^2 + 2u + 4\ln u$] substituting correct limits
(iv) $e^y = \frac{x^2+3}{x-1}$ $\Rightarrow e^y \frac{dy}{dx} = \frac{x^2-2x-3}{(x-1)^2}$ $\Rightarrow \frac{dy}{dx} = e^{-y} \frac{x^2-2x-3}{(x-1)^2}$ When $x = 2$, $e^y = 7 \Rightarrow$ $\Rightarrow dy/dx = \frac{1}{7} \cdot \frac{4-4-3}{1} = -\frac{3}{7}$	M1 A1ft B1 A1cao [4]	$e^y dy/dx = \text{their } f'(x)$ or $xe^y - e^y = x^2 + 3$ $\Rightarrow e^y + xe^y \frac{dy}{dx} - e^y \frac{dy}{dx} = 2x$ $\Rightarrow \frac{dy}{dx} = \frac{2x - e^y}{e^y(x-1)}$ $y = \ln 7$ or 1.95... or $e^y = 7$ or $\frac{dy}{dx} = \frac{4-7}{7(2-1)} = -\frac{3}{7}$ or -0.43 or better

8 (i) (A)  (B) 	B1 B1 M1 A1 [4]	Zeros shown every $\pi/2$. Correct shape, from $-\pi$ to π Translated in x-direction π to the left
(ii) $f(x) = -\frac{1}{5}e^{\frac{1}{5}x} \sin x + e^{\frac{1}{5}x} \cos x$ $f(x) = 0$ when $-\frac{1}{5}e^{\frac{1}{5}x} \sin x + e^{\frac{1}{5}x} \cos x = 0$ $\Rightarrow \frac{1}{5}e^{\frac{1}{5}x} (-\sin x + 5 \cos x) = 0$ $\Rightarrow \sin x = 5 \cos x$ $\Rightarrow \frac{\sin x}{\cos x} = 5$ $\Rightarrow \tan x = 5^*$ $\Rightarrow x = 1.37(34\dots)$ $\Rightarrow y = 0.75$ or $0.74(5\dots)$	B1 B1 M1 E1 B1 B1 [6]	$e^{\frac{1}{5}x} \cos x$ $\dots -\frac{1}{5}e^{\frac{1}{5}x} \sin x$ dividing by $e^{\frac{1}{5}x}$ www 1.4 or better, must be in radians 0.75 or better
(iii) $f(x + \pi) = e^{\frac{1}{5}(x+\pi)} \sin(x + \pi)$ $= e^{\frac{1}{5}x} e^{\frac{1}{5}\pi} \sin(x + \pi)$ $= -e^{\frac{1}{5}x} e^{\frac{1}{5}\pi} \sin x$ $= -e^{\frac{1}{5}\pi} f(x)^*$ $\int_{\pi}^{2\pi} f(x) dx$ let $u = x - \pi$, $du = dx$ $= \int_0^{\pi} f(u + \pi) du$ $= \int_0^{\pi} -e^{\frac{1}{5}\pi} f(u) du$ $= -e^{\frac{1}{5}\pi} \int_0^{\pi} f(u) du^*$ Area enclosed between π and 2π $= (-) e^{\frac{1}{5}\pi} \times \text{area between } 0 \text{ and } \pi.$	M1 A1 A1 E1 B1 B1dep E1 B1 [8]	$e^{\frac{1}{5}(x+\pi)} = e^{\frac{1}{5}x} \cdot e^{\frac{1}{5}\pi}$ $\sin(x + \pi) = -\sin x$ www $\int f(u + \pi) du$ limits changed using above result or repeating work or multiplied by 0.53 or better