

Mark Scheme 4734
June 2006

1	Add two Poisson distributions With mean 17 $P(27) = e^{-17} 17^{27} / 27!$ or $P(\leq 27) - P(\leq 26)$ 0.00634 or 0.0063, 0.0064 from tables	M1 A1 M1 A1	4	Use formula or table M1A1 0.0052 from N(17,17)
2	$H_0: p_1 = p_2 = p_3 = p_4$, (H_1 : They are not all equal) Expected values under $H_0 = 150$ $X^2 = (12^2 + 23^2 + 15^2 + 20^2) / 150$ = 8.653 Critical value with 3 d.f. = 7.815 ($X^2 > 7.815$ so) reject H_0 and accept that proportions are different.	B1 B1 M1 B1 B1✓	A1 6	Indication of equality of proportions At least one correct term Accept art 8.65 or 8.66 ft critical value
3	Assume population of differences has a normal distribution. or sample random $H_0: \mu_B - \mu_A = 0$, $H_1: \mu_B - \mu_A > 0$ $t = (23.43 - 22.84) / \sqrt{(0.548/10)}$ = 2.520 CV = 1.833 2.52 > CV so reject H_0 1.812, 1.734 Accept that there is evidence that mean time has reduced.	B1 B1 M1 B1 M1 A1✓	A1 7	Either assumption. AEF Seen Allow from CV 2.262 (2-tail), ft wrong CV
4	(i) EITHER: $\int_{q_3}^4 \frac{1}{12} x dx = \frac{1}{4}$ or $\int_1^2 \frac{4}{3x^3} dx + \int_2^{q_3} \frac{1}{12} x dx = \frac{3}{4}$ [$x^2/24$] OR [-2/(3x ²) + $x^2/24$] (16 - q_3^2)/24 = 1/4 or $1/3 + q_3^2/24 = 3/4$ $q_3 = \sqrt{10}$ If they find F(x): M1A1, M1A1	M1* A1 dep *M1 A1	4	Either Form equation and attempt to solve Accept to 3 SF
(ii)	$E(X^2) = \int_1^2 \frac{4}{3x} dx + \int_2^4 \frac{x^3}{12} dx$ $E(X) = \int_1^2 \frac{4}{3x^2} dx + \int_2^4 \frac{x^2}{12} dx$ $\left[\frac{4}{3} \ln x \right]_1^2 + \left[\frac{x^4}{48} \right]_2^4$ $\left[\frac{-4}{3x} \right]_1^2 + \left[\frac{x^3}{36} \right]_2^4$ $a = E(X^2)/E(X)$ $a = 2.6659, 2.67$	M1 A1 A1 M1 A1	5	Either correct Or exact value, (3ln2)/5 + 9/4 or equiv.

5	(i)	$(48 \times 72/150)$ or $(48/150)(72/150) \times 150$	M1 A1	2	Multiply and divide relevant values All correct
	(ii)	No, no expected value less than 5	B1	1	
	(iii)	H_0 : Volume and day are independent (H_1 : Volume and day are not independent) Critical value for 4 df=13.28 Test statistic > 13.28, reject H_0 Accept that volume and day are not independent	B1 B1 M1 A1	4	Attributes specified
	(iv)	Choose Friday Highest volume	B1	B1	2 Not reference to E values
6	(i)	(a) No 0.43 belongs to relevant interval (b) Yes 0.43 is outside relevant interval	B1 B1 B1	3	Must be with reason
	(ii)	$H_0: p_R = p_T$, $H_1: p_R \neq p_T$ Estimate of $p = 74/165$ Variance estimate of difference $= \left(\frac{74}{165}\right)\left(\frac{91}{165}\right)\left(\frac{1}{80} + \frac{1}{85}\right)$ $z = (28/80 - 46/85)/\sigma_{\text{est}}$ $= -2.468$ Compare correctly with CV $-2.468 < -2.326$, or $2.468 > 2.326$ Reject H_0 and accept that the proportions differ on the island.	B1 B1 B1 M1 A1 A1 M1 A1	8	Proportions May be implied by later work Standardising Completely correct expression + or - , 2.47 Conclusion in context
7	(i)	$T_1 \sim N(2.2, 0.75^2)$, $T_2 \sim N(1.8, 0.70^2)$ Use $T_2 - \frac{1}{2} T_1$ normal $\mu = 0.7$ $\sigma^2 = 0.7^2 + \frac{1}{4} \times 0.75^2$ (0.630625) $(0 - \mu)/\sigma$ -0.881 Probability 0.189	M1 A1 A1 M1 A1	A1	Or $\frac{1}{2}T_1 - T_2$ From reasonable σ^2 not just sum + or - 6
	(ii)	Use sum of 5 Ts $\mu = 9.4$ $\sigma^2 = 2.5225$ $z = (10 - \mu)/\sigma$ Probability 0.6473, 0.647	M1 A1 A1 M1 A1	5	Standardising, must be σ
	(iii)	Calculation of variance	B1	1	

8	(i)	$s_B^2 = \frac{1}{49}(630.194 - \frac{176.35^2}{50})$	M1		Any equivalent formula
		=0.1675	A1		May be implied by later work
		$H_0: \mu_B - \mu_A = 0, H_1: \mu_B - \mu_A > 0$		B1	aef
		$z=0.115/\sqrt{(0.049/40 + 0.1675/50)}$	M1		Standardising but not from pooled variance estimate
		=1.700	A1		art 1.70
		$z > 1.645$, reject H_0	M1		Compare correctly with 1.645
		and accept that $\mu_B > \mu_A$	A1	✓ 7	ft their calculated z
--					
	(ii)	$z = 0.09/\sqrt{(0.004575)}$	M1		Correct form
		= 1.331	A1		
		H_0 not rejected for $\alpha < 9.16$	M1	A1	Accept $< 9.2, \leq 9.2$. M1 for correct method for 9.2, A1 for inequality
				4	
	-----	(iii) (a) Not necessary		B1	Ignore any reason
		(b) Not necessary since samples large enough for CLT to be applied (normality of sample means giving normality of difference)	M1		Mention of CLT implied by “sample large”
			A1	3	Sample mean (approx) normal. (Do not award if population or sample said to be normal)