

**ADVANCED SUBSIDIARY GCE UNIT
MATHEMATICS**

Further Pure Mathematics 1
MONDAY 11 JUNE 2007

4725/01

Afternoon

Time: 1 hour 30 minutes

Additional Materials: Answer Booklet (8 pages)
List of Formulae (MF1)

INSTRUCTIONS TO CANDIDATES

- Write your name, centre number and candidate number in the spaces provided on the answer booklet.
- Answer **all** the questions.
- Give non-exact numerical answers correct to 3 significant figures unless a different degree of accuracy is specified in the question or is clearly appropriate.
- You are permitted to use a graphical calculator in this paper.

INFORMATION FOR CANDIDATES

- The number of marks is given in brackets [] at the end of each question or part question.
- The total number of marks for this paper is 72.

ADVICE TO CANDIDATES

- Read each question carefully and make sure you know what you have to do before starting your answer.
- **You are reminded of the need for clear presentation in your answers.**

This document consists of **4** printed pages.

1 The complex number $a + ib$ is denoted by z . Given that $|z| = 4$ and $\arg z = \frac{1}{3}\pi$, find a and b . [4]

2 Prove by induction that, for $n \geq 1$, $\sum_{r=1}^n r^3 = \frac{1}{4}n^2(n+1)^2$. [5]

3 Use the standard results for $\sum_{r=1}^n r$ and $\sum_{r=1}^n r^2$ to show that, for all positive integers n ,

$$\sum_{r=1}^n (3r^2 - 3r + 1) = n^3. \quad [6]$$

4 The matrix \mathbf{A} is given by $\mathbf{A} = \begin{pmatrix} 1 & 1 \\ 3 & 5 \end{pmatrix}$.

(i) Find \mathbf{A}^{-1} . [2]

The matrix \mathbf{B}^{-1} is given by $\mathbf{B}^{-1} = \begin{pmatrix} 1 & 1 \\ 4 & -1 \end{pmatrix}$.

(ii) Find $(\mathbf{AB})^{-1}$. [4]

5 (i) Show that

$$\frac{1}{r} - \frac{1}{r+1} = \frac{1}{r(r+1)}. \quad [1]$$

(ii) Hence find an expression, in terms of n , for

$$\frac{1}{2} + \frac{1}{6} + \frac{1}{12} + \dots + \frac{1}{n(n+1)}. \quad [3]$$

(iii) Hence find the value of $\sum_{r=n+1}^{\infty} \frac{1}{r(r+1)}$. [3]

6 The cubic equation $3x^3 - 9x^2 + 6x + 2 = 0$ has roots α , β and γ .

(i) (a) Write down the values of $\alpha + \beta + \gamma$ and $\alpha\beta + \beta\gamma + \gamma\alpha$. [2]

(b) Find the value of $\alpha^2 + \beta^2 + \gamma^2$. [2]

(ii) (a) Use the substitution $x = \frac{1}{u}$ to find a cubic equation in u with integer coefficients. [2]

(b) Use your answer to part (ii) (a) to find the value of $\frac{1}{\alpha} + \frac{1}{\beta} + \frac{1}{\gamma}$. [2]

7 The matrix \mathbf{M} is given by $\mathbf{M} = \begin{pmatrix} a & 4 & 0 \\ 0 & a & 4 \\ 2 & 3 & 1 \end{pmatrix}$.

(i) Find, in terms of a , the determinant of \mathbf{M} . [3]

(ii) In the case when $a = 2$, state whether \mathbf{M} is singular or non-singular, justifying your answer. [2]

(iii) In the case when $a = 4$, determine whether the simultaneous equations

$$\begin{aligned} ax + 4y &= 6, \\ ay + 4z &= 8, \\ 2x + 3y + z &= 1, \end{aligned}$$

have any solutions. [3]

8 The loci C_1 and C_2 are given by $|z - 3| = 3$ and $\arg(z - 1) = \frac{1}{4}\pi$ respectively.

(i) Sketch, on a single Argand diagram, the loci C_1 and C_2 . [6]

(ii) Indicate, by shading, the region of the Argand diagram for which

$$|z - 3| \leq 3 \text{ and } 0 \leq \arg(z - 1) \leq \frac{1}{4}\pi. \quad [2]$$

9 (i) Write down the matrix, \mathbf{A} , that represents an enlargement, centre $(0, 0)$, with scale factor $\sqrt{2}$. [1]

(ii) The matrix \mathbf{B} is given by $\mathbf{B} = \begin{pmatrix} \frac{1}{2}\sqrt{2} & \frac{1}{2}\sqrt{2} \\ -\frac{1}{2}\sqrt{2} & \frac{1}{2}\sqrt{2} \end{pmatrix}$. Describe fully the geometrical transformation represented by \mathbf{B} . [3]

(iii) Given that $\mathbf{C} = \mathbf{AB}$, show that $\mathbf{C} = \begin{pmatrix} 1 & 1 \\ -1 & 1 \end{pmatrix}$. [1]

(iv) Draw a diagram showing the unit square and its image under the transformation represented by \mathbf{C} . [2]

(v) Write down the determinant of \mathbf{C} and explain briefly how this value relates to the transformation represented by \mathbf{C} . [2]

10 (i) Use an algebraic method to find the square roots of the complex number $16 + 30i$. [6]

(ii) Use your answers to part (i) to solve the equation $z^2 - 2z - (15 + 30i) = 0$, giving your answers in the form $x + iy$. [5]

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (OCR) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

OCR is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.