

**ADVANCED GCE**

**4753/01**

**MATHEMATICS (MEI)**

Methods for Advanced Mathematics (C3)

**MONDAY 2 JUNE 2008**

Morning

Time: 1 hour 30 minutes

**Additional materials (enclosed):** None

**Additional materials (required):**

Answer Booklet (8 pages)

Graph paper

MEI Examination Formulae and Tables (MF2)

**INSTRUCTIONS TO CANDIDATES**

- Write your name in capital letters, your Centre Number and Candidate Number in the spaces provided on the Answer Booklet.
- Read each question carefully and make sure you know what you have to do before starting your answer.
- Answer **all** the questions.
- You are permitted to use a graphical calculator in this paper.
- Final answers should be given to a degree of accuracy appropriate to the context.

**INFORMATION FOR CANDIDATES**

- The number of marks is given in brackets [ ] at the end of each question or part question.
- The total number of marks for this paper is **72**.
- You are advised that an answer may receive **no marks** unless you show sufficient detail of the working to indicate that a correct method is being used.

This document consists of **4** printed pages.

## Section A (36 marks)

- 1 Solve the inequality  $|2x - 1| \leq 3$ . [4]
- 2 Find  $\int xe^{3x} dx$ . [4]
- 3 (i) State the algebraic condition for the function  $f(x)$  to be an even function.  
What geometrical property does the graph of an even function have? [2]
- (ii) State whether the following functions are odd, even or neither.  
(A)  $f(x) = x^2 - 3$   
(B)  $g(x) = \sin x + \cos x$   
(C)  $h(x) = \frac{1}{x + x^3}$  [3]
- 4 Show that  $\int_1^4 \frac{x}{x^2 + 2} dx = \frac{1}{2} \ln 6$ . [4]
- 5 Show that the curve  $y = x^2 \ln x$  has a stationary point when  $x = \frac{1}{\sqrt{e}}$ . [6]
- 6 In a chemical reaction, the mass  $m$  grams of a chemical after  $t$  minutes is modelled by the equation  
$$m = 20 + 30e^{-0.1t}.$$
  
(i) Find the initial mass of the chemical.  
What is the mass of chemical in the long term? [3]
- (ii) Find the time when the mass is 30 grams. [3]
- (iii) Sketch the graph of  $m$  against  $t$ . [2]
- 7 Given that  $x^2 + xy + y^2 = 12$ , find  $\frac{dy}{dx}$  in terms of  $x$  and  $y$ . [5]

## Section B (36 marks)

8 Fig. 8 shows the curve  $y = f(x)$ , where  $f(x) = \frac{1}{1 + \cos x}$ , for  $0 \leq x \leq \frac{1}{2}\pi$ .

P is the point on the curve with  $x$ -coordinate  $\frac{1}{3}\pi$ .




Fig. 8

- (i) Find the  $y$ -coordinate of P. [1]
- (ii) Find  $f'(x)$ . Hence find the gradient of the curve at the point P. [5]
- (iii) Show that the derivative of  $\frac{\sin x}{1 + \cos x}$  is  $\frac{1}{1 + \cos x}$ . Hence find the exact area of the region enclosed by the curve  $y = f(x)$ , the  $x$ -axis, the  $y$ -axis and the line  $x = \frac{1}{3}\pi$ . [7]
- (iv) Show that  $f^{-1}(x) = \arccos\left(\frac{1}{x} - 1\right)$ . State the domain of this inverse function, and add a sketch of  $y = f^{-1}(x)$  to a copy of Fig. 8. [5]

[Question 9 is printed overleaf.]

9 The function  $f(x)$  is defined by  $f(x) = \sqrt{4 - x^2}$  for  $-2 \leq x \leq 2$ .

- (i) Show that the curve  $y = \sqrt{4 - x^2}$  is a semicircle of radius 2, and explain why it is not the whole of this circle. [3]

Fig. 9 shows a point  $P(a, b)$  on the semicircle. The tangent at  $P$  is shown.




Fig. 9

- (ii) (A) Use the gradient of  $OP$  to find the gradient of the tangent at  $P$  in terms of  $a$  and  $b$ .  
 (B) Differentiate  $\sqrt{4 - x^2}$  and deduce the value of  $f'(a)$ .  
 (C) Show that your answers to parts (A) and (B) are equivalent. [6]

The function  $g(x)$  is defined by  $g(x) = 3f(x - 2)$ , for  $0 \leq x \leq 4$ .

- (iii) Describe a sequence of two transformations that would map the curve  $y = f(x)$  onto the curve  $y = g(x)$ .

Hence sketch the curve  $y = g(x)$ . [6]

- (iv) Show that if  $y = g(x)$  then  $9x^2 + y^2 = 36x$ . [3]