


# 4753 (C3) Methods for Advanced Mathematics

## Section A

|                                                                                                                                                                                                                 |                                                   |                                                                                                                                                                                                                                                                                                           |
|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <p><b>1</b></p> $\int_0^{\pi/6} \sin 3x \, dx = \left[ -\frac{1}{3} \cos 3x \right]_0^{\pi/6}$ $= -\frac{1}{3} \cos \frac{\pi}{2} + \frac{1}{3} \cos 0$ $= \frac{1}{3}$                                         | <p>B1</p> <p>M1</p> <p>A1cao<br/>[3]</p>          | $\left[ -\frac{1}{3} \cos 3x \right] \text{ or } \left[ -\frac{1}{3} \cos u \right]$<br>substituting correct limits in $\pm k \cos \dots$<br>0.33 or better.                                                                                                                                              |
| <p><b>2(i)</b></p> $100 = Ae^0 = A \Rightarrow A = 100$<br>$50 = 100 e^{-1500k}$<br>$\Rightarrow e^{-1500k} = 0.5$<br>$\Rightarrow -1500k = \ln 0.5$<br>$\Rightarrow k = -\ln 0.5 / 1500 = 4.62 \times 10^{-4}$ | <p>M1A1</p> <p>M1</p> <p>M1</p> <p>A1<br/>[5]</p> | $50 = A e^{-1500k}$ ft their 'A' if used<br>taking lns correctly<br>0.00046 or better                                                                                                                                                                                                                     |
| <p><b>(ii)</b></p> $1 = 100e^{-kt}$<br>$\Rightarrow -kt = \ln 0.01$<br>$\Rightarrow t = -\ln 0.01 / k$<br>= 9966 years                                                                                          | <p>M1</p> <p>M1</p> <p>A1<br/>[3]</p>             | ft their A and k<br>taking lns correctly<br>art 9970                                                                                                                                                                                                                                                      |
| <p><b>3</b></p>                                                                                                              | <p>M1</p> <p>B1</p> <p>A1<br/>[3]</p>             | Can use degrees or radians<br>reasonable shape (condone extra range)<br>passes through $(-1, 2\pi)$ , $(0, \pi)$ and $(1, 0)$<br>good sketches – look for curve reasonably vertical at $(-1, 2\pi)$ and $(1, 0)$ , negative gradient at $(0, \pi)$ . Domain and range must be clearly marked and correct. |
| <p><b>4</b></p> $g(x) = 2 x-1 $<br>$\Rightarrow b = 2 0-1  = 2$ or $(0, 2)$<br>$2 x-1  = 0$<br>$\Rightarrow x = 1$ , so $a = 1$ or $(1, 0)$                                                                     | <p>B1</p> <p>M1</p> <p>A1<br/>[3]</p>             | Allow unsupported answers.<br>www<br>$ x  = 1$ is A0<br>www                                                                                                                                                                                                                                               |

|                                                                                                                                                                                                                                                      |                                                   |                                                                                                                                                           |
|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>5(i)</b> $e^{2y} = 1 + \sin x$<br>$\Rightarrow 2e^{2y} dy/dx = \cos x$<br>$\Rightarrow dy/dx = \frac{\cos x}{2e^{2y}}$                                                                                                                            | M1<br>B1<br><br>A1<br>[3]                         | Their $2e^{2y} \times dy/dx$<br>$2e^{2y}$<br><br>o.e. cao                                                                                                 |
| <b>(ii)</b> $2y = \ln(1 + \sin x)$<br>$\Rightarrow y = \frac{1}{2} \ln(1 + \sin x)$<br>$\Rightarrow dy/dx = \frac{1}{2} \frac{\cos x}{1 + \sin x}$<br>$= \frac{\cos x}{2e^{2y}}$ as before                                                           | B1<br>M1<br><br>B1<br>E1<br>[4]                   | chain rule (can be within 'correct' quotient rule with $dv/dx = 0$ )<br>$1/u$ or $1/(1 + \sin x)$ soi<br>www                                              |
| <b>6</b> $f(x) = \frac{\frac{x+1}{x-1} + 1}{\frac{x+1}{x-1} - 1}$<br>$= \frac{x+1+x-1}{x+1-x+1}$<br>$= 2x/2 = x^*$<br><br>$f^{-1}(x) = f(x)$<br><br>Symmetrical about $y = x$ .                                                                      | M1<br><br>M1<br><br>E1<br><br>B1<br><br>B1<br>[5] | correct expression<br><br>without subsidiary denominators<br>e.g. $= \frac{x+1+x-1}{x-1} \times \frac{x-1}{x+1-x+1}$<br><br>stated, or shown by inverting |
| <b>7(i)</b> (A) $(x-y)(x^2 + xy + y^2)$<br>$= x^3 + x^2y + xy^2 - yx^2 - xy^2 - y^3$<br>$= x^3 - y^3$ *<br><br>(B) $(x + \frac{1}{2}y)^2 + \frac{3}{4}y^2$<br>$= x^2 + xy + \frac{1}{4}y^2 + \frac{3}{4}y^2$<br>$= x^2 + xy + y^2$                   | M1<br><br>E1<br><br>M1<br><br>E1<br>[4]           | expanding - allow tabulation<br><br>www<br><br>$(x + \frac{1}{2}y)^2 = x^2 + \frac{1}{2}xy + \frac{1}{2}xy + \frac{1}{4}y^2$ o.e.<br><br>cao www          |
| <b>(ii)</b> $x^3 - y^3 = (x-y)[(x + \frac{1}{2}y)^2 + \frac{3}{4}y^2]$<br>$(x + \frac{1}{2}y)^2 + \frac{3}{4}y^2 > 0$ [as squares $\geq 0$ ]<br><br>$\Rightarrow$ if $x - y > 0$ then $x^3 - y^3 > 0$<br>$\Rightarrow$ if $x > y$ then $x^3 > y^3$ * | M1<br><br>M1<br><br>E1<br>[3]                     | substituting results of (i)                                                                                                                               |

|                                                                                                                                                                                                                                                                                                                                                         |                                                               |                                                                                                                                                                           |
|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>8(i)</b> A: $1 + \ln x = 0$<br>$\Rightarrow \ln x = -1$ so A is $(e^{-1}, 0)$<br>$\Rightarrow x = e^{-1}$<br>B: $x = 0, y = e^{0-1} = e^{-1}$ so B is $(0, e^{-1})$<br><br>C: $f(1) = e^{1-1} = e^0 = 1$<br>$g(1) = 1 + \ln 1 = 1$                                                                                                                   | M1<br><br>A1<br>B1<br><br>E1<br>E1<br>[5]                     | SC1 if obtained using symmetry<br>condone use of symmetry<br>Penalise $A = e^{-1}$ , $B = e^{-1}$ , or co-ords wrong way round, but condone labelling errors.             |
| <b>(ii)</b> Either by inversion:<br>e.g. $y = e^{x-1} \quad x \leftrightarrow y$<br>$x = e^{y-1}$<br>$\Rightarrow \ln x = y - 1$<br>$\Rightarrow 1 + \ln x = y$                                                                                                                                                                                         | M1<br>E1                                                      | taking lns or exps                                                                                                                                                        |
| or by composing<br>e.g. $f(g(x)) = f(1 + \ln x)$<br>$= e^{1 + \ln x - 1}$<br>$= e^{\ln x} = x$                                                                                                                                                                                                                                                          | M1<br><br>E1<br>[2]                                           | $e^{1 + \ln x - 1}$ or $1 + \ln(e^{x-1})$                                                                                                                                 |
| <b>(iii)</b> $\int_0^1 e^{x-1} dx = \left[ e^{x-1} \right]_0^1$<br>$= e^0 - e^{-1}$<br>$= 1 - e^{-1}$                                                                                                                                                                                                                                                   | M1<br><br>M1<br>A1cao<br>[3]                                  | $\left[ e^{x-1} \right]$ o.e or $u = x - 1 \Rightarrow \left[ e^u \right]$<br>substituting correct limits for $x$ or $u$<br>o.e. not $e^0$ , must be exact.               |
| <b>(iv)</b> $\int \ln x dx = \int \ln x \frac{d}{dx}(x) dx$<br>$= x \ln x - \int x \cdot \frac{1}{x} dx$<br>$= x \ln x - x + c$<br>$\Rightarrow \int_{e^{-1}}^1 g(x) dx = \int_{e^{-1}}^1 (1 + \ln x) dx$<br>$= \left[ x + x \ln x - x \right]_{e^{-1}}^1$<br>$= \left[ x \ln x \right]_{e^{-1}}^1$<br>$= 1 \ln 1 - e^{-1} \ln(e^{-1})$<br>$= e^{-1} *$ | M1<br><br>A1<br><br>A1cao<br><br>B1ft<br><br>DM1<br>E1<br>[6] | parts: $u = \ln x, du/dx = 1/x, v = x, dv/dx = 1$<br><br>condone no 'c'<br><br>ft their ' $x \ln x - x$ ' (provided 'algebraic')<br><br>substituting limits dep B1<br>www |
| <b>(v)</b> Area = $\int_0^1 f(x) dx - \int_{e^{-1}}^1 g(x) dx$<br>$= (1 - e^{-1}) - e^{-1}$<br>$= 1 - 2/e$                                                                                                                                                                                                                                              | M1<br><br>A1cao                                               | Must have correct limits<br><br>0.264 or better.                                                                                                                          |
| or<br>Area OCB = area under curve – triangle<br>$= 1 - e^{-1} - \frac{1}{2} \times 1 \times 1$<br>$= \frac{1}{2} - e^{-1}$<br><br>or<br>Area OAC = triangle – area under curve<br>$= \frac{1}{2} \times 1 \times 1 - e^{-1}$<br>$= \frac{1}{2} - e^{-1}$<br>Total area = $2(\frac{1}{2} - e^{-1}) = 1 - 2/e$                                            | M1<br><br>A1cao<br>[2]                                        | OCA or OCB = $\frac{1}{2} - e^{-1}$<br><br>0.264 or better                                                                                                                |

|                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                          |                                                              |                                                                                                                                                                                                                                                                                                                             |
|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| 9(i) $a = 1/3$                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                           | B1<br>[1]                                                    | or 0.33 or better                                                                                                                                                                                                                                                                                                           |
| (ii) $\frac{dy}{dx} = \frac{(3x-1)2x-x^2 \cdot 3}{(3x-1)^2}$<br>$= \frac{6x^2-2x-3x^2}{(3x-1)^2}$<br>$= \frac{3x^2-2x}{(3x-1)^2}$<br>$= \frac{x(3x-2)}{(3x-1)^2} *$                                                                                                                                                                                                                                                                                                                                                                                                                                      | M1<br>A1<br><br><br><br>E1<br>[3]                            | quotient rule<br><br><br><br><br>www – must show both steps; penalise missing brackets.                                                                                                                                                                                                                                     |
| (iii) $dy/dx = 0$ when $x(3x-2) = 0$<br>$\Rightarrow x = 0$ or $x = 2/3$ , so at P, $x = 2/3$<br>when $x = \frac{2}{3}$ , $y = \frac{(2/3)^2}{3 \times (2/3) - 1} = \frac{4}{9}$<br><br>when $x = 0.6$ , $dy/dx = -0.1875$<br>when $x = 0.8$ , $dy/dx = 0.1633$<br>Gradient increasing $\Rightarrow$ minimum                                                                                                                                                                                                                                                                                             | M1<br>A1<br>M1<br>A1cao<br><br>B1<br>B1<br>E1<br>[7]         | if denom = 0 also then M0<br>o.e e.g. 0.6, but must be exact<br>o.e e.g. 0.4, but must be exact<br><br>-3/16, or -0.19 or better<br>8/49 or 0.16 or better<br>o.e. e.g. 'from negative to positive'. Allow ft on their gradients, provided -ve and +ve respectively.<br>Accept table with indications of signs of gradient. |
| (iv) $\int \frac{x^2}{3x-1} dx$ $u = 3x-1 \Rightarrow du = 3dx$<br>$\frac{(u+1)^2}{9} \cdot \frac{1}{3} du$<br>$= \frac{1}{27} \int \frac{(u+1)^2}{u} du = \frac{1}{27} \int \frac{u^2+2u+1}{u} du$<br>$= \frac{1}{27} \int (u+2+\frac{1}{u}) du *$<br><br>Area = $\int_{2/3}^1 \frac{x^2}{3x-1} dx$<br>When $x = 2/3$ , $u = 1$ , when $x = 1$ , $u = 2$<br>$= \frac{1}{27} \int_1^2 (u+2+1/u) du$<br>$= \frac{1}{27} \left[ \frac{1}{2} u^2 + 2u + \ln u \right]_1^2$<br>$= \frac{1}{27} [(2+4+\ln 2) - (\frac{1}{2} + 2 + \ln 1)]$<br>$= \frac{1}{27} (3\frac{1}{2} + \ln 2) [= \frac{7+2\ln 2}{54}]$ | B1<br><br>M1<br>M1<br>E1<br><br><br>B1<br>M1<br>A1cao<br>[7] | $\frac{(u+1)^2}{9} \cdot \frac{1}{u}$ o.e.<br>$\times 1/3 (du)$<br>expanding<br>Condone missing $du$ 's<br><br><br>$\left[ \frac{1}{2} u^2 + 2u + \ln u \right]$<br>substituting correct limits, dep integration<br>o.e., but must evaluate $\ln 1 = 0$ and collect terms.                                                  |