

4727 Further Pure Mathematics 3

1	$\left(\frac{1}{2}\sqrt{3} + \frac{1}{2}i\right)^{\frac{1}{3}} = \left(\cos \frac{1}{6}\pi + i \sin \frac{1}{6}\pi\right)^{\frac{1}{3}}$	B1	For $\arg z = \frac{1}{6}\pi$ seen or implied
	$= \cos \frac{1}{18}\pi + i \sin \frac{1}{18}\pi,$	M1	For dividing $\arg z$ by 3
	$\cos \frac{13}{18}\pi + i \sin \frac{13}{18}\pi,$	A1	For any one correct root
	$\cos \frac{25}{18}\pi + i \sin \frac{25}{18}\pi$	A1 4	For 2 other roots and no more in range $0 \leq \theta < 2\pi$
4			
2 (i)	$\frac{1}{5}e^{-\frac{1}{3}\pi i}$	B1 1	For stating correct inverse in the form $re^{i\theta}$
(ii)	$r_1e^{i\theta} \times r_2e^{i\phi} = r_1r_2e^{i(\theta+\phi)}$	M1	For stating 2 distinct elements multiplied
		A1 2	For showing product of correct form
(iii)	$Z^2 = e^{2i\gamma}$	B1	For $e^{2i\gamma}$ seen or implied
	$\Rightarrow e^{2i\gamma-2\pi i}$	B1 2	For correct answer. aef
5			
3 (i)	$[6-4\lambda, -7+8\lambda, -10+7\lambda]$ on p	B1	For point on l seen or implied
	$\Rightarrow 3(6-4\lambda) - 4(-7+8\lambda) - 2(-10+7\lambda) = 8$	M1	For substituting into equation of p
	$\Rightarrow \lambda = 1 \Rightarrow (2, 1, -3)$	A1 3	For correct point. Allow position vector
(ii)	METHOD 1		
	$\mathbf{n} = [-4, 8, 7] \times [3, -4, -2]$	M1*	For direction of l and normal of p seen
		M1	For attempting to find $\mathbf{n}_1 \times \mathbf{n}_2$
		(*dep)	
	$\mathbf{n} = k[12, 13, -8]$	A1	For correct vector
	$(2, 1, -3)$ OR $(6, -7, -10)$	M1	For finding scalar product of their point on l with their attempt at \mathbf{n} , or equivalent
	$\Rightarrow 12x + 13y - 8z = 61$	A1 5	For correct equation, aef cartesian
	METHOD 2		
	$\mathbf{r} = [2, 1, -3]$ OR $[6, -7, -10]$	M1	For stating eqn of plane in parametric form (may be implied by next stage), using $[2, 1, -3]$ (ft from (i))
	$+ \lambda[-4, 8, 7] + \mu[3, -4, -2]$	A1✓	Or $[6, -7, -10]$, \mathbf{n}_1 and \mathbf{n}_2 (as above)
	$x = 2 - 4\lambda + 3\mu$	M1	For writing as 3 linear equations
	$y = 1 + 8\lambda - 4\mu$	M1	For attempting to eliminate λ and μ
	$z = -3 + 7\lambda - 2\mu$		
	$\Rightarrow 12x + 13y - 8z = 61$	A1	For correct equation aef cartesian
	METHOD 3		
	$3(6+3\mu) - 4(-7-4\mu) - 2(-10-2\mu) = 8$	M1	For finding foot of perpendicular from point on l to p
	$\Rightarrow \mu = -2 \Rightarrow (0, 1, -6)$	A1	For correct point or position vector
	From 3 points $(2, 1, -3)$, $(6, -7, -10)$, $(0, 1, -6)$,		
	\mathbf{n} = vector product of 2 of		
	$[2, 0, 3]$, $[6, -8, -4]$, $[-4, 8, 7]$	M1	Use vector product of 2 vectors in plane
	$\Rightarrow \mathbf{n} = k[12, 13, -8]$		
	$(2, 1, -3)$ OR $(6, -7, -10)$	M1	For finding scalar product of their point on l with their attempt at \mathbf{n} , or equivalent
	$\Rightarrow 12x + 13y - 8z = 61$	A1	For correct equation aef cartesian
8			

4	(i)	IF $e^{\int \frac{1}{1-x^2} dx} = e^{\frac{1}{2} \ln \frac{1+x}{1-x}} = \left(\frac{1+x}{1-x} \right)^{\frac{1}{2}}$	M1	For IF stated or implied. Allow $\pm \int$ and omission of
			A1	dx For integration and simplification to AG (intermediate step must be seen)
<hr/>				
	(ii)	$\frac{d}{dx} \left(y \left(\frac{1+x}{1-x} \right)^{\frac{1}{2}} \right) = (1+x)^{\frac{1}{2}}$	M1*	For multiplying both sides by IF
		$y \left(\frac{1+x}{1-x} \right)^{\frac{1}{2}} = \frac{2}{3} (1+x)^{\frac{3}{2}} + c$	M1	For integrating RHS to $k(1+x)^n$
			A1	For correct equation (including + c)
		$(0, 2) \Rightarrow 2 = \frac{2}{3} + c \Rightarrow c = \frac{4}{3}$	M1 (*dep)	In either order: For substituting (0, 2) into their GS (including +c)
			M1 (*dep)	For dividing solution through by IF, including dividing c or their numerical value for c
		$y = \frac{2}{3} (1+x) (1-x)^{\frac{1}{2}} + \frac{4}{3} \left(\frac{1-x}{1+x} \right)^{\frac{1}{2}}$	A1	For correct solution aef (even unsimplified) in form $y = f(x)$
<hr/>				
			8	
<hr/>				
5	(i)	$m^2 - 6m + 9 (= 0) \Rightarrow m = 3$	M1	For attempting to solve correct auxiliary equation
			A1	For correct m
		CF = $(A + Bx)e^{3x}$	A1	For correct CF
<hr/>				
	(ii)	ke^{3x} and kxe^{3x} both appear in CF	B1	For correct statement
<hr/>				
	(iii)	$y = kx^2e^{3x} \Rightarrow y' = 2kxe^{3x} + 3kx^2e^{3x}$	M1	For differentiating kx^2e^{3x} twice
			A1	For correct y' aef
		$\Rightarrow y'' = 2ke^{3x} + 12kxe^{3x} + 9kx^2e^{3x}$	A1	For correct y'' aef
		\Rightarrow		
		$ke^{3x} (2 + 12x + 9x^2 - 12x - 18x^2 + 9x^2) = e^{3x}$	M1	For substituting y'', y', y into DE
		$\Rightarrow k = \frac{1}{2}$	A1	For correct k
<hr/>				
			9	

6 (i)	METHOD 1		
	$\mathbf{n}_1 = [1, 1, 0] \times [1, -5, -2]$	M1	For attempting to find vector product of the pair of direction vectors
	$= [-2, 2, -6] = k[1, -1, 3]$	A1	For correct \mathbf{n}_1
	Use $(2, 2, 1)$	M1	For substituting a point into equation
	$\Rightarrow \mathbf{r} \cdot [-2, 2, -6] = -6 \Rightarrow \mathbf{r} \cdot [1, -1, 3] = 3$	A1 4	For correct equation. aef in this form
	METHOD 2		
	$x = 2 + \lambda + \mu$	M1	For writing as 3 linear equations
	$y = 2 + \lambda - 5\mu$	M1	For attempting to eliminate λ and μ
	$z = 1 - 2\mu$		
	$\Rightarrow x - y + 3z = 3$	A1	For correct cartesian equation
$\Rightarrow \mathbf{r} \cdot [1, -1, 3] = 3$	A1	For correct equation. aef in this form	
(ii)	For $\mathbf{r} = \mathbf{a} + t\mathbf{b}$		
	METHOD 1		
	$\mathbf{b} = [1, -1, 3] \times [7, 17, -3]$	M1	For attempting to find $\mathbf{n}_1 \times \mathbf{n}_2$
	$= k[2, -1, -1]$	A1✓	For a correct vector. ft from \mathbf{n}_1 in (i)
	e.g. x, y or $z = 0$ in $\begin{cases} x - y + 3z = 3 \\ 7x + 17y - 3z = 21 \end{cases}$	M1	For attempting to find a point on the line
	$\Rightarrow \mathbf{a} = \left[0, \frac{3}{2}, \frac{3}{2}\right]$ OR $[3, 0, 0]$ OR $[1, 1, 1]$	A1✓	For a correct vector. ft from equation in (i) SR a correct vector may be stated without working
	Line is (e.g.) $\mathbf{r} = [1, 1, 1] + t[2, -1, -1]$	A1✓ 5	For stating equation of line ft from \mathbf{a} and \mathbf{b} SR for $\mathbf{a} = [2, 2, 1]$ stated award M0
	METHOD 2		
	Solve $\begin{cases} x - y + 3z = 3 \\ 7x + 17y - 3z = 21 \end{cases}$	M1	In either order: For attempting to solve equations
	by eliminating one variable (e.g. z) Use parameter for another variable (e.g. x) to find other variables in terms of t	M1	For attempting to find parametric solution
	(eg) $y = \frac{3}{2} - \frac{1}{2}t, z = \frac{3}{2} - \frac{1}{2}t$	A1✓ A1✓	For correct expression for one variable For correct expression for the other variable ft from equation in (i) for both
	Line is (eg) $\mathbf{r} = \left[0, \frac{3}{2}, \frac{3}{2}\right] + t[2, -1, -1]$	A1✓	For stating equation of line. ft from parametric solutions
	METHOD 3		
	eg x, y or $z = 0$ in $\begin{cases} x - y + 3z = 3 \\ 7x + 17y - 3z = 21 \end{cases}$	M1	For attempting to find a point on the line
	$\Rightarrow \mathbf{a} = \left[0, \frac{3}{2}, \frac{3}{2}\right]$ OR $[3, 0, 0]$ OR $[1, 1, 1]$	A1✓	For a correct vector. ft from equation in (i) SR a correct vector may be stated without working SR for $\mathbf{a} = [2, 2, 1]$ stated award M0
	eg $[3, 0, 0] - [1, 1, 1]$	M1	For finding another point on the line and using it with the one already found to find \mathbf{b}
	$\mathbf{b} = k[2, -1, -1]$	A1✓	For a correct vector. ft from equation in (i)
	Line is (eg) $\mathbf{r} = [1, 1, 1] + t[2, -1, -1]$	A1✓	For stating equation of line. ft from \mathbf{a} and \mathbf{b}

6 (ii) contd	METHOD 4		
	A point on Π_1 is $[2 + \lambda + \mu, 2 + \lambda - 5\mu, 1 - 2\mu]$	M1	For using parametric form for Π_1 and substituting into Π_2
	On $\Pi_2 \Rightarrow$ $[2 + \lambda + \mu, 2 + \lambda - 5\mu, 1 - 2\mu] \cdot [7, 17, -3] = 21$	A1	For correct unsimplified equation
	$\Rightarrow \lambda - 3\mu = -1$	A1	For correct equation
	Line is (e.g.) $\mathbf{r} = [2, 2, 1] + (3\mu - 1)[1, 1, 0] + \mu[1, -5, -2]$	M1	For substituting into Π_1 for λ or μ
	$\Rightarrow \mathbf{r} = [1, 1, 1] \text{ or } \left[\frac{7}{3}, \frac{1}{3}, \frac{1}{3}\right] + t[2, -1, -1]$	A1	For stating equation of line
9			
7 (i)	$\cos 3\theta + i \sin 3\theta = c^3 + 3ic^2s - 3cs^2 - is^3$	M1	For using de Moivre with $n = 3$
	$\Rightarrow \cos 3\theta = c^3 - 3cs^2$ and $\sin 3\theta = 3c^2s - s^3$	A1	For both expressions in this form (seen or implied) SR For expressions found without de Moivre M0 A0
	$\Rightarrow \tan 3\theta = \frac{3c^2s - s^3}{c^3 - 3cs^2}$	M1	For expressing $\frac{\sin 3\theta}{\cos 3\theta}$ in terms of c and s
	$\tan 3\theta = \frac{3 \tan \theta - \tan^3 \theta}{1 - 3 \tan^2 \theta} = \frac{\tan \theta (3 - \tan^2 \theta)}{1 - 3 \tan^2 \theta}$	A1	4 For simplifying to AG
(ii) (a)	$\theta = \frac{1}{12}\pi \Rightarrow \tan 3\theta = 1$		
	$\Rightarrow 1 - 3t^2 = t(3 - t^2) \Rightarrow$ $t^3 - 3t^2 - 3t + 1 = 0$	B1	1 For both stages correct AG
(b)	$(t+1)(t^2 - 4t + 1) = 0$	M1	For attempt to factorise cubic
		A1	For correct factors
	$\Rightarrow (t = -1), t = 2 \pm \sqrt{3}$	A1	For correct roots of quadratic
	– sign for smaller root \Rightarrow $\tan \frac{1}{12}\pi = 2 - \sqrt{3}$	A1	4 For choice of – sign and correct root AG
(iii)	$dt = (1 + t^2) d\theta$	B1	For differentiation of substitution and use of $\sec^2 \theta = 1 + \tan^2 \theta$
	$\Rightarrow \int_0^{\frac{1}{12}\pi} \tan 3\theta d\theta$	B1	For integral with correct θ limits seen
	$= \left[\frac{1}{3} \ln(\sec 3\theta) \right]_0^{\frac{1}{12}\pi} = \frac{1}{3} \ln\left(\sec \frac{1}{4}\pi\right)$	M1	For integrating to $k \ln(\sec 3\theta)$ OR $k \ln(\cos 3\theta)$
	$= \frac{1}{3} \ln \sqrt{2} = \frac{1}{6} \ln 2$	M1	For substituting limits and $\sec \frac{1}{4}\pi = \sqrt{2}$ OR $\cos \frac{1}{4}\pi = \frac{1}{\sqrt{2}}$ seen
		A1	5 For correct answer aef
14			

8	(i)	$a^2 = (ap)^2 = apap \Rightarrow a = pap$	B1	For use of given properties to obtain AG																									
		$p^2 = (ap)^2 = apap \Rightarrow p = apa$	B1 2	For use of given properties to obtain AG SR allow working from AG to obtain relevant properties																									
<hr/>																													
(ii)		$(p^2)^2 = p^4 = e \Rightarrow \text{order } p^2 = 2$	B1	For correct order with no incorrect working seen																									
		$(a^2)^2 = (p^2)^2 = e \Rightarrow \text{order } a = 4$	B1	For correct order with no incorrect working seen																									
		$(ap)^4 = a^4 = e \Rightarrow \text{order } ap = 4$	B1	For correct order with no incorrect working seen																									
		$(ap^2)^2 = ap^2 ap^2 = ap \cdot a \cdot p = a^2$	M1	For relevant use of (i) or given properties																									
		OR $ap^2 = a \cdot a^2 = a^3 \Rightarrow$	A1 5	For correct order with no incorrect working seen																									
		$(ap^2)^2 = a^6 = a^2$																											
<hr/>																													
(iii)	METHOD 1		M2	For use of the given properties to simplify p^2 and ap^2																									
	$p^2 = a^2, ap^2 = a^3$																												
	$\Rightarrow \{e, a, p^2, ap^2\} = \{e, a, a^2, a^3\}$		A1	For obtaining a^2 and a^3																									
	which is a cyclic group		A1 4	For justifying that the set is a group																									
<hr/>																													
METHOD 2																													
		<table><tr><td></td><td>e</td><td>a</td><td>p^2</td><td>ap^2</td></tr><tr><td>e</td><td>e</td><td>a</td><td>p^2</td><td>ap^2</td></tr><tr><td>a</td><td>a</td><td>p^2</td><td>ap^2</td><td>e</td></tr><tr><td>p^2</td><td>p^2</td><td>ap^2</td><td>e</td><td>a</td></tr><tr><td>ap^2</td><td>ap^2</td><td>e</td><td>a</td><td>p^2</td></tr></table>		e	a	p^2	ap^2	e	e	a	p^2	ap^2	a	a	p^2	ap^2	e	p^2	p^2	ap^2	e	a	ap^2	ap^2	e	a	p^2	M1	For attempting closure
	e	a	p^2	ap^2																									
e	e	a	p^2	ap^2																									
a	a	p^2	ap^2	e																									
p^2	p^2	ap^2	e	a																									
ap^2	ap^2	e	a	p^2																									
				A1	For all 9 non-trivial products seen																								
					For all 16 products correct																								
Completed table is a cyclic group			B2	For justifying that the set is a group																									
<hr/>																													
METHOD 3																													
		<table><tr><td></td><td>e</td><td>a</td><td>p^2</td><td>ap^2</td></tr><tr><td>e</td><td>e</td><td>a</td><td>p^2</td><td>ap^2</td></tr><tr><td>a</td><td>a</td><td>p^2</td><td>ap^2</td><td>e</td></tr><tr><td>p^2</td><td>p^2</td><td>ap^2</td><td>e</td><td>a</td></tr><tr><td>ap^2</td><td>ap^2</td><td>e</td><td>a</td><td>p^2</td></tr></table>		e	a	p^2	ap^2	e	e	a	p^2	ap^2	a	a	p^2	ap^2	e	p^2	p^2	ap^2	e	a	ap^2	ap^2	e	a	p^2	M1	For attempting closure
	e	a	p^2	ap^2																									
e	e	a	p^2	ap^2																									
a	a	p^2	ap^2	e																									
p^2	p^2	ap^2	e	a																									
ap^2	ap^2	e	a	p^2																									
				A1	For all 9 non-trivial products seen																								
					For all 16 products correct																								
Identity = e			B1	For stating identity																									
Inverses exist since			B1	For justifying inverses ($e^{-1} = e$ may be assumed)																									
EITHER: e is in each row/column																													
OR: p^2 is self-inverse; a, ap^2 form an																													
inverse pair																													

(iv)	METHOD 1	M1	For attempting to find a non-commutative pair of elements, at least one involving a (may be embedded in a full or partial table)
	e.g. $\left. \begin{array}{l} a \cdot ap = a^2 p = p^3 \\ ap \cdot a = p \end{array} \right\} \Rightarrow \text{not}$	M1	For simplifying elements both ways round
	commutative	B1	For a correct pair of non-commutative elements
		A1 4	For stating Q non-commutative, with a clear argument
<hr/>			
	METHOD 2		
	Assume commutativity, so (eg) $ap = pa$	M1	For setting up proof by contradiction
	(i) \Rightarrow		
	$p = ap \cdot a \Rightarrow p = pa \cdot a = pa^2 = pp^2 = p^3$	M1	For using (i) and/or given properties
	But p and p^3 are distinct	B1	For obtaining and stating a contradiction
	$\Rightarrow Q$ is non-commutative	A1	For stating Q non-commutative, with a clear argument
<hr/>			
15			
<hr/>			