GCE

Mathematics

Advanced GCE

Unit 4727: Further Pure Mathematics 3

Mark Scheme for June 2013

OCR (Oxford Cambridge and RSA) is a leading UK awarding body, providing a wide range of qualifications to meet the needs of candidates of all ages and abilities. OCR qualifications include AS/A Levels, Diplomas, GCSEs, Cambridge Nationals, Cambridge Technicals, Functional Skills, Key Skills, Entry Level qualifications, NVQs and vocational qualifications in areas such as IT, business, languages, teaching/training, administration and secretarial skills.

It is also responsible for developing new specifications to meet national requirements and the needs of students and teachers. OCR is a not-for-profit organisation; any surplus made is invested back into the establishment to help towards the development of qualifications and support, which keep pace with the changing needs of today's society.

This mark scheme is published as an aid to teachers and students, to indicate the requirements of the examination. It shows the basis on which marks were awarded by examiners. It does not indicate the details of the discussions which took place at an examiners' meeting before marking commenced.

All examiners are instructed that alternative correct answers and unexpected approaches in candidates' scripts must be given marks that fairly reflect the relevant knowledge and skills demonstrated.

Mark schemes should be read in conjunction with the published question papers and the report on the examination.

OCR will not enter into any discussion or correspondence in connection with this mark scheme.

© OCR 2013

(Question		Answer	Marks	G	uidance
1	(i)		vectors in plane: two of $\begin{pmatrix} -4\\4\\1 \end{pmatrix}$, $\begin{pmatrix} 0\\6\\4 \end{pmatrix} = 2 \begin{pmatrix} 0\\3\\2 \end{pmatrix}$, $\begin{pmatrix} 4\\2\\3 \end{pmatrix}$	M1	Differences between two pairs	Any multiple
			$\mathbf{r} = \begin{pmatrix} 1\\6\\2 \end{pmatrix} + \lambda \begin{pmatrix} 0\\3\\2 \end{pmatrix} + \mu \begin{pmatrix} 4\\2\\3 \end{pmatrix}$	A1	Aef of parametric equation	Must have " r ="
1	(ii)		(0) (4) (5)			
			$\begin{bmatrix} 0\\3\\2 \end{bmatrix} \times \begin{bmatrix} 2\\3 \end{bmatrix} = \begin{bmatrix} 8\\-12 \end{bmatrix}$	M1 A1	Calculate vector product or multiple	M1 can be awarded where vector product has method shown or only one term wrong
			$\left(\mathbf{r} - \begin{pmatrix} 1\\6\\2 \end{pmatrix} \right) \cdot \begin{pmatrix} 5\\8\\-12 \end{pmatrix} = 0$	M1		Or Cartesian form = d with attempt to compute d
			5x + 8y - 12z = 29	A1	Aef of cartesian equation, isw.	
				[4]		
			Alternate method			
				M1 A1 M1A1	EITHER x, y, z in parametric form both parameters in terms of e.g. x, y substitute into parametric form of z	
				M1 A1 M1 A1	OR <i>x, y, z</i> in parametric form 2 equations in <i>x, y, z</i> and one parameter eliminate parameter	

(Juestior	Answer	Marks	G	uidance
2	(i)	$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$	B2	-1 each error	
		From table clearly closed	B1		Must be clear they are referring to tabulated results
		1 is identity	B1		
		$3^{-1} \equiv 3, 5^{-1} \equiv 5, 7^{-1} \equiv 7 \pmod{8}$	B1		Or "1 appears in every row"
			[5]	Superfluous fact/s gets -1	
2	(ii)	1 has order 1 and 3, 5, 7 all have order 2	B1		
			[1]		
2	(iii)	$\{1, 3\}, \{1, 5\}, \{1, 7\} \text{ (and } \{1\})$	B1	All correct, no extras	Allow {1} included or omitted
			[1]		
2	(iv)	in $H 3^2 \equiv 9 \pmod{10}$ so 3 not order 2	M1	Shows and states that 3 or that 7 is not order 2 (or is order 4)	
		no element of order > 2 in <i>G</i> so not isomorphic	A1	Completely correct reasoning	
			[2]	Or, if zero, then SC1 for merely stating comparable orders and then saying that "orders don't correspond, so not isomorphic" Or table for H with saying "not all elements self inverse, so not isomorphic"	

Question	Answer	Marks	G	uidance
3	$u = y^3 \Longrightarrow \frac{\mathrm{d}u}{\mathrm{d}x} = 3y^2 \frac{\mathrm{d}y}{\mathrm{d}x}$	M1		Or $\frac{\mathrm{d}y}{\mathrm{d}x} = \frac{1}{3}u^{-\frac{2}{3}}\frac{\mathrm{d}u}{\mathrm{d}x}$
	in DE gives $x \frac{\mathrm{d}u}{\mathrm{d}x} + 2u = \frac{\cos x}{x}$	A1		
	$\frac{\mathrm{d}u}{\mathrm{d}x} + \frac{2}{x}u = \frac{\cos x}{x^2}$	B1	Divide	Both sides
	$I = \exp\left(\int \frac{2}{x} dx\right) = e^{2\ln x}$	M1	Correctly integrates	Must have form $\frac{du}{dx} + f(x)u = g(x)$
	$=x^2$	A1		Can be implied by subsequent work
	$x^2 \frac{\mathrm{d}u}{\mathrm{d}x} + 2xu = \cos x$			
	$\frac{\mathrm{d}}{\mathrm{d}x}\left(x^2u\right) = \cos x$			
	$x^2 u = \sin x (+A)$	M1	Integrate	
	$u = \frac{\sin x + A}{x^2}$	A1	Or gives GS in implicit form	Must include constant at this stage
	$y = \left(\frac{\sin x + A}{x^2}\right)^{\frac{1}{3}}$	A1		
		[8]		

(Question		Answer	Marks	G	uidance
4	(i)		Sketch $OA = 3 = 3, OB = 3e^{\frac{1}{3}\pi i} = 3$	B1		Must have axes, A shown 3 across and either scale (or co-ordinates) with B in rough position, or angle and distance on argand diagram. No inconsistencies
			and $\angle BOA = \frac{1}{3}\pi$ hence $\triangle OAB$ equilateral	M1 A1 [3]	Can be seen on diagram	Alt. Attempts AB or second angle
4	(ii)		$3e^{-\frac{1}{3}\pi i}$	M1A1	Or $3e^{\frac{5}{3}\pi i}$. Isw M1 for evidence they are considering BA, or for $\frac{3}{2} - \frac{3}{2}\sqrt{3}i$	For full marks can use CiS form, or clear polar co-ordinates, in radians. Not C-iS
4	(iii)		$\left(3 - 3e^{\frac{1}{3}\pi i}\right)^5 = 3^5 e^{-\frac{5}{3}\pi i}$ $= 243\left(\cos\frac{5}{3}\pi - i\sin\frac{5}{3}\pi\right)$ $= \frac{243}{2} + \frac{243}{2}\sqrt{3}i$	M1 A1ft B1	For mod ⁵ and arg \times 5 aef	"Hence" so must use 'their $3e^{-\frac{1}{3}\pi i}$, Condone use of "121.5".
				[2]		

Question		Answer	Marks	Guidance		
5		AE: $\lambda^2 + 2\lambda + 5 = 0$	M1			
		$\lambda = -1 \pm 2i$	A1			
		CF: $e^{-x} (A\cos 2x + B\sin 2x)$	A1ft		Or $Ae^{-x}\cos(2x+\alpha)$ Must be in real form	
		PI: $y = a e^{-x}$	B1		If PI $y = ax e^{-x}$, then max of M1,A1,A1, B0,M1,A0,A0 (since cannot be consistent) M1, M1, A1.	
		$a e^{-x} - 2a e^{-x} + 5a e^{-x} = e^{-x}$ 4a = 1	M1	Differentiate & substitute	Must have a constant in "their PI"	
		$a = \frac{1}{4}$	A1			
		GS: $y = e^{-x} \left(\frac{1}{4} + A \cos 2x + B \sin 2x \right)$	A1ft		Must have " <i>y</i> ="	
		$\frac{\mathrm{d}y}{\mathrm{d}x} = -\mathrm{e}^{-x} \left(\frac{1}{4} + A\cos 2x + B\sin 2x \right)$ $+ \mathrm{e}^{-x} \left(-2A\sin 2x + 2B\cos 2x \right)$	M1*	Differentiate their GS of form $y = e^{-x} (P + A\cos nx + B\sin nx)$ where P is constant or linear term, n not 0 or 1	Allow one error	
		$x = 0, \frac{dy}{dx} = 0 \Longrightarrow -\left(\frac{1}{4} + A\right) + 2B = 0$ $x = 0, y = 0 \Longrightarrow \frac{1}{4} + A = 0$	*M1	Use conditions	But M0 if leads to solution of $y = 0$	
		$A = -\frac{1}{4}, B = 0$	A1ft	From their GS		
		$y = \frac{1}{4} \mathrm{e}^{-x} \left(1 - \cos 2x \right)$	A1 [11]		Must have ' $y =$ ' and be in real form	
6	(i)	x = 2t + 1, y = 5t - 1, z = t + 2	B1	Parameterise	or B1 for y and z correctly in terms of x e.g. $2y = 5x - 7$, $2z = x + 3$	
		(2t+1)+2(5t-1)-2(t+2)=5		Substitute into plane	Then M1 for full simultaneous equations method.	
		$\Rightarrow 10t = 10 \Rightarrow t = 1$ Intersect at (3, 4, 3)	M1 A1 [3]	Solve cao	Accept vector form	

(Questi	on	Answer	Marks	Guidance		
6	(ii)		$\cos\left(\frac{1}{2}\pi - \theta\right) = \frac{\begin{vmatrix} 2\\5\\1 \end{vmatrix} \begin{pmatrix} 1\\2\\-2 \end{vmatrix}}{\begin{vmatrix} 2\\-2 \end{vmatrix}} = \frac{10}{3\sqrt{30}}$	M1A1		Attempt to find angle or its complement	
			$\theta = 0.654$	A1	or 37.5°		
6	(iii)		If P is point of intersection and Q is required point, $\overrightarrow{PQ} = \lambda \begin{pmatrix} 2 \\ 5 \\ 1 \end{pmatrix}$ so $\sin \theta = \frac{2}{PQ} = \frac{2}{ \lambda \sqrt{30}}$	 M1*	or $\overrightarrow{PQ} \cdot \hat{\mathbf{n}} = \pm 2$ where $\mathbf{n} = \begin{pmatrix} 1 \\ 2 \\ -2 \end{pmatrix}$	Use \overrightarrow{PQ} with right angled triangle or consider component of \overrightarrow{PQ} in direction of normal vector.	
			$\frac{10}{3\sqrt{30}} = \frac{2}{ \lambda \sqrt{30}}$	M1		Valid method to set up equation in λ alone.	
			$\lambda = \pm \frac{3}{5}$	A1		(May work from general point on original equation)	
			points have position vectors $\begin{pmatrix} 3\\4\\3 \end{pmatrix} \pm \frac{3}{5} \begin{pmatrix} 2\\5\\1 \end{pmatrix}$	*M1	Dep on 1 st M1		
			points at (1.8, 1, 2.4) and (4.2, 7, 3.6)	A1	сао		
			Alternative:				
			Distance = $\frac{ 2t+1+2(5t-1)-2(t+2)-5 }{\sqrt{1^2+2^2+2^2}} = 2$	M1* A1		Zero if formula used without substitution in of parametric form.	
			$\Rightarrow t = 0.4 \text{ or } 1.6$ (1.8, 1, 2.4) and (4.2, 7, 3.6)	*M1 A1 A1 [5]	Solve At least one value found		

(Question		Answer	Marks	Guidance		
7	(i)		$(ab)^6 = ababab = a^6b^6$ as commutative	M1	Must give reason	Some demonstration that they understand commutativity	
			$=(a^2)^3(b^3)^2 = e^3e^2 = e^3e^2$	A1	Using orders of a and b		
			So <i>ab</i> has order 1, 2, 3, or 6				
			$(b \neq a \Rightarrow ab \neq a^2 \Rightarrow ab \neq e \text{ so } ab \text{ not order } 1)$			Condone absence of this line	
			$(ab)^2 = a^2b^2 = eb^2 = b^2$ and b not order 2, so ab not order 2	M1	Consider other cases	Insufficient to merely have simplified all $(ab)^n$	
			$(ab)^3 = a^3b^3 = aa^2e = aee = a \neq e$, so ab not order 3				
			(So must be order 6)	A1 [4]	AG Complete argument		
7	(ii)		ac has order 18	B1		Or <i>abc</i> or generator	
			18 is LCM of 2 and 9, (so we can use a similar argument to part (i))	M1	or explicit consideration of other cases		
			So as G has an element of order 18 it must be cyclic.	A1	AG Complete argument		
				[3]			
8	(i)		$\cos 5\theta + i \sin 5\theta = (\cos \theta + i \sin \theta)^5$	B 1	Or $\cos 5\theta = re\{(\cos \theta + i \sin \theta)^5\}$		
			$=c^{5}+5ic^{4}s-10c^{3}s^{2}-10ic^{2}s^{3}+5cs^{4}+is^{5}$	M1		No more than 1 error, can be unsimplified	
			$\cos 5\theta = c^5 - 10c^3s^2 + 5cs^4$	M1	Take real parts		
			$= c^{5} - 10c^{3}(1 - c^{2}) + 5c(1 - c^{2})^{2}$	M1			
			$= c^{5} - 10c^{3} + 10c^{5} + 5c - 10c^{3} + 5c^{5}$				
			$\cos 5\theta = 16c^5 - 20c^3 + 5c$	A1	AG		
			1	[ວ]			

(Question		Answer	Marks	Guidance	
8	(ii)		Multiplying by x gives $16x^5 - 20x^3 + 5x = 0$			Hence, so no marks for using quadratic at this stage.
			letting $x = \cos \alpha$ gives $\cos 5\alpha = 0$	M1		
			hence $5\alpha = \frac{1}{2}\pi, \frac{3}{2}\pi, \frac{5}{2}\pi, \frac{7}{2}\pi, \frac{9}{2}\pi$	A1		
			$\alpha = \frac{1}{10}\pi, \frac{3}{10}\pi, \frac{5}{10}\pi, \frac{7}{10}\pi, \frac{9}{10}\pi$			
			$\cos\frac{5}{10}\pi = 0$ which is not a root	A1		
			so roots $x = \cos \frac{1}{10}\pi$, $\cos \frac{3}{10}\pi$, $\cos \frac{7}{10}\pi$, $\cos \frac{9}{10}\pi$	A1		
				[4]		
8	(iii)		$16x^4 - 20x^2 + 5 = 0 \Leftrightarrow x^2 = \frac{20 \pm \sqrt{80}}{32}$	B1		Can be gained if seen in (ii)
			cos decreases between 0 and π so $\cos \frac{1}{10}\pi$ is			
			greatest root	M1		
			so $\cos\frac{1}{10}\pi = \sqrt{\frac{20 + \sqrt{80}}{32}} = \sqrt{\frac{5 + \sqrt{5}}{8}}$	A1	Dep on full marks in (ii)	
				[3]		

OCR (Oxford Cambridge and RSA Examinations) 1 Hills Road Cambridge CB1 2EU

OCR Customer Contact Centre

Education and Learning

Telephone: 01223 553998 Facsimile: 01223 552627 Email: general.qualifications@ocr.org.uk

www.ocr.org.uk

For staff training purposes and as part of our quality assurance programme your call may be recorded or monitored