

OXFORD CAMBRIDGE AND RSA EXAMINATIONS

**Advanced Subsidiary General Certificate of Education
Advanced General Certificate of Education**

MATHEMATICS

4725

Further Pure Mathematics 1

Specimen Paper

Additional materials:
Answer booklet
Graph paper
List of Formulae (MF 1)

TIME 1 hour 30 minutes

INSTRUCTIONS TO CANDIDATES

- Write your Name, Centre Number and Candidate Number in the spaces provided on the answer booklet.
- Answer **all** the questions.
- Give non-exact numerical answers correct to 3 significant figures, unless a different degree of accuracy is specified in the question or is clearly appropriate.
- You are permitted to use a graphic calculator in this paper.

INFORMATION FOR CANDIDATES

- The number of marks is given in brackets [] at the end of each question or part question.
- The total number of marks for this paper is 72.
- Questions carrying smaller numbers of marks are printed earlier in the paper, and questions carrying larger numbers of marks later in the paper.
- **You are reminded of the need for clear presentation in your answers.**

This question paper consists of 4 printed pages.

- 1 Use formulae for $\sum_{r=1}^n r$ and $\sum_{r=1}^n r^2$ to show that

$$\sum_{r=1}^n r(r+1) = \frac{1}{3}n(n+1)(n+2). \quad [5]$$

- 2 The cubic equation $x^3 - 6x^2 + kx + 10 = 0$ has roots $p - q$, p and $p + q$, where q is positive.

(i) By considering the sum of the roots, find p . [2]

(ii) Hence, by considering the product of the roots, find q . [3]

(iii) Find the value of k . [3]

- 3 The complex number $2 + i$ is denoted by z , and the complex conjugate of z is denoted by z^* .

(i) Express z^2 in the form $x + iy$, where x and y are real, showing clearly how you obtain your answer. [2]

(ii) Show that $4z - z^2$ simplifies to a real number, and verify that this real number is equal to zz^* . [3]

(iii) Express $\frac{z+1}{z-1}$ in the form $x + iy$, where x and y are real, showing clearly how you obtain your answer. [3]

- 4 A sequence u_1, u_2, u_3, \dots is defined by

$$u_n = 3^{2n} - 1.$$

(i) Write down the value of u_1 . [1]

(ii) Show that $u_{n+1} - u_n = 8 \times 3^{2n}$. [3]

(iii) Hence prove by induction that each term of the sequence is a multiple of 8. [4]

- 5 (i) Show that

$$\frac{1}{2r-1} - \frac{1}{2r+1} = \frac{2}{4r^2-1}. \quad [2]$$

- (ii) Hence find an expression in terms of n for

$$\frac{2}{3} + \frac{2}{15} + \frac{2}{35} + \dots + \frac{2}{4n^2-1}. \quad [4]$$

- (iii) State the value of

(a) $\sum_{r=1}^{\infty} \frac{2}{4r^2-1}, \quad [1]$

(b) $\sum_{r=n+1}^{\infty} \frac{2}{4r^2-1}. \quad [1]$

- 6 In an Argand diagram, the variable point P represents the complex number $z = x + iy$, and the fixed point A represents $a = 4 - 3i$.

- (i) Sketch an Argand diagram showing the position of A , and find $|a|$ and $\arg a$. [4]

- (ii) Given that $|z - a| = |a|$, sketch the locus of P on your Argand diagram. [3]

- (iii) Hence write down the non-zero value of z corresponding to a point on the locus for which

- (a) the real part of z is zero, [1]

- (b) $\arg z = \arg a$. [2]

- 7 The matrix \mathbf{A} is given by $\mathbf{A} = \begin{pmatrix} 1 & -2 \\ 2 & 1 \end{pmatrix}$.

- (i) Draw a diagram showing the unit square and its image under the transformation represented by \mathbf{A} . [3]

- (ii) The value of $\det \mathbf{A}$ is 5. Show clearly how this value relates to your diagram in part (i). [3]

\mathbf{A} represents a sequence of two elementary geometrical transformations, one of which is a rotation R .

- (iii) Determine the angle of R , and describe the other transformation fully. [3]

- (iv) State the matrix that represents R , giving the elements in an exact form. [2]

8 The matrix \mathbf{M} is given by $\mathbf{M} = \begin{pmatrix} a & 2 & -1 \\ 2 & 3 & -1 \\ 2 & -1 & 1 \end{pmatrix}$, where a is a constant.

(i) Show that the determinant of \mathbf{M} is $2a$. [2]

(ii) Given that $a \neq 0$, find the inverse matrix \mathbf{M}^{-1} . [4]

(iii) Hence or otherwise solve the simultaneous equations

$$\begin{aligned} x + 2y - z &= 1, \\ 2x + 3y - z &= 2, \\ 2x - y + z &= 0. \end{aligned} \quad [3]$$

(iv) Find the value of k for which the simultaneous equations

$$\begin{aligned} 2y - z &= k, \\ 2x + 3y - z &= 2, \\ 2x - y + z &= 0, \end{aligned}$$

have solutions. [3]

(v) Do the equations in part (iv), with the value of k found, have a solution for which $x = z$? Justify your answer. [2]